

5-28-2010

Reducing the effects of intracellular accumulation of thermolabile collagen II mutants by increasing their thermostability in cell culture conditions.

Katarzyna Gawron
Thomas Jefferson University


Deborah A. Jensen
Thomas Jefferson University

Andrzej Steplewski
Thomas Jefferson University

Andrzej Fertala
Thomas Jefferson University, andrzej.fertala@jefferson.edu

[Let us know how access to this document benefits you](#)

Follow this and additional works at: <http://jdc.jefferson.edu/dcbfp>

 Part of the [Dermatology Commons](#), and the [Medical Biochemistry Commons](#)

Recommended Citation

Gawron, Katarzyna; Jensen, Deborah A.; Steplewski, Andrzej; and Fertala, Andrzej, "Reducing the effects of intracellular accumulation of thermolabile collagen II mutants by increasing their thermostability in cell culture conditions." (2010). *Department of Dermatology and Cutaneous Biology Faculty Papers*. Paper 9.

<http://jdc.jefferson.edu/dcbfp/9>

As submitted to:

Biochemical and BioPhysical Research Communications

And later published as:

“Reducing the effects of intracellular accumulation of thermolabile collagen II mutants by increasing their thermostability in cell culture conditions”

Volume 396, Issue 2, 28 May 2010, Pages 213-218

doi:10.1016/j.bbrc.2010.04.056

Katarzyna Gawron, Deborah A. Jensen, Andrzej Steplewski and Andrzej Fertala *

Department of Dermatology and Cutaneous Biology, Jefferson Medical College, Thomas

Jefferson University, Philadelphia, PA, 19107, USA

*Correspondence to Andrzej Fertala, Department of Dermatology and Cutaneous Biology, Jefferson Medical College, Thomas Jefferson University, BLSB, Room 424, 233 S. 10th Street, Philadelphia, PA 19107, U.S.A.

Tel: 215-503-0113

Fax: 215-503-5788

E-mail: andrzej.fertala@jefferson.edu

Abstract

Mutations in collagen II are associated with spondyloepiphyseal dysplasia, a group of heritable diseases whose common features include aberrations of skeletal growth. The mechanisms through which mutations in collagen II affect the cartilaginous tissues are complex and include both intracellular and extracellular processes. One of those mechanisms involves cellular stress caused by excessive accumulation of misfolded collagen II mutants. We investigated whether stabilizing the structure of thermolabile R789C and R992C collagen II mutants would improve their secretion from cells, thereby reducing cellular stress and apoptosis. Employing glycerol and trimethylamine N-oxide (TMAO), chemicals that increase the thermostability of collagen triple helices, we demonstrated that those compounds function as chaperones and stabilize the R789C and R992C mutants, accelerate their secretion, and improve cell survival. Our study provides a scientific basis for considering misfolded triple helices of collagen mutants a target for reducing the deleterious effects caused by their excessive intracellular accumulation.

Introduction

Collagen II is a component of heterotypic fibrils which form scaffolds of mature cartilaginous tissues and provide a template that guides the development of skeletal tissues [1]. Processes leading from the biosynthesis of pro- α 1(II) chains in cells to the formation of collagen fibrils in the extracellular space are complex and include: (i) posttranslational modifications of nascent pro- α 1(II) chains, (ii) their folding into triple-helical procollagen molecules and secretion from cells, and (iii) self-assembly of collagen molecules into fibrils [2].

Fundamental for the mechanical and biological functions of collagen II is its stable triple-helical conformation. Such a conformation is achieved by the folding of individual pro- α 1(II) chains whose main collagenous domain consists of uninterrupted repetitions of the G-X-Y motif in which the "X" and the "Y" positions are frequently occupied by proline and hydroxyproline residues, respectively [2]. The accurate amino acid sequence of collagen II is critical for the correct functions of this protein at the tissue level and mutations in *COL2A1* (OMIM#120140) are associated with a broad spectrum of chondrodysplasia phenotypes, generally described as spondyloepiphyseal dysplasia (SED), the main feature of which is abnormal skeletal development [3]

One of the pathological consequences of mutations in collagenous proteins is a loss-of-function effect due to the reduced incorporation of mutants into the extracellular matrix (ECM). It has been suggested that the fundamental cause for the reduced amount of collagen molecules containing mutant α chains is excessive intracellular accumulation and subsequent degradation in a process described as "procollagen suicide" [4].

Due to the possibility of excessive intracellular retention, some diseases that involve collagen mutants are categorized as storage diseases, a group that also includes diseases caused by the excessive accumulation of non-collagenous proteins, lipids, glycoproteins, and other macromolecules [5]. In our studies published elsewhere we demonstrated that the R789C and R992C substitutions in collagen II cause misfolding of mutant molecules, decrease their thermostabilities, promote excessive intracellular retention and aggregation, and increase apoptosis [6; 7; 8]. Thus, in addition to the loss-of-function effect, certain mutations in collagen II may also cause a gain-of-toxicity effect.

Here, we studied the feasibility of reducing cellular stress associated with the R789C and R992C substitutions by stabilizing the mutant molecules and decreasing their intracellular accumulation. Employing glycerol and trimethylamine N-oxide (TMAO), chemicals that stabilize the collagen triple helix, we demonstrated that these compounds increase the thermostability of the R789C and R992C mutants, decrease their intracellular retention, and improve cell survival [9; 10; 11]. Our study provides bases for considering triple helices of thermolabile collagen mutants a new target to reduce the deleterious effects caused by their atypical retention.

Materials and Methods

Mutation nomenclature -- The amino acid substitutions are named according to the literature, with amino acid residues numbered from the first glycine of the collagen II triple helix.

Procollagen II mutants -- Procollagen II mutants employed in this study are associated with SED [12; 13]. To facilitate various biochemical and microscopic assays, wild type (WT) and mutant procollagen II variants were engineered as chimeras fused with green fluorescent protein (Pro-GFP), as described [6; 8].

Cells expressing collagen II mutants -- For the production of Pro-GFP variants in quantities sufficient for assays of their thermostability, these proteins were expressed in HT-1080 cells (ATCC; CCL-121), as described [6; 8; 14]. For analyses of cellular responses to the presence of procollagen II mutants, the chondrocytic cell line SW-1353 (ATCC; HTB-94) was employed, as described [6; 7; 8]. Both cell lines were cultured in Dulbecco's Modified Eagle Medium supplemented with 10% fetal bovine serum and 40 µg/ml of L-ascorbic acid phosphate magnesium salt (Wako Chemicals, Inc.), as described [8; 14].

Cell culture in the presence of glycerol and TMAO -- Glycerol and TMAO, compounds with chemical chaperone characteristics, were employed to determine the effects of their collagen-stabilizing activities on the structure of the R789C and R992C mutants, and the faith of cells harboring them. For

analyses of cell survival, we followed the experimental design described by Hintze *et al* [7]. In brief, SW-1353 cells were cultured for 12 days post-confluence, and then cell culture media were supplemented with glycerol added to a final concentration of 0.1 M or 0.5 M, or TMAO added at 30 mM or 0.1 M. Subsequently, cell culture was continued for five additional days. After the five-day treatment, cell culture media and cell lysates were collected for further assays. For thermostability analyses, collagen II variants were isolated from HT-1080 cells cultured in the presence of 0.5 M glycerol or 0.1 M TMAO, as described [14].

Working concentrations of selected compounds used in this study were determined in pilot experiments. In these experiments cells were cultured in increasing concentrations of glycerol or TMAO. During culture, cells were monitored for proliferation, changes in morphology, and production of procollagen II, as described [6; 8]. The highest concentrations that did not have any apparent negative effects on cultured cells were applied in the presented studies.

Cleaved poly-(ADP-ribose) polymerase (cPARP) and TdT-mediated dUTP Nick-End Labeling (TUNEL) assays -- Glycerol and TMAO-treated cells were analyzed for apoptotic markers, as described [7]. In these assays cPARP was detected in cell lysates by Western blot with primary rabbit anti-cPARP antibodies (Cell Signaling Technology, Inc.) and secondary anti-rabbit IgG conjugated to horseradish peroxidase (HRP; Sigma-Aldrich). The same nitrocellulose membranes were also probed for glyceraldehyde-3-phosphate dehydrogenase (GAPDH). The relative amount of cPARP in the analyzed samples was determined by measuring the ratios of the pixel intensities of the cPARP-positive bands and GAPDH-positive bands. In any given group of cells, *i.e.* expressing GFP-tagged WT, R789C or R992C collagen II, the cPARP/GAPDH ratio derived from non-treated cells was considered 100%. Statistical significances of differences between the means calculated for the relative contents of cPARP in cells cultured in the presence vs. the absence of tested compounds were evaluated by the Student's *t* test (GraphPad Software, Inc.).

In another set of experiments the presence of fragmented DNA was analyzed by TUNEL assays that utilize tetramethylrodamine (TMR) as a red fluorophore (Roche). Subsequently the apoptotic index,

defined as the percentage of TUNEL-positive nuclei out of the total number of DAPI-positive nuclei, was calculated for each analyzed group, as described [6; 7].

Thermostability assays of collagen II variants -- Two experimental approaches were designed; (i) in the first approach we analyzed the thermostabilities of purified collagen II mutants produced by HT-1080 cells cultured in the presence of 0.5 M glycerol or 0.1 M TMAO, and (ii) in the second approach these compounds were added to purified collagen II variants produced by cells cultured in the absence of these compounds. The thermostabilities were measured by brief protease digestion, as described [6; 15]. Based on collagen II-melting curves, we determined the temperatures at which the triple helical structure of analyzed collagens was preserved in 75% ($T_{m_{75}}$), 50% ($T_{m_{50}}$), and 25% ($T_{m_{25}}$) of maximum.

Intracellular Pro-GFP variants produced in the presence of glycerol and TMAO -- We analyzed whether glycerol and TMAO-mediated stabilization of collagen molecules could also influence their intracellular accumulation. In brief, glycerol or TMAO-treated cells were lysed, and then Pro-GFP variants were detected by Western blot assays with an anti-GFP antibody, as described [6; 8]. The relative content of GFP-positive bands was assayed by densitometry. The results of measurements were normalized to GAPDH and plotted as a percent of the non-treated control (expressed as 100%).

We also studied the effects of glycerol and TMAO on the accumulation of collagen molecules misfolded due to inhibition of critical hydroxylation of proline residues by 2,2' dipyridyl (Sigma-Aldrich) [16]. The intracellular content of Pro-GFP variants produced in the presence of 2,2' dipyridyl was assayed as described above.

Results

Apoptotic markers in cells harboring the R789C and R992C mutants cultured in the presence of glycerol and TMAO -- Cells cultured in the presence of 0.5 M glycerol or 0.1 M TMAO had a statistically significant decrease in the relative amount of cPARP, indicating an increase of cell survival (Fig. 1 and Supplementary Fig. S1). In the same experimental conditions, the overall amounts of Pro-GFP

produced by cells did not change (Fig. 1). The lower concentrations of these compounds used in the pilot studies did not have significant effects on relative amounts of cPARP (data not shown).

TUNEL analyses (Supplementary Fig. S2) of cells cultured in the presence of 0.5 M glycerol or 0.1 M TMAO support the results of cPARP assays. Specifically, we determined that in comparison to the non-treated cells harboring the WT, R789C or the R992C mutants, the apoptotic indexes for cells cultured in the presence of 0.5 M glycerol decreased from 0.89% (± 0.15 SEM), 3.45% (± 0.73 SEM), and 0.96% (± 0.12 SEM), to 0.82% (± 0.14 SEM), 2.34% (± 0.3 SEM), and 0.5% (± 0.11 SEM), respectively. A similar trend was observed in an analogous experiment where cells were treated with 0.1 M TMAO. Specifically, we determined that in comparison to the non-treated cells harboring the WT, R789C or the R992C mutants (see above) the apoptotic indexes for cells cultured in the presence of 0.1 M TMAO decreased to 0.56% (± 0.14 SEM), 2.24% (± 0.22 SEM), and 0.67% (± 0.08 SEM), respectively.

Thermostabilities of the R789C and R992C mutants produced by cells cultured in the presence of glycerol or TMAO -- In comparison to the non-treated controls, the thermostabilities of the WT collagen II and the analyzed mutants produced by cells cultured in the presence of 0.5 M glycerol or 0.1 M TMAO have increased (Fig. 2A, Supplementary Fig. 3S, and Tab. 1). Based on this observation, however, it has not been clear whether the thermostabilizing effect depended on intracellular interactions of glycerol and TMAO with newly synthesized pro- $\alpha 1(\text{II})$ chains or whether these compounds were able exert a stabilizing effect extracellularly. Thus, in another set of experiments, we first prepared collagen II variants isolated from cells cultured in the absence of chaperones. Next, purified collagens were exposed to these compounds for 24-h at 4°C, and then thermostability measurements were performed. Because the thermostabilities of analyzed mutants have increased, these assays suggest that intracellular interaction between the chemical chaperones and procollagen is not a defining factor in stabilizing the collagen II variants (Fig. 2B, Supplementary Fig. 3S, and Tab. 1). When comparing the thermostabilities of the collagen variants purified from cultures of cells grown in the presence of chaperones to those isolated from cells grown in chaperone-free media and treated with chaperones right before the stability assays, it is apparent that the increase in thermostability was greater in the latter group. (Fig. 2 and Tab. 1).

Note: As glycerol and TMAO do not change the enzymatic activities of trypsin and chymotrypsin, it is important to emphasize that the observed increase of thermostabilities of collagen II variants was not a result of the decrease of the enzymatic activities of employed proteases [17].

Intracellular accumulation of the R789C and R992C mutants -- We analyzed the intracellular content of Pro-GFP variants in the presence of glycerol and TMAO. In comparison to the non-treated controls, in cells treated with these compounds the relative amounts of the R789C and R992C variants were reduced (Fig. 3 and Fig. 4). Of note was the observation that in cells expressing the R789C mutant, partial degradation of this protein was observed.

As shown in Fig. 3 and Fig. 4 adding 2,2' dipyridyl into the cell culture increased accumulation of Pro-GFP in all analyzed groups. Moreover, as evident by the presence of GFP-positive fragments, increased intracellular accumulation was associated with degradation of all Pro-GFP variants. In addition, in cells cultured in the presence of this compound, neither glycerol nor TMAO reduced the intracellular accumulation of the Pro-GFP variants (Fig. 3 and Fig. 4).

Discussion

Employing a simple experimental system, we tested whether adverse consequences of intracellular accumulation of the thermolabile R789C and R992C collagen II mutants could be partially reduced by improving their thermostabilities. The strategy proposed here was to employ glycerol and TMAO, compounds that at 1 M concentration increase the thermostability of collagen triple helices by 1°C and 5°C, respectively [9; 10].

We demonstrated that the presence of glycerol and TMAO was associated with reduction of the relative amounts of cPARP in cells harboring the R789C and R992C mutants. At the same time, the presence of these compounds did not affect the overall production of collagen II. In cells expressing WT collagen II, the relative amount of background-level cPARP also decreased, but this change was relatively small compared to that observed for cells harboring collagen II mutants. The decrease in the apoptotic indexes for cells expressing the R789C and R992C variants in the presence of glycerol and TMAO further corroborates the results of quantitative assays of cPARP. Changes in the relative amounts of apoptotic markers in cells cultured in the presence of glycerol or TMAO correlated with the decrease of the

intracellular pool of procollagen molecules, suggesting the increase of their secretion. Because the efficient secretion of procollagens depends on the formation of thermostable triple-helical structures, the decrease of intracellular accumulation of inherently thermolabile R789C and R992C procollagen II mutants in the presence of glycerol and TMAO further indicate the chaperone-like collagen-stabilizing effect of these compounds [6; 7; 16; 18; 19]. The collagen-stabilizing activity of selected chemical chaperones, however, is probably limited to mutant molecules with localized structural aberrations. This notion is based on the observation that collagen molecules whose structures are extensively misfolded due to the inhibition of hydroxylation of proline residues by 2,2' dipyridyl are not stabilized by selected chemical chaperones..

The decrease in the intracellular accumulation of studied mutants indicates that the stabilizing effects of glycerol and TMAO were primarily a result of the intracellular influence of these compounds on the structure of a collagen triple helix. In comparison to the stabilities of collagen II variants purified from cell culture media of cells grown in the presence of glycerol or TMAO, the increase in the thermostability was even greater in another experimental group in which these compounds were added to the purified-collagen samples before thermostability assays and remained in the analyzed samples during these assays. This apparent difference was most likely due to the partial reversal of the stabilizing effects in the first experimental group due to the decrease in the concentration of glycerol and TMAO occurring during the purification of collagen II.

The concept of improving the structure of misfolded mutant proteins by employing chemical chaperones to reduce the harmful effects that such mutants have on cells and tissues has been proposed and tested with a number of proteins associated with various diseases [20; 21; 22; 23; 24]. Specific examples of such an approach include employing TMAO to reduce the effects of mutant keratins that cause various skin disorders and to restore the activity of mutant branched-chain α -ketoacid dehydrogenase associated with maple syrup urine disease [25; 26]. Moreover, chemical chaperones were shown to be effective in stabilizing the structure and accelerating the secretion of mutant proteins involved in cystic fibrosis and Huntington's disease, as well as liver injury and emphysema caused by an α 1-antitrypsin deficiency [27; 28; 29]. Although most studies on chemical chaperones as therapeutic agents were done in cell culture conditions, reduction of ER stress and restoration of glucose homeostasis

by administration of 4-phenylbutyric acid, yet another compound with chaperone-like activities, in a mouse model of type 2 diabetes was also demonstrated, thereby indicating a potential utility of chemical chaperones *in vivo* [30].

The studies presented here provide experimental evidence that the aberrant structure of collagen mutants could be considered a target for approaches aimed at reducing the harmful effects imposed by these mutants. Preliminary tests presented here focused on the thermolabile R789C and R992C collagen II mutants that are associated with apoptosis and whose structural alterations are readily detectable [6; 7; 8; 31]. The structural and biological effects of these mutations, however, do not represent the full spectrum of potential effects that different mutations may impose. For instance, in cells expressing the thermostable R75C, R519C, and G853E collagen II mutants which are also associated with skeletal diseases, an increase in apoptosis was not readily apparent [7]. Thus, the question remains whether targeting the structure of collagens should be considered only for the thermolabile mutants or also for the thermostable mutants whose ER stress-related effects may be long-term and accumulative.

Acknowledgements

This work was supported by the National Health Institutes [5R01AR049537-06].

References

- [1] E. Gustafsson, A. Aszodi, N. Ortega, E.B. Hunziker, H.W. Denker, Z. Werb, and R. Fassler, Role of collagen type II and perlecan in skeletal development. *Ann N Y Acad Sci* 995 (2003) 140-50.
- [2] D.J. Prockop, and K.I. Kivirikko, Collagens: molecular biology, diseases, and potentials for therapy. *Annu Rev Biochem* 64 (1995) 403-34.
- [3] B.R. Olsen, Mutations in collagen genes resulting in metaphyseal and epiphyseal dysplasias. *Bone* 17 (1995) 45S-49S.
- [4] C.J. Williams, and D.J. Prockop, Synthesis and processing of a type I procollagen containing shortened pro-alpha 1(I) chains by fibroblasts from a patient with osteogenesis imperfecta. *J Biol Chem* 258 (1983) 5915-21.

- [5] J. Rutishauser, and M. Spiess, Endoplasmic reticulum storage diseases. *Swiss Med Wkly* 132 (2002) 211-22.
- [6] H.J. Chung, D.A. Jensen, K. Gawron, A. Steplewski, and A. Fertala, R992C (p.R1192C) Substitution in collagen II alters the structure of mutant molecules and induces the unfolded protein response. *J Mol Biol* 390 (2009) 306-18.
- [7] V. Hintze, A. Steplewski, H. Ito, D.A. Jensen, U. Rodeck, and A. Fertala, Cells expressing partially unfolded R789C/p.R989C type II procollagen mutant associated with spondyloepiphyseal dysplasia undergo apoptosis. *Hum Mutat* 29 (2008) 841-51.
- [8] H. Ito, E. Rucker, A. Steplewski, E. McAdams, R.J. Brittingham, T. Alabyeva, and A. Fertala, Guilty by association: some collagen II mutants alter the formation of ECM as a result of atypical interaction with fibronectin. *J Mol Biol* 352 (2005) 382-95.
- [9] K. Beck, V.C. Chan, N. Shenoy, A. Kirkpatrick, J.A. Ramshaw, and B. Brodsky, Destabilization of osteogenesis imperfecta collagen-like model peptides correlates with the identity of the residue replacing glycine. *Proc Natl Acad Sci U S A* 97 (2000) 4273-8.
- [10] R. Penkova, I. Goshev, S. Gorinstein, and P. Nedkov, Stability of collagen during denaturation. *J Protein Chem* 18 (1999) 397-401.
- [11] A.E. Russell, Effect of alcohols and neutral salt on the thermal stability of soluble and precipitated acid-soluble collagen. *Biochem J* 131 (1973) 335-42.
- [12] D. Chan, T.K. Taylor, and W.G. Cole, Characterization of an arginine 789 to cysteine substitution in alpha 1 (II) collagen chains of a patient with spondyloepiphyseal dysplasia. *J Biol Chem* 268 (1993) 15238-45.
- [13] L.R. Donahue, B. Chang, S. Mohan, N. Miyakoshi, J.E. Wergedal, D.J. Baylink, N.L. Hawes, C.J. Rosen, P. Ward-Bailey, Q.Y. Zheng, R.T. Bronson, K.R. Johnson, and M.T. Davisson, A missense mutation in the mouse *Col2a1* gene causes spondyloepiphyseal dysplasia congenita, hearing loss, and retinoschisis. *J Bone Miner Res* 18 (2003) 1612-21.
- [14] A. Fertala, A.L. Sieron, A. Ganguly, S.W. Li, L. Ala-Kokko, K.R. Anumula, and D.J. Prockop, Synthesis of recombinant human procollagen II in a stably transfected tumour cell line (HT1080). *Biochem J* 298 (1994) 31-7.

- [15] A.L. Sieron, A. Fertala, L. Ala-Kokko, and D.J. Prockop, Deletion of a large domain in recombinant human procollagen II does not alter the thermal stability of the triple helix. *J Biol Chem* 268 (1993) 21232-7.
- [16] W.W. Kao, D.J. Prockop, and R.A. Berg, Kinetics for the secretion of nonhelical procollagen by freshly isolated tendon cells. *J Biol Chem* 254 (1979) 2234-43.
- [17] R. Kumar, J.M. Serrette, and E.B. Thompson, Osmolyte-induced folding enhances tryptic enzyme activity. *Arch Biochem Biophys* 436 (2005) 78-82.
- [18] P. Bornstein, The biosynthesis of collagen. *Annu Rev Biochem* 43 (1974) 567-603.
- [19] G.S. Barsh, and P.H. Byers, Reduced secretion of structurally abnormal type I procollagen in a form of osteogenesis imperfecta. *Proc Natl Acad Sci U S A* 78 (1981) 5142-6.
- [20] I. Kim, W. Xu, and J.C. Reed, Cell death and endoplasmic reticulum stress: disease relevance and therapeutic opportunities. *Nat Rev Drug Discov* 7 (2008) 1013-30.
- [21] F.E. Cohen, and J.W. Kelly, Therapeutic approaches to protein-misfolding diseases. *Nature* 426 (2003) 905-9.
- [22] D.H. Perlmutter, Chemical chaperones: a pharmacological strategy for disorders of protein folding and trafficking. *Pediatr Res* 52 (2002) 832-6.
- [23] I. Baskakov, and D.W. Bolen, Forcing thermodynamically unfolded proteins to fold. *J Biol Chem* 273 (1998) 4831-4.
- [24] T. Arakawa, D. Ejima, Y. Kita, and K. Tsumoto, Small molecule pharmacological chaperones: From thermodynamic stabilization to pharmaceutical drugs. *Biochim Biophys Acta* 1764 (2006) 1677-87.
- [25] D. Lee, D. Santos, H. Al-Rawi, A.M. McNeill, and E.L. Rugg, The chemical chaperone trimethylamine N-oxide ameliorates the effects of mutant keratins in cultured cells. *Br J Dermatol* 159 (2008) 252-5.
- [26] J.L. Song, and D.T. Chuang, Natural osmolyte trimethylamine N-oxide corrects assembly defects of mutant branched-chain alpha-ketoacid decarboxylase in maple syrup urine disease. *J Biol Chem* 276 (2001) 40241-6.

- [27] C.R. Brown, L.Q. Hong-Brown, and W.J. Welch, Strategies for correcting the delta F508 CFTR protein-folding defect. *J Bioenerg Biomembr* 29 (1997) 491-502.
- [28] J.A. Burrows, L.K. Willis, and D.H. Perlmutter, Chemical chaperones mediate increased secretion of mutant alpha 1-antitrypsin (alpha 1-AT) Z: A potential pharmacological strategy for prevention of liver injury and emphysema in alpha 1-AT deficiency. *Proc Natl Acad Sci U S A* 97 (2000) 1796-801.
- [29] S. Reijonen, N. Putkonen, A. Norremolle, D. Lindholm, and L. Korhonen, Inhibition of endoplasmic reticulum stress counteracts neuronal cell death and protein aggregation caused by N-terminal mutant huntingtin proteins. *Exp Cell Res* 314 (2008) 950-60.
- [30] U. Ozcan, E. Yilmaz, L. Ozcan, M. Furuhashi, E. Vaillancourt, R.O. Smith, C.Z. Gorgun, and G.S. Hotamisligil, Chemical chaperones reduce ER stress and restore glucose homeostasis in a mouse model of type 2 diabetes. *Science* 313 (2006) 1137-40.
- [31] A. Steplewski, I. Majsterek, E. McAdams, E. Rucker, R.J. Brittingham, H. Ito, K. Hirai, E. Adachi, S.A. Jimenez, and A. Fertala, Thermostability gradient in the collagen triple helix reveals its multi-domain structure. *J Mol Biol* 338 (2004) 989-98.

Table 1

Thermostabilities of the R789C and R992C mutant in the presence of glycerol and TMAO.

	Collagen II variant	^a Tm ₇₅ [°C]			^a Tm ₅₀ [°C]			^a Tm ₂₅ [°C]		
		C	0.5M Glycerol	0.1M TMAO	C	0.5 M Glycerol	0.1M TMAO	C	0.5M Glycerol	0.1M TMAO
Chaperones added during cell culture	WT	40.4	40.6	40.5	41.3	41.5	41.3	42.2	42.3	42.2
	R789C	37.7	37.8	37.9	38.6	38.6	38.7	39.6	39.5	39.6
	R992C	38.7	39.4	39.1	39.3	39.8	39.6	40.1	40.1	40.3
Chaperones added to purified collagen isolated from non-treated cells	WT	40.4	41.0	41.5	41.5	43.0	43.4	42.7	44.9	45.2
	R789C	38.6	39.0	39.4	39.1	39.7	40.1	39.6	40.6	40.9
	R992C	38.5	39.2	38.7	39.3	40.0	39.7	40.2	40.7	40.6

^aTm₇₅, Tm₅₀, and Tm₂₅ indicate the temperatures at which the triple helical structure of analyzed collagens was preserved in 75%, 50%, and 25% of maximum, respectively; C indicates control samples not treated with chemical chaperones.

Figure Legends

Figure 1 - Relative amounts of Pro-GFP variants (A) and cPARP (B) in cells cultured in the presence of 0.5 M glycerol or 0.1 M TMAO. The relative amounts of Pro-GFP and cPARP in specific groups of cells treated with glycerol or TMAO are compared corresponding control groups cultured in the absence of these compounds (controls expressed as 100%). Each point represents the mean value from multiple independent assays \pm standard error of the mean (SEM). Symbols: WT, samples that include wild type Pro-GFP; R789C and R992C, Pro-GFP mutants; ns, not statistically significant; P values for statistically significant differences between means calculated for analyzed group and non-treated controls are also presented (expressed as 100%).

Figure 2 – A, Assays of the thermostabilities of the WT, R789C, and R992C collagen II variants isolated from HT-1080 cells cultured in the presence of 0.5 M glycerol or 0.1 M TMAO. B, Similar assays of the thermostability of the WT, R789C, and R992C collagen II variants isolated from HT-1080 cells cultured in chaperone-free media. Proteins were isolated from the media then incubated with 0.5 M glycerol or 0.1 M TMAO.

Following preincubation at increasing temperatures, collagen II samples were subjected to brief protease digestion. The digested samples were electrophoresed in 7.5% polyacrylamide gels in reducing conditions and then $\alpha 1(\text{II})$ chains were visualized by Western blot (Supplementary Figure S3). Subsequently, the relative pixel intensities of bands corresponding to $\alpha 1(\text{II})$ chains were measured by densitometry. Graphic representations of changes in the relative amount of triple-helical collagen II in response to increasing temperature are demonstrated (see Table 1). Symbols: WT, wild type collagen II; R789C and R992C, collagen II mutant harboring specific substitution; solid circles (●) and solid line, non-treated control; open circles (○) and dashed line, group treated with 0.5 M glycerol; triangles (▲) and dotted lines, a group treated with 0.1 M TMAO.

Figure 3 - Intracellular pool of Pro-GFP variants in cells cultured in the presence of glycerol or TMAO with or without addition of 2,2' dipyridyl. A, Site-by-site comparison of Pro-GFP variants from cells grown in the presence (+) or the absence (-) of 2,2' dipyridyl but in the absence of glycerol and TMAO. B, Intracellular pool of the Pro-GFP variants in cells grown in the presence of glycerol or TMAO but without 2,2' dipyridyl. C, Intracellular pool of the Pro-GFP variants in cells grown in the presence of 2,2' dipyridyl and glycerol or TMAO. Symbols: WT, samples that include wild type Pro-GFP; R789C and R992C, Pro-GFP mutants. Symbols: C, controls grown in the absence of glycerol (G) or TMAO (T).

Figure 4. Graphic representation of intracellular accumulation of Pro-GFP variants in cells cultured in the presence of glycerol or TMAO with or without addition of 2,2' dipyridyl. The relative amounts of Pro-GFP in specific groups of cells treated with glycerol or TMAO are compared to control groups cultured in the absence of these compounds (controls expressed as 100%). Each point represents the mean value from multiple independent assays \pm standard error of the mean (SEM).




Figure 1.




Figure 2.




Figure 3.




Figure 4.