

11-1-2009

Combination of vandetanib, radiotherapy, and irinotecan in the LoVo human colorectal cancer xenograft model.

Phyllis Wachsberger

Thomas Jefferson University, phyllis.wachsberger@mail.tju.edu

Randy Burd

University of Arizona

Anderson Ryan

AstraZeneca Pharmaceuticals

Constantine Daskalakis

Thomas Jefferson University

Adam P. Dicker

Thomas Jefferson University

[Let us know how access to this document benefits you](#)

Follow this and additional works at: <http://jdc.jefferson.edu/radoncfp>Part of the [Radiology Commons](#)

Recommended Citation

Wachsberger, Phyllis; Burd, Randy; Ryan, Anderson; Daskalakis, Constantine; and Dicker, Adam P., "Combination of vandetanib, radiotherapy, and irinotecan in the LoVo human colorectal cancer xenograft model." (2009). *Department of Radiation Oncology Faculty Papers*. Paper 12.
<http://jdc.jefferson.edu/radoncfp/12>

As submitted to:

International Journal of Radiation Oncology Biology

Physics

And later published as:

International Journal of Radiation Oncology Biology

Physics

Volume 75, Issue 3, 1 November 2009, Pages 854-861

DOI: 10.1016/j.ijrobp.2009.06.016

PubMed ID: 19801101

**“Combination of vandetanib, radiotherapy and
irinotecan in the LoVo human colorectal cancer
xenograft model”**

Phyllis Wachsberger¹, Ph.D., Randy Burd², Ph.D., Anderson Ryan³, Ph.D.,

Constantine Daskalakis⁴, Ph.D. and Adam P. Dicker¹, M.D., Ph.D.

¹Department of Radiation Oncology, Thomas Jefferson University,

Philadelphia, PA ; ²Department of Nutritional Sciences, University of

Arizona, Tucson, AZ; ³AstraZeneca Pharmaceuticals, Alderley Park,

Macclesfield, United Kingdom, ⁴Department of Pharmacology and

**Experimental Therapeutics, Thomas Jefferson University, Philadelphia,
PA.**

Corresponding Author

**Phyllis Wachsberger, Ph.D.
Department of Radiation Oncology, Thomas Jefferson University
111 South 11th Street
Philadelphia PA 19107
215 955 6659
FAX: 215 955 2052
Phyllis.wachsberger@jeffersonhospital.org**

**Running Title: Vandetanib, radiotherapy and irinotecan in LoVo
colorectal cancer**

Conflict of Interest: Statement

Anderson Ryan is an employee of AstraZeneca Pharmaceuticals.

Abstract:

Purpose: The tumor growth kinetics of the human LoVo colorectal xenograft model was assessed in response to vandetanib, an orally available receptor tyrosine kinase inhibitor, radiotherapy (RT) or irinotecan (CPT-11), as single therapies and in combination.

Methods and Materials: LoVo cells were injected subcutaneously into the right hind limb (5×10^6 cells in 100 μ l PBS) of athymic NCR NUM mice and tumors were grown to a volume of 200-300 mm^3 before treatment. Vandetanib was administered at 50 mg/kg daily p.o. for 14 days starting on day 1. RT was given as three fractions (3 x 3 Gy) on days 1, 2 and 3. CPT-

11 was given at 15 mg/kg i.p. on days 1 and 3. Tumor volumes were measured on a daily basis and calculated by measuring tumor diameters with digital calipers in two orthogonal dimensions.

Results: All three single treatments (vandetanib, CPT-11 and radiation) significantly slowed LoVo colorectal tumor growth. Vandetanib significantly increased the antitumor effects of CPT-11 and radiation, when given in combination with either of these treatments. These treatment combinations resulted in a slow tumor growth rate during the two weeks of vandetanib administration. The triple combination of vandetanib, CPT-11, and radiation produced the most marked improvement in response as observed by measurable shrinkage of tumors during the first week of treatment.

Conclusions: The tumor growth delay kinetics observed in this study of the LoVo colorectal model suggest concurrent and sustained post-sequencing of vandetanib with cytotoxic therapy may be beneficial in tumors of this type.

Key Words:

Vandetanib, radiotherapy, CPT-11, LoVo colorectal cancer, angiogenesis inhibitor

Introduction

Colorectal cancer (CRC) remains one of the leading causes of mortality worldwide. In recent years, the most widely used chemotherapy for metastatic CRC, fluoropyrimidine (5-fluorouracil (5-FU)) in combination with folinic acid (FA), has been combined with newer, highly active cytotoxic agents. Among these agents is the DNA topoisomerase I inhibitor, irinotecan (CPT-11)¹, a potent DNA-targeting drug used in patients with CRC that is refractory to treatment with fluorouracil and leucovorin. This cytotoxic agent is, in turn,

currently being combined with new molecular therapies targeting the tumor vasculature and key signaling pathways controlling tumor cell proliferation, angiogenesis and survival in CRC. In this regard, the epidermal growth factor receptor (EGFR) plays an important role in CRC tumor growth and progression ², and cetuximab, a monoclonal antibody (MAB) specific for EGFR, has been approved for use in combination with CPT-11 in patients with EGFR-expressing CRC refractory to CPT-11-based chemotherapy ³. In addition, bevacizumab, a MAB specific for vascular endothelial growth factor (VEGF-A), a key player in tumor angiogenesis in CRC as well as other solid tumors, has been approved for the treatment of metastatic CRC in combination with intravenous 5-FU-based chemotherapies ⁴. Despite recent improvements in treatment for CRC, a need still remains to improve the performance of existing treatments and to establish the optimum scheduling and dosing of combined therapies.

Vandetanib (ZACTIMATM) is an oral receptor tyrosine kinase inhibitor that, in recombinant enzyme assays, demonstrates potent activity against VEGFR-2 tyrosine kinase ($IC_{50} = 40$ nmol/L) with additional activity against EGFR ($IC_{50} = 500$ nmol/L) and RET ($IC_{50} = 130$ nM) tyrosine kinases ⁵⁻⁸. Vandetanib currently has orphan-drug status in the USA and Europe for medullary thyroid cancer (MTC) (in which RET activity is important) and is in Phase III development in non-small-cell lung cancer and MTC. Phase II studies are

ongoing to investigate its efficacy in other tumor types, thyroid cancer, hepatocellular carcinoma and glioblastoma.

Vandetanib has been shown to enhance the efficacy of radiotherapy in subcutaneous (s.c.) and orthotopic tumor xenograft models⁹⁻¹³. The combination of vandetanib, radiation and current chemotherapeutic agents used in CRC treatment has not been studied to date. Preclinical demonstration of efficacy of a combination protocol with novel agents plus radiation is usually considered crucial prior to clinical evaluation. The purpose of the present study was to examine the effect of vandetanib on the radiation response of a colorectal tumor model when administered in combination with CPT-11. It was hypothesized that simultaneous inhibition of VEGFR and EGFR by vandetanib in combination with the cytotoxic agent CPT-11 would interact to enhance radiation response and tumor control in the human LoVo colorectal tumor cell model. The LoVo colorectal model expresses activated EGFR^{14, 15} and is highly vascularized, and therefore is an appropriate model to test the hypothesis.

Materials and Methods

Animal and Tumor Model: LoVo cell suspensions (5×10^6 cells in 100 μ l phosphate buffered saline) were implanted s.c. into the right hind limbs of 6-8 week-old athymic NCR NUM mice (Taconic Farms, Hudson, NY). A s.c. xenograft model was chosen to facilitate radiation dosing and ease of tumor

measurements. Tumors were allowed to grow for approximately 25 days, until reaching an approximate volume of 200-300 mm³ at the start of treatment (day 1). All animals were randomized among treatment groups.

Drug Treatment: Vandetanib (AstraZeneca, Macclesfield, UK) was administered by oral gavage (p.o.) at 50 mg/kg daily for 14 days, starting on day 1. Vandetanib dosing in this study was based on previous pharmacokinetic studies in mouse models predicting relevance of this dosing to clinical drug exposure in human patients ¹⁶. CPT-11 was given at 15 mg/kg intraperitoneally (i.p.) on days 1 and 3.

Radiation Treatment: Irradiation was performed on anesthetized mice using X-rays generated by a PanTak, 310 kV_e x-ray machine, 0.25 mm Cu + 1 mm Al added filtration, at 125 cGy per min. Dosimetry was performed by an in-the-beam ionization chamber calibrated against a primary standard. Corrections were made daily for humidity, temperature and barometric pressure. Mice were anesthetized with a combination of ketamine and acepromazine at a concentration of 37.5 mg/kg and 0.2 mg/kg, respectively, to provide 25-30 minutes of sedation. Each mouse was confined in a lead casing with its tumor-bearing leg extended through an opening on the side to allow the tumor to be irradiated locally. Radiation was administered as three daily fractions of 3 Gy each on days 1, 2, and 3. On days when radiation was

administered with vandetanib and/or CPT-1, vandetanib and CPT-11 were given approximately 2 hours before radiation, with vandetanib preceding CPT-11 administration.

Tumor Measurement: Tumors were synchronized to be approximately 250 mm³ at the start of treatment (day 1) and were measured four to five times per week, for up to six weeks of follow-up, or until they reached 2,000 mm³. Tumor size was determined by direct measurement with calipers and calculated by the formula: (smallest diameter² x widest diameter)/2. Tumors were not allowed to grow beyond 2,000 mm³ in accordance with Institutional Animal Care and Use Committee regulations.

In-vivo Tumor Necrosis: Tumors were collected from animals on day 14 following start of treatment for fixation and staining with H&E. The area of necrosis was evaluated by image analysis and expressed as the percentage of the total tumor area.

Statistical Analysis Tumor growth was analyzed via mixed-effects regression, as previously described. The method was used because it does not depend on an arbitrary endpoint target tumor size, but utilizes the repeated tumor size measurements obtained over the entire study period, while appropriately handling unbalanced data (i.e., different number of measurements for different animals) and the correlation of each animal's

measurements over time. Mixed-effects regression yields generalizable parameters of interest (e.g., average daily tumor growth rate and tumor doubling time), and can investigate treatment interactions and non-linear patterns of tumor growth. The base-10 logarithm of tumor volume was modeled as a function of time and treatment. Linear or quadratic growth curves over time were fitted to the log-transformed data, depending on growth patterns in each treatment group. All statistical analyses were conducted in SAS 9.2 (SAS Institute Inc., Cary, NC, 1999-2001).

Results

The experiment involved three different treatments (vandetanib, CPT-11, and radiotherapy), as described above and summarized in Figure 1. Data were collected from a total of 104 animals in eight experimental groups (11-16 animals per group) and are summarized in Figure 2. Starting tumor sizes were comparable across groups, with geometric means ranging from 230 to 257 mm³ ($p = 0.771$). All treatments were well tolerated in the animals with no observable loss of body weight.

The three single-treatment groups (CPT-11, radiation, or vandetanib), as well as the combination of CPT-11 with radiation (Figure 2) were fitted to log transformed curves, while the three remaining groups that received combination treatments involving vandetanib showed a significantly non-linear tumor growth and were fitted to quadratic curves.

Figure 3 shows the measured geometric mean tumor size graphically over time. Table 1 shows the corresponding calculated tumor growth parameters (daily tumor growth rate and tumor doubling time). Table 2 shows p-values for group comparisons at 7, 14, and 21 days after start of treatment.

The control group had an estimated average daily tumor growth rate of 9.9%, corresponding to an average tumor doubling time of about 7 days (Table 1). All three single treatments resulted in a significant inhibition of tumor growth, compared with the control group (average daily tumor growth rates: CPT-11: 7.1%, $p = 0.015$; radiation: 5.6%, $p = 0.001$; vandetanib: 5.0%, $p = 0.001$). Vandetanib inhibited tumor growth significantly more than CPT-11 ($p = 0.043$) but not radiation ($p = 0.514$); radiation and CPT-11 were not significantly different ($p = 0.139$). The combination of CPT-11 with radiation produced a daily tumor growth rate of 5.1%, which was significantly lower than CPT-11 alone ($p = 0.015$) but comparable to radiation alone ($p = 0.560$). There was no significant (additive) interaction between CPT-11 and radiation ($p = 0.105$).

The remaining three groups which received treatment combinations involving vandetanib (with either CPT-11 or radiation, or with both CPT-11 and radiation), showed significant treatment interactions ($p = 0.001$ for the interaction between vandetanib and CPT-11 and between vandetanib and radiation) and non-linear tumor growth patterns. Compared to single-treatment groups, growth was significantly delayed (and, in the triple-

treatment combination, tumor volume actually decreased) early on, but progressively accelerated later, although it never exceeded that of the untreated controls (Figure 3). Because of the non-linearity of tumor growth in these groups, tumor growth parameters are not constant over time and comparisons depend on the timepoint referenced. Table 2 shows p-values for days 7, 14, and 21.

During the first week of treatment, animals receiving the combination of vandetanib with CPT-11 had average daily tumor growth rate of less than 3.5%, significantly lower than CPT-11 alone and marginally so compared to vandetanib alone ($p = 0.001$ and 0.058 , respectively, after 7 days). By the end of the two-week vandetanib treatment, the tumor growth rate in the combination group (4.6%) was still significantly lower than for CPT-11 alone ($p = 0.015$) but comparable to that for vandetanib alone ($p = 0.682$). By the third and fourth weeks, tumor growth had reached levels similar to those seen in the single-treatment groups (Figure 3, Table 1).

The combination of vandetanib with radiation resulted in a similar pattern of non-linear tumor growth inhibition. After the first 7 days, the average daily tumor growth rate of 2.1% was significantly lower than for either radiation alone or vandetanib alone ($p = 0.005$ and 0.019 , respectively). After 14 days, the tumor growth rate in the combination group had accelerated to 3.4% and was only marginally lower than for radiation alone and comparable to that for

vandetanib alone ($p = 0.080$ and 0.212 , respectively). By the third and fourth weeks, tumor growth had become similar to that seen in the single-treatment groups (Figure 3, Table 1).

Despite delaying tumor growth in the initial weeks, the treatment combinations induced only modest levels of tumor necrosis (10-20%), with no significant differences between treatment groups (Figure 4).

The pattern of tumor growth in the group that received the triple-treatment combination reflected both the interaction between vandetanib and CPT-11 and that between vandetanib and radiation (as mentioned previously). Thus, during the first week, instead of the delayed tumor growth seen in the two-treatment combinations, tumor volume in the triple-treatment combination actually decreased ($p = 0.001$ versus vandetanib plus CPT-11, and 0.052 versus vandetanib plus radiation). After that time, similar to the two-treatment combinations that involved vandetanib, tumor growth started accelerating. By the end of the third week, tumor growth in the triple-treatment combination group was similar to that in the two-treatment combination groups involving vandetanib, and by the fourth week, it was similar to that in the single-treatment groups.

Discussion

Relatively little is known about the antitumor effects of combining cytotoxic drugs, radiotherapy, and novel targeted therapies that specifically interfere with signaling pathways controlling cancer proliferation, angiogenesis and survival. In the present study, vandetanib, a potent inhibitor of both VEGFR and EGFR signaling, was combined with CPT-11 and/or radiation, to determine if greater anti-colorectal tumor activity can be obtained.

This study demonstrated that all three single treatments (vandetanib, CPT-11 and radiation) significantly slowed LoVo colorectal tumor growth. Previous studies with single-agent vandetanib demonstrated that chronic oral administration reduced tumor vascularity and tumor growth in a variety of xenograft models, including CRC^{7,17}. In the clinic, the safety and tolerability of vandetanib has been demonstrated in patients with advanced colorectal cancer as well as other solid tumors¹⁸. Vandetanib induced manageable normal tissue toxicities related to inhibition of EGFR and VEGFR signaling such as diarrhea, rash and hypertension^{19, 20}. The effect of combining radiation and vandetanib on normal tissue is currently unknown, however it has been shown in both preclinical and clinical trials that use of VEGF inhibitors with radiation may result in higher rates of normal tissue toxicity such as induction of thrombosis, hemorrhage and bowel toxicities²¹⁻²³. In contrast, it was postulated that combination of radiotherapy with inhibitors of angiogenesis may actually decrease these risks because radiotherapy has been used to prevent hemorrhage²⁴. Overall the investigation of agents such

as vandetanib in combination with radiation in normal tissue are lacking, and thus will be a major focus in the future.

As previously discussed, single agent vandetanib has dual tyrosine kinase inhibitory activity against VEGFR-2 and EGFR which allows it to target two key pathways responsible for tumor growth, i.e., tumor angiogenic signaling and tumor cell proliferation. It has been speculated that dual suppression may be critical for sustained suppression of tumor growth, especially since the EGFR and VEGFR pathways are linked and exhibit cross-talk²⁵. In addition, vandetanib can also enhance the antiproliferative activity of selective EGFR inhibitors such as cetuximab, thereby potentiating suppression of EGFR signaling¹⁷.

The present study confirmed that vandetanib, chronically administered over two weeks, slowed tumor growth in a colorectal tumor model, and, under the dosing conditions of this study, slowed tumor growth to a greater extent than CPT-11 alone and to a similar level to radiation alone. Moreover, vandetanib significantly increased the antitumor effects of CPT-11 and radiation, when given in combination with either of these treatments. In particular, these treatment combinations resulted in a slow tumor growth rate during the two weeks of vandetanib administration. These results confirm an earlier study by Troiani et al.²⁶, in which vandetanib (25 mg/kg/day) administered in combination with CPT-11 exhibited high antitumor activity in HT29-tumor-

bearing nude mice. Troiani et al. showed a correlation between this dosing schedule and enhanced EGFR and VEGFR signal inhibition.

In the present study, the triple combination of vandetanib, CPT-11, and radiation produced the most marked improvement in response in the LoVo-tumor-bearing mice. The triple treatment produced a measurable shrinkage of tumors during the first week of treatment. The combination of vandetanib, chemotherapy (gemcitabine), and radiation has also been previously shown to significantly inhibit tumor progression in a pancreatic tumor model ²⁷. Importantly, the present study also investigated the kinetics of tumor growth, both during and after a course of treatment. It was demonstrated that the addition of vandetanib significantly enhanced the initial antitumor effect of chemo-radiation. However, when vandetanib treatment ended, tumor growth returned to near control (untreated) levels. Therefore, these data support the rationale of adding an anti-vascular agent to cytotoxic therapies and provide valuable information for the design of therapeutic protocols.

The precise mechanisms leading to initial tumor regression with the combined therapies in this study are not known. Analysis of interactions between cytotoxic agents and vandetanib is complex, given that both the tumor cells and the tumor microenvironment are affected. In this connection, radiation can kill not only tumor cells but also endothelial cells of the tumor vasculature, thereby affecting the radiosensitivity of the tumor ^{28, 29}. In addition, cytotoxic agents have mechanisms of cell killing that are different from the targeted

agent. Both radiation and CPT-11 kill cells through DNA damage. Both chemotherapy and radiation can also alter cellular signaling pathways by inducing EGFR phosphorylation and through the growth factor signaling pathway, contribute to tumor cell proliferation and survival³⁰⁻³². Preclinical studies have also shown that cytotoxic therapy alone, such as radiation, can result in intensification of angiogenic processes³³. After cytotoxic treatment, up-regulation of vascular growth factors and their receptors occurs which contributes to tumor recurrence and progression³⁴. Direct up-regulation of VEGF after irradiation of various cancer cell lines has been reported³⁵. Radiation also induces transient tumor hypoxia which results in upregulation of hypoxia inducible factor -1 (HIF-1) which can stimulate VEGF and VEGFR2 expression. Therefore, simultaneous inhibition of both VEGFR and EGFR signaling through chronic administration of vandetanib in combination with cytotoxic therapy is expected to suppress the upsurge in pro-proliferative and angiogenic signaling resulting from CPT-11 and radiation-induced EGFR and VEGF. This suppression will thereby lead to inhibition of vascular protective mechanisms and growth factor mechanisms contributing to tumor regrowth.

The increased tumor growth that was seen in this study following discontinuation of vandetanib suggests that inhibition of angiogenic and pro-proliferative signaling is readily reversed. The current observations are in agreement with a number of both preclinical and clinical studies showing that tumors can adapt to anti-angiogenic treatment by undergoing “evasive

resistance” to angiogenesis inhibitors ³⁶. Mechanisms of resistance include upregulation of alternative proangiogenic signaling pathways as well as recruitment of bone marrow-derived proangiogenic cells ^{37, 38}. In addition, administration of vandetanib itself has been observed to increase VEGF production in certain cancer cell lines as well as in tumor xenografts ^{39, 40}, thereby suggesting an additional contributing mechanism to tumor relapse. More studies will be needed to determine whether additional angiogenic pathways may be induced by triple modality treatment.

Conclusions

The results of this study provide a scientific rationale for testing the combination of vandetanib, CPT-11, and radiation in patients with CRC. Although the best schedule and sequencing for this triple modality treatment has yet to be determined, the tumor growth delay kinetics observed in this study suggest that improvement in colorectal tumor response can be obtained by concurrent and sustained post-sequencing of vandetanib with cytotoxic therapy, keeping in mind that prolonged chronic administration of the receptor tyrosine kinase inhibitors may lead to the development of resistance and the requirement for additional therapeutic agents as seen with other targeted agents, such as imatinib and gefitinib ^{41, 42}.

Declaration of Interest: Phyllis Wachsberger received an unrestricted grant from AstraZeneca Pharmaceuticals.

Adam P. Dicker is supported by the National Institute of Health CA10663, Tobacco Research Settlement Fund (Commonwealth of Pennsylvania), USDA # 2006-03152 and the Christine Baxter Fund.

ZACTIMA is a registered trademark of the AstraZeneca group of companies.

Acknowledgments: The authors wish to acknowledge the expert technical assistance of Tara Strickler and Kari Kulp.

Reference List

- (1) Tewes M, Schleucher N, Achterrath W et al. Capecitabine and irinotecan as first-line chemotherapy in patients with metastatic colorectal cancer: results of an extended phase I study. *Ann Oncol* 2003 September;14(9):1442-8.
- (2) Roberts RB, Min L, Washington MK et al. Importance of epidermal growth factor receptor signaling in establishment of adenomas and maintenance of carcinomas during intestinal tumorigenesis. *Proc Natl Acad Sci U S A* 2002 February 5;99(3):1521-6.
- (3) Cunningham D, Humblet Y, Siena S et al. Cetuximab monotherapy and cetuximab plus irinotecan in irinotecan-refractory metastatic colorectal cancer. *N Engl J Med* 2004 July 22;351(4):337-45.
- (4) Hurwitz H, Fehrenbacher L, Novotny W et al. Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl J Med* 2004 June 3;350(23):2335-42.
- (5) Carlomagno F, Vitagliano D, Guida T et al. ZD6474, an orally available inhibitor of KDR tyrosine kinase activity, efficiently blocks oncogenic RET kinases. *Cancer Res* 2002 December 15;62(24):7284-90.
- (6) Ciardiello F, Caputo R, Damiano V et al. Antitumor effects of ZD6474, a small molecule vascular endothelial growth factor receptor tyrosine kinase inhibitor, with additional activity against epidermal growth factor receptor tyrosine kinase. *Clin Cancer Res* 2003 April;9(4):1546-56.

- (7) Ryan AJ, Wedge SR. ZD6474--a novel inhibitor of VEGFR and EGFR tyrosine kinase activity. *Br J Cancer* 2005 June;92 Suppl 1:S6-13.
- (8) Wedge SR, Ogilvie DJ, Dukes M et al. ZD6474 inhibits vascular endothelial growth factor signaling, angiogenesis, and tumor growth following oral administration. *Cancer Res* 2002 August 15;62(16):4645-55.
- (9) Brazelle WD, Shi W, Siemann DW. VEGF-associated tyrosine kinase inhibition increases the tumor response to single and fractionated dose radiotherapy. *Int J Radiat Oncol Biol Phys* 2006 July 1;65(3):836-41.
- (10) Damiano V, Melisi D, Bianco C et al. Cooperative antitumor effect of multitargeted kinase inhibitor ZD6474 and ionizing radiation in glioblastoma. *Clin Cancer Res* 2005 August 1;11(15):5639-44.
- (11) Frederick B, Gustafson D, Bianco C, Ciardiello F, Dimery I, Raben D. ZD6474, an inhibitor of VEGFR and EGFR tyrosine kinase activity in combination with radiotherapy. *Int J Radiat Oncol Biol Phys* 2006 January 1;64(1):33-7.
- (12) Sandstrom M, Johansson M, Bergstrom P, Bergenheim AT, Henriksson R. Effects of the VEGFR inhibitor ZD6474 in combination with radiotherapy and temozolomide in an orthotopic glioma model. *J Neurooncol* 2008 May;88(1):1-9.
- (13) Williams KJ, Telfer BA, Brave S et al. ZD6474, a potent inhibitor of vascular endothelial growth factor signaling, combined with radiotherapy: schedule-dependent enhancement of antitumor activity. *Clin Cancer Res* 2004 December 15;10(24):8587-93.

- (14) Koizumi F, Kanzawa F, Ueda Y et al. Synergistic interaction between the EGFR tyrosine kinase inhibitor gefitinib ("Iressa") and the DNA topoisomerase I inhibitor CPT-11 (irinotecan) in human colorectal cancer cells. *Int J Cancer* 2004 January 20;108(3):464-72.
- (15) Nyati MK, Maheshwari D, Hanasoge S et al. Radiosensitization by pan ErbB inhibitor CI-1033 in vitro and in vivo. *Clin Cancer Res* 2004 January 15;10(2):691-700.
- (16) Gustafson DL, Bradshaw-Pierce EL, Merz AL, Zirrolli JA. Tissue distribution and metabolism of the tyrosine kinase inhibitor ZD6474 (Zactima) in tumor-bearing nude mice following oral dosing. *J Pharmacol Exp Ther* 2006 August;318(2):872-80.
- (17) Morelli MP, Cascone T, Troiani T et al. Anti-tumor activity of the combination of cetuximab, an anti-EGFR blocking monoclonal antibody and ZD6474, an inhibitor of VEGFR and EGFR tyrosine kinases. *J Cell Physiol* 2006 August;208(2):344-53.
- (18) Morabito A, Piccirillo MC, Falasconi F et al. Vandetanib (ZD6474), a dual inhibitor of vascular endothelial growth factor receptor (VEGFR) and epidermal growth factor receptor (EGFR) tyrosine kinases: current status and future directions. *Oncologist* 2009 April;14(4):378-90.
- (19) Michael M, Gibbs P, Smith R, Godwood A, Oliver S, Tebbutt N. Open-label phase I trial of vandetanib in combination with mFOLFOX6 in patients with advanced colorectal cancer. *Invest New Drugs* 2009 June;27(3):253-61.

- (20) Natale RB, Bodkin D, Govindan R et al. Vandetanib Versus Gefitinib in Patients With Advanced Non-Small-Cell Lung Cancer: Results From a Two-Part, Double-Blind, Randomized Phase II Study. *J Clin Oncol* 2009 March 30.
- (21) Johnson DH, Fehrenbacher L, Novotny WF et al. Randomized phase II trial comparing bevacizumab plus carboplatin and paclitaxel with carboplatin and paclitaxel alone in previously untreated locally advanced or metastatic non-small-cell lung cancer. *J Clin Oncol* 2004 June 1;22(11):2184-91.
- (22) Levine RJ, Maynard SE, Qian C et al. Circulating angiogenic factors and the risk of preeclampsia. *N Engl J Med* 2004 February 12;350(7):672-83.
- (23) Senan S, Smit EF. Design of clinical trials of radiation combined with antiangiogenic therapy. *Oncologist* 2007 April;12(4):465-77.
- (24) Shibuya K, Komaki R, Shintani T et al. Targeted therapy against VEGFR and EGFR with ZD6474 enhances the therapeutic efficacy of irradiation in an orthotopic model of human non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2007 December 1;69(5):1534-43.
- (25) Pennell NA, Lynch TJ, Jr. Combined inhibition of the VEGFR and EGFR signaling pathways in the treatment of NSCLC. *Oncologist* 2009 April;14(4):399-411.
- (26) Troiani T, Serkova NJ, Gustafson DL et al. Investigation of two dosing schedules of vandetanib (ZD6474), an inhibitor of vascular endothelial growth factor receptor and epidermal growth factor receptor signaling, in combination with irinotecan in a human colon cancer xenograft model. *Clin Cancer Res* 2007 November 1;13(21):6450-8.

- (27) Bianco C, Giovannetti E, Ciardiello F et al. Synergistic antitumor activity of ZD6474, an inhibitor of vascular endothelial growth factor receptor and epidermal growth factor receptor signaling, with gemcitabine and ionizing radiation against pancreatic cancer. *Clin Cancer Res* 2006 December 1;12(23):7099-107.
- (28) Garcia-Barros M, Paris F, Cordon-Cardo C et al. Tumor response to radiotherapy regulated by endothelial cell apoptosis. *Science* 2003 May 16;300(5622):1155-9.
- (29) Paris F, Fuks Z, Kang A et al. Endothelial apoptosis as the primary lesion initiating intestinal radiation damage in mice. *Science* 2001 July 13;293(5528):293-7.
- (30) Balaban N, Moni J, Shannon M, Dang L, Murphy E, Goldkorn T. The effect of ionizing radiation on signal transduction: antibodies to EGF receptor sensitize A431 cells to radiation. *Biochim Biophys Acta* 1996 November 8;1314(1-2):147-56.
- (31) Schmidt-Ullrich RK, Mikkelsen RB, Dent P et al. Radiation-induced proliferation of the human A431 squamous carcinoma cells is dependent on EGFR tyrosine phosphorylation. *Oncogene* 1997 September 4;15(10):1191-7.
- (32) Van SS, Kyula J, Kelly DM et al. Chemotherapy-induced epidermal growth factor receptor activation determines response to combined gefitinib/chemotherapy treatment in non-small cell lung cancer cells. *Mol Cancer Ther* 2006 May;5(5):1154-65.

- (33) Koukourakis MI, Giatromanolaki A, Sivridis E et al. Squamous cell head and neck cancer: evidence of angiogenic regeneration during radiotherapy. *Anticancer Res* 2001 November;21(6B):4301-9.
- (34) Taylor AP, Osorio L, Craig R et al. Tumor-specific regulation of angiogenic growth factors and their receptors during recovery from cytotoxic therapy. *Clin Cancer Res* 2002 April;8(4):1213-22.
- (35) Gorski DH, Beckett MA, Jaskowiak NT et al. Blockage of the vascular endothelial growth factor stress response increases the antitumor effects of ionizing radiation. *Cancer Res* 1999 July 15;59(14):3374-8.
- (36) Bergers G, Hanahan D. Modes of resistance to anti-angiogenic therapy. *Nat Rev Cancer* 2008 August;8(8):592-603.
- (37) Casanovas O, Hicklin DJ, Bergers G, Hanahan D. Drug resistance by evasion of antiangiogenic targeting of VEGF signaling in late-stage pancreatic islet tumors. *Cancer Cell* 2005 October;8(4):299-309.
- (38) Du R, Lu KV, Petritsch C et al. HIF1alpha induces the recruitment of bone marrow-derived vascular modulatory cells to regulate tumor angiogenesis and invasion. *Cancer Cell* 2008 March;13(3):206-20.
- (39) Sasaki T, Tanno S, Shibukawa K, Osanai S, Kawabe J, Ohsaki Y. Administration of VEGF receptor tyrosine kinase inhibitor increases VEGF production causing angiogenesis in human small-cell lung cancer xenografts. *Int J Oncol* 2008 September;33(3):525-32.
- (40) Showalter TN, Darocz B, Halko R, Liu Y, Marrero N, Wachsberger PR. ZD6474 enhances radiation therapy in human glioblastoma xenografts which

contain the EGFRvIII mutation or overexpress wtEGFR. *Abstract, Radiation Research Society Annual Meeting 2008 (Boston, MA) 2008*;(Radiation Research Society Annual Meeting 2008).

- (41) Komarova NL, Katouli AA, Wodarz D. Combination of two but not three current targeted drugs can improve therapy of chronic myeloid leukemia. *PLoS ONE* 2009;4(2):e4423.
- (42) Okabe T, Okamoto I, Tsukioka S et al. Addition of S-1 to the epidermal growth factor receptor inhibitor gefitinib overcomes gefitinib resistance in non-small cell lung cancer cell lines with MET amplification. *Clin Cancer Res* 2009 February 1;15(3):907-13.

Tables

Table 1. Estimates of the average daily tumor growth rate and average tumor doubling time, by treatment group.

Table 2. P-values for comparisons of treatment groups, on days 7, 14 and 21, after the start of treatment.

Figures

Figure 1. Summary of treatment groups.

LoVo cells were implanted s.c. into the right hind limbs of athymic NCR NUM male mice. Mice were randomized into eight experimental groups (11-16 animals per group). Vandetanib was administered at 50 mg/kg daily p.o. for 14 days, starting on day 1. CPT-11 was given at 15 mg/kg i.p. on days 1 and 3. Radiation was given as three fractions (3 x 3 Gy) on days 1, 2, and 3.

Figure 2. Tumor growth curves in LoVo xenografts treated with vandetanib, CPT-11, and/or radiation.

Individual mouse data for eight treatment groups (11-16 animals per group), along with fitted group curves. Vandetanib was administered at 50 mg/kg daily p.o. for 14 days, starting on day 1. CPT-11 was given at 15 mg/kg i.p. on days 1 and 3. Radiation was given as three fractions (3 x 3 Gy) on days 1, 2, and 3.

Figure 3. Estimated geometric mean tumor volume over time in LoVo xenografts treated with vandetanib, CPT-11, and/or radiation.

Vandetanib was administered at 50 mg/kg daily p.o. for 14 days, starting on day 1. CPT-11 was given at 15 mg/kg i.p. on days 1 and 3. Radiation was given as three fractions (3 x 3 Gy) on days 1, 2, and 3.

Figure 4. H&E stained sections of LoVo colorectal xenografts.

All tumors were collected from animals on day 14 following start of treatment.

Areas of necrosis are denoted by nec. Magnification 20 X.

(A) Control (untreated) tumor, showing 2% necrosis.

(B) Tumor from animal after administration of last dose of vandetanib, showing 15% necrosis.

(C) Tumor from animal after administration of CPT-11 and RT, showing 20% necrosis.

(D) Tumor from animal after administration of vandetanib and CPT-11, showing 10% necrosis.