

7-1-2013

Angular cheilitis: a maligned condition.

Caren Campbell

Albert Einstein Medical Center, carencampbell@gmail.com

Lawrence Parish

Thomas Jefferson University, larryderm@yahoo.com

Let us know how access to this document benefits you

Follow this and additional works at: <http://jdc.jefferson.edu/dcbfp>



Part of the [Dermatology Commons](#)

Recommended Citation

Campbell, Caren and Parish, Lawrence, "Angular cheilitis: a maligned condition." (2013).

Department of Dermatology and Cutaneous Biology Faculty Papers. Paper 23.

<http://jdc.jefferson.edu/dcbfp/23>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Dermatology and Cutaneous Biology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Editorial

As submitted to:

Skinmed

And later published as:

Angular Cheilitis – A Maligned Condition

Volume 11, Issue 3, pp. 198-200.

July-August 2013

PMID: 24053003

Caren Campbell, MD 1

Medical Service, Albert Einstein Medical Center,

Philadelphia, PA

Lawrence Charles Parish, MD, MD(Hon)

Editor-in-Chief

Department of Dermatology and Cutaneous Biology

and the Jefferson Center for International Dermatology

Jefferson Medical College of Thomas Jefferson University

Philadelphia, Pennsylvania

1 E-mail: carencampbell@gmail.com

2 Address for Correspondence: Lawrence Charles Parish, MD, MD (Hon), 1760 Market Street, Philadelphia, PA 19103 E-mail: larryderm@yahoo.com

Why an essay on angular cheilitis, an easily recognized cutaneous entity? One suggestion rests with the observation that angular cheilitis could be one of the most maligned conditions seen in contemporary practice. The fissuring, redness, and soreness at the angles of the mouth may not be catastrophic, but the embarrassment and soreness are disconcerting. Another that is more disturbing concerns the unnecessary testing and treatment often seen in the community.

When a middle aged man presented with angular cheilitis on routine examination, he asked if this was due to a vitamin deficiency. Fig 1 He had been told by a physician and, of course, by Aunt Mabel that his diet was the problem.¹ In addition to having been prescribed multivitamins, a high potency steroid cream, and an antifungal preparation for presumed *Candida albicans* infection, he was instructed to eliminate several foods from his diet including anything spicy, plus carbonated beverages.

Causation

Angular cheilitis, also known as perleche, cheilosis, or angular stomatitis, is simply an inflammatory condition of the labial commissures, characterized by cracking, crusting, and in severe cases bleeding. Probably, the most common cause is the recession of the boney support of the lower aspects of the mouth. This can result in an overbite with the upper lip protruding over the lower. The situation can then be aggravated by dentition in less than stellar condition, or dentures that have not been adjusted in some time. A set-up for the problem may even have been initiated by thumb-sucking that continued long after the toddler years.

There are innumerable other causes that could be listed in a differential diagnosis, such as lip licking or actinic cheilitis Fig 2 & 3 For example, unilateral lesions due to trauma are short-lived, while bilateral lesions are more likely long-standing.² Peri-oral dermatitis

may also extend to the commissures. Fig 4 Oral thrush could also be manifested at the commissures, but these causes are infrequent.

Ultimately, mechanical trauma to the area is likely to be the primary culprit, but less common etiologies in practice, better known by physicians in training, are nutritional deficiency, particularly of riboflavin, iron, cobalamin, or zinc. These deficiencies are often cited whether due to malabsorption from diseases such as celiac or malnutrition due to anorexia nervosa, as well as bulimia nervosa.³ Patients suffering from diabetes, chronic renal failure, hepatitis, Sjogren's, Plummer Vinson or Crohn's can present with angular cheilitis, just to make the listing more complete.⁴⁻⁶ Medications have also been shown to cause angular cheilitis. Antineoplastic agents, such as sorafenib and selumetinib, can cause cheilitis in patients being treated with them for various types of malignancies.^{7,8} The condition is uncommonly caused or exacerbated by oral candidiasis or secondary bacterial infections.⁹ In patients who are immunocompromised or have diabetes, malignancy, or anemia the likelihood of infection is increased.¹⁰

Intervention

With the multifactorial origin of angular cheilitis, treatment focuses on eliminating precipitating factors.¹¹ If the patient is a cigarette, cigar, or pipe smoker, this could be a problem. Mid- to low- potency steroid ointment is helpful. Almost never is a lipstick allergy, or for that matter a reaction to toothpaste, the **cause**. While angular cheilitis could masquerade as contact dermatitis, Fig 5 the chronic nature of the condition and the ineffectiveness of the mid- to low- potency steroids can indicate that another condition should be considered, such as impetigo or herpes zoster.

Comment [CC1]: Mention other contact allergies – toothpaste ??

Conclusions

Angular cheilitis remains a chronic problem and is usually mechanical. While there are esoteric causes to be considered, more mundane etiologies are more likely. Identifying the underlying etiology is useful and allows for a more appropriate therapeutic approach, but looking for zebras should be reserved for the next safari!

1. Parish LC, Witkowski JA. The most important medical source: Aunt Mabel knows best. *Skinmed*. 2010 ;8:7-8.
2. Konstantinidis AB, Hatziotis JH. Angular cheilosis: an analysis of 156 cases. *J Oral Med*. 1984;39:199–206.
3. Strumia R. Dermatologic signs in patients with eating disorders. *Am J Clin Dermatol*. 2005;6:165–173.

4. Udayakumar P, Balasubramanian S, Ramalingam KS, et al. Cutaneous manifestations in patients with chronic renal failure on hemodialysis. *Indian J Dermatol Venereol Leprol.* 2006;72:119–125.
5. Soy M, Piskin S. Cutaneous findings in patients with primary Sjogren’s syndrome. *Clin. Rheumatol.* 2007;26:1350–1352.
6. Novacek G. Plummer-Vinson syndrome. *Orphanet J Rare Dis.* 2006;1:36.
7. Yang C-H, Lin W-C, Chuang C-K, et al. Hand-foot skin reaction in patients treated with sorafenib: a clinicopathological study of cutaneous manifestations due to multitargeted kinase inhibitor therapy. *Br. J. Dermatol.* 2008;158:592–596.
8. Balagula Y, Barth Huston K, Busam KJ, et al. Dermatologic side effects associated with the MEK 1/2 inhibitor selumetinib (AZD6244, ARRY-142886). *Invest New Drugs.* 2010. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20978926>. Accessed August 10, 2012.
9. Sharon V, Fazel N. Oral candidiasis and angular cheilitis. *Dermatol Ther.* 2010;23:230–242.
10. Rogers RS 3rd, Bekic M. Diseases of the lips. *Semin Cutan Med Surg.* 1997;16:328–336.
11. Park KK, Brodell RT, Helms SE. Angular cheilitis, part 2: nutritional, systemic, and drug-related causes and treatment. *Cutis.* 2011;88:27–32.

Fig 1 Angular cheilitis which had been incorrectly diagnosed as candidosis, as well as a vitamin deficiency

Fig 2 Lip licking in a 10 year old girl

Fig 3 Peri-oral dermatitis extending to the commissures

Fig 4 Contact dermatitis due to neomycin ointment