

## Thomas Jefferson University **Jefferson Digital Commons**

Department of Pathology, Anatomy and Cell Biology Faculty Papers Department of Pathology, Anatomy and Cell Biology

7-19-2008

# Transcriptional regulation network analysis of the hypertension-perturbed nucleus tractus solitarius

### Gregory E. Gonye

Daniel Baugh Institute for Functional Genomics/Computational Biology, Department of Pathology, Thomas Jefferson University, Philadelphia, PA 19107, USA, ggonye@mail.dbi.tju.edu

### Rajanikanth Vadigepalli

Daniel Baugh Institute for Functional Genomics/Computational Biology, Department of Pathology, Thomas Jefferson University, Philadelphia, PA 19107, USA, Rajanikanth. Vadigepalli@jefferson.edu

### Haiping Hao

Daniel Baugh Institute for Functional Genomics/Computational Biology, Department of Pathology, Thomas Jefferson University, Philadelphia, PA 19107, USA

### James S. Schwaber

Daniel Baugh Institute for Functional Genomics/Computational Biology, Department of Pathology, Thomas Jefferson University, Philadelphia, PA 19107, USA, James.Schwaber@jefferson.edu

### Let us know how access to this document benefits you

Follow this and additional works at: http://jdc.jefferson.edu/pacbfp

Part of the Genetic Processes Commons, Medical Cell Biology Commons, and the Neurosciences Commons

### Recommended Citation

Gonye, Gregory E.; Vadigepalli, Rajanikanth; Hao, Haiping; and Schwaber, James S., "Transcriptional regulation network analysis of the hypertension-perturbed nucleus tractus solitarius" (2008). *Department of Pathology, Anatomy and Cell Biology Faculty Papers*. Paper 54. http://jdc.jefferson.edu/pacbfp/54

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's Center for Teaching and Learning (CTL). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Pathology, Anatomy and Cell Biology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

### **BMC Neuroscience**



Poster presentation

**Open Access** 

# Transcriptional regulation network analysis of the hypertension-perturbed nucleus tractus solitarius

Gregory E Gonye\*, Rajanikanth Vadigepalli, Haiping Hao and James S Schwaber

Address: Daniel Baugh Institute for Functional Genomics/Computational Biology, Department of Pathology, Thomas Jefferson University, Philadelphia, PA 19107, USA

Email: Gregory E Gonye\* - ggonye@mail.dbi.tju.edu

\* Corresponding author

from Seventeenth Annual Computational Neuroscience Meeting: CNS\*2008 Portland, OR, USA. 19–24 July 2008

Published: II July 2008

BMC Neuroscience 2008, 9(Suppl 1):P31 doi:10.1186/1471-2202-9-S1-P31

This abstract is available from: http://www.biomedcentral.com/1471-2202/9/S1/P31

© 2008 Gonye et al; licensee BioMed Central Ltd.

#### Introduction

Neurons expressing the angiotensin II (angII) AT1 receptor (AT1R) in the nucleus tractus solitarius (NTS) have been implicated in many of the functions assigned to angII in the CNS including roles in neural control of cardiovascular function, in particular blood pressure regulation, the baroreflex, stress responses and maintenance of homeostasis (thirst and salt appetite). AngII, acting through the AT1 receptor has two main effects on neurons: a rapid and transient modulation of K+ and Ca2+ currents and longer lasting changes in gene expression, ultimately leading to modulation of noradrenergic neuronal networks regulating cardiovascular function [1]. Using a combined experimental and computational systems level analysis, we have investigated the transcriptional regulatory network downstream of AT1R activation by using N1E-115 neuroblastoma cells engineered to express AT1Rs as a model system and have recently extended this to the NTS.

### **Methods**

We have applied transcriptional regulatory network analysis (TRNA), using our bioinformatics framework, Promoter Analysis and Interaction Network Toolset (PAINT) [2] to the response of the NTS both directly to angII and to phenylephrine-induced acute hypertension. TRNA makes the fundamental assumption that many coexpressed genes share transcription regulatory elements (TRE's), typically transcription factor binding sites, in their promoters, leading to co-regulation. This results in a

higher frequency of the suspect binding sites in the promoters of sets of co-regulated genes relative to randomly associated sets of genes. PAINT uses bioinformatics in combination with robust statistical approaches to identify those regulatory elements significantly enriched in the promoters of the genes of interest relative to random gene sets. Predicted TF:promoter interactions are tested using chromatin immunoprecipitation PCR (ChIP-PCR) with chromatin from untreated and perturbed (angII or hypertension) NTS or cell lines.

### **Results and conclusions**

TRNA predictions generated from microarray-based gene expression analysis of stimulated NTS ex vivo by 100 nM angII for 4 hours (n = 4) are shown in Table 1. Predicted TF's include those previously described as angII-responsive in neurons (CREB, KROX), angII-responsive in nonneuronal cells (E2F, ATF and MEF2), and novel associations (Pax, cRel). Validation studies using chromatin immunoprecipitation (ChIP) to test specific TF::promoter interactions in the in vivo NTS and in an engineered AT1Rexpressing neuronal cell line (N1E-AT1). Several predicted interactions have been confirmed, demonstrating both angII and hypertension-dependent binding dynamics. TRNA has successfully predicted TF activity dynamics at specific promoters relevant to neuronal function in hypertension and angII perturbed NTS. Going forward, these validated interactions will be manipulated in vivo to directly demonstrate a role in CNS cardiovascular regulation.

Table I: TRNA of ATIR-dependent NTS Expression

TF Family	Enriched TRE	Fisher's p-value
CREB	V\$CREB_Q2_01	0.045
ATF	V\$ATFI_Q6	0.016
ATF	V\$ATF3_Q6	0.035
ATF	V\$ATF6_01	0.027
E2F	V\$E2F1_Q3_01	0.016
MEF2	V\$ZID_01	0.0065
Pax	V\$PAX3_B	0.0012
cRel	V\$CREL_01	0.0023
KROX/egrI	V\$KROX_Q6	0.036

### Acknowledgements

The authors wish to acknowledge the expert technical assistance of H. Liu, Y. Liu and M. Payne in support of these studies and the intellectual input of the other members of the Daniel Baugh Institute for Functional Genomics and Computational Biology. This work was supported by grants from the NIGMS and NHLBI (J. Schwaber, PI).

#### References

- Phillips MI, Sumners C: Angiotensin II in central nervous system
- physiology. Regul Pept 1998, 78(1-3):1-11. Vadigepalli R, Chakravarthula P, Zak DE, Schwaber JS, Gonye GE: PAINT: a promoter analysis and interaction network generation tool for gene regulatory network identification. Omics 2003, **7(3):**235-252.

Publish with **Bio Med Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours you keep the copyright

Submit your manuscript here: http://www.biomedcentral.com/info/publishing\_adv.asp

