

9-15-2010

Using hospital discharge abstract data to identify incident breast cancer cases and assess quality of care

Elaine J. Yuen
Thomas Jefferson University

Daniel Z. Louis
Thomas Jefferson University

Carol Rabinowitz
Thomas Jefferson University

Vittorio maio

L. Cisbani
Agenzia Sanitaria e Sociale Regionale, Regione Emilia-Romagna, Bologna, Italy

Follow this and additional works at: <https://jdc.jefferson.edu/healthpolicyfaculty>



Part of [Digital Commons](#), and the [Oncology Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Yuen, Elaine J.; Louis, Daniel Z.; Rabinowitz, Carol; maio, Vittorio; Cisbani, L.; DePalma, R.; and Grilli, R., "Using hospital discharge abstract data to identify incident breast cancer cases and assess quality of care" (2010). *College of Population Health Faculty Papers*. Paper 57.
<https://jdc.jefferson.edu/healthpolicyfaculty/57>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in College of Population Health Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Elaine J. Yuen, Daniel Z. Louis, Carol Rabinowitz, Vittorio maio, L. Cisbani, R. DePalma, and R. Grilli

MEETING ABSTRACT

Open Access

Using hospital discharge abstract data to identify incident breast cancer cases and assess quality of care

E J Yuen^{1*}, D Z Louis¹, C Rabinowitz¹, V Maio¹, L Cisbani², R DePalma², R Grilli²

From 26th Patient Classification Systems International (PCSI) Working Conference Munich, Germany. 15-18 September 2010

Introduction

This study assesses the feasibility of using hospital discharge data to identify women with incident breast cancer, and it evaluates the use of these data to assess compliance with an indicator of quality of care. The study's goals are to: 1) compare breast cancer case-finding algorithms based upon ICD-9 CM codes found in hospital discharge data with cancer registry data and 2) compare the utility of these algorithms for use with a selected quality indicator.

Methods

Women aged 20 or greater who had continuous residence in the Emilia-Romagna Region, Italy, between 2002 and 2005 (N=1,887,212) were included in the analysis. The regional breast cancer tumor registry was used as the "gold standard" in identifying incident cases. Algorithms using ICD-9 CM codes on the Italian hospital discharge data were developed to identify women with incident breast cancer requiring a diagnosis code for cancer, as well as a surgical code for lumpectomy or mastectomy.

These hospital discharge-based algorithms were then compared to the registry data. Sensitivity, specificity and positive predictive value were computed overall and by age and cancer stage. These two case-finding approaches were also used to evaluate the use of radiation therapy one year post diagnosis for eligible women.

Results

15,469 women were identified by the regional cancer registry as incident cases, while 15,218 women were

identified by the hospital discharge data algorithm. Overall sensitivity was 87.5%, specificity was 99.9%, and positive predictive value (PPV) was 89.0%. Of the 1,677 who were false positives, 1,060 (64%) were women who had an incident case prior to 2000. There were 1,928 women who were false negatives. Many of these women were older and did not receive a surgical procedure. Therefore, they were not included in our hospital data-based, case-finding algorithm.

Sensitivity declined as the patient population became older; however, we observed relatively good PPV for all age groups. For the radiation therapy quality indicator, we identified 5,847 women using the cancer registry who were included as the population at risk (denominator), while hospital data-based algorithms identified 7,262 women. The estimated overall compliance with the radiation therapy indicator using the registry was 90.2%, compared to 78.8% using the hospital-based case-finding algorithm. This lower rate for the hospital-based population was chiefly due to a larger identified population at risk (false positives).

Conclusions

This project confirms the feasibility of using routinely available hospital discharge abstract data to identify women with incident breast cancer with good sensitivity, specificity, and positive predictive value when compared to cancer registry data. These hospital claims-based algorithms allow for analyses of large populations for which registry data may not be routinely available, thus allowing for the study of subsequent patterns and outcomes of care.

¹Thomas Jefferson University, Philadelphia, PA, USA
Full list of author information is available at the end of the article

Author details

¹Thomas Jefferson University, Philadelphia, PA, USA. ²Agenzia Sanitaria e Sociale Regionale, Regione Emilia-Romagna, Bologna, Italy.

Published: 6 October 2010

doi:10.1186/1472-6963-10-S2-A1

Cite this article as: Yuen *et al.*: Using hospital discharge abstract data to identify incident breast cancer cases and assess quality of care. *BMC Health Services Research* 2010 **10**(Suppl 2):A1.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

