

# Thomas Jefferson University Jefferson Digital Commons

Department of Medicine Faculty Papers

Department of Medicine

8-30-2011

# 16-year trends in the infection burden for pacemakers and implantable cardioverterdefibrillators in the United States 1993 to 2008.

Arnold J Greenspon Thomas Jefferson University, arnold.greenspon@jefferson.edu

Jasmine D Patel *Drexel University* 

Edmund Lau Drexel University

Jorge A Ochoa Drexel University

Daniel R Frisch Thomas Jefferson University

See next page for additional authors

Follow this and additional works at: http://jdc.jefferson.edu/medfp Part of the <u>Cardiology Commons</u>, and the <u>Medical Genetics Commons</u>

# **Recommended** Citation

Greenspon, Arnold J; Patel, Jasmine D; Lau, Edmund; Ochoa, Jorge A; Frisch, Daniel R; Ho, Reginald T; Pavri, Behzad B; and Kurtz, Steven M, "16-year trends in the infection burden for pacemakers and implantable cardioverter-defibrillators in the United States 1993 to 2008." (2011). *Department of Medicine Faculty Papers*. Paper 60. http://jdc.jefferson.edu/medfp/60

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's Academic & Instructional Support & Resources Department (AISR). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Medicine Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

#### Authors

Arnold J Greenspon, Jasmine D Patel, Edmund Lau, Jorge A Ochoa, Daniel R Frisch, Reginald T Ho, Behzad B Pavri, and Steven M Kurtz

As submitted to: Journal of the American College of Cardiology

And later published as:

Sixteen Year Trends in the Infection Burden for Pacemakers and Implantable Cardioverter-Defibrillators in the United States: 1993-2008

doi:10.1016/j.jacc.2011.04.033 Volume: 58, Issue: 10, Pages: 1001-1006, 2011

Arnold J. Greenspon M.D. <u>FACC</u><sup>1</sup>, Jasmine D. Patel PhD<sup>2,3</sup>, Edmund Lau MS<sup>2,3</sup>, Jorge Ochoa PhD<sup>3</sup>, Daniel Frisch M.D. FACC<sup>1</sup>, Reginald Ho M.D. FACC<sup>1</sup>, Behzad Pavri M.D. FACC<sup>1</sup>, Steven Kurtz PhD<sup>2,3</sup>

<sup>1</sup> Thomas Jefferson University Hospital, Philadelphia, PA

<sup>2</sup> Drexel University, Philadelphia, PA

<sup>3</sup> Exponent, Philadelphia, PA

Key words: endocarditis, infection, pacemakers, ICDs.

Word count-Title=17, Text (including abstract) = 1896, References= 525, Figure Legends= 53 Total=2491

Running Title: Infection Trends for Pacemakers and ICDs

Address for correspondence: Arnold J. Greenspon M.D. FACC Jefferson Heart Institute 925 Chestnut Street, Mezzanine Philadelphia, PA 19107 215-955-8659 Email: arnold.greenspon@jefferson.edu

Field Code Changed

# Abstract

**Objectives:** We analyzed the infection burden associated with the implantation of cardiac implantable electrophysiologic devices (CIED) in the United States for the years 1993-2008.

**Background:** Recent data suggests that the rate of infection following CIED implantation may be increasing.

**Methods:** The Nationwide Inpatient Sample (NIS) discharge records were queried between 1993-2008 using the 9th Revision of the International Classification of Diseases (ICD-9-CM). CIED infection was defined as either: (1) ICD-9 code for device related infection (996.61) and any CIED procedure or removal code, or (2) CIED procedure code along with systemic infection. Patient health profile was evaluated by coding for renal failure, heart failure, respiratory failure, and diabetes mellitus. The infection burden and patient health profile were calculated for each year and linear regression was used to test for changes over time.

**Results:** During the study period (1993-2008), the incidence of CIED infection was 1.61%. The annual rate of infections remained constant until 2004 when a marked increase was observed, which coincided with an increase in the incidence of major co-morbidities. This was associated with a marked increase in mortality and in-hospital charges.

**Conclusions:** The infection burden associated with CIED implantation is increasing over time and is associated with prolonged hospital stays and high financial costs.

# Abbreviations:

PM= Permanent pacemaker ICD= Implantable cardioverter defibrillator CIED= Cardiac implantable electrophysiologic device ICD-9= 9<sup>th</sup> Revision of the International Classification of Diseases NIS= National In-Patient Sample CRT= Cardiac resynchronization therapy

# Introduction

Implantation of cardiac implantable electrophysiology devices (CIED) which include permanent pacemakers (PM) and implantable cardioverter defibrillators (ICDs) has dramatically increased over the past several years.<sup>1,2,3</sup> This is largely due to the expanded indications for CIED implantation based on the results of large clinical trials of ICDs for primary prevention as well as the aging of the general population.<sup>4,5,6</sup> Infection associated with CIEDs is a serious complication with high morbidity and mortality.<sup>7,8,9</sup> Previous studies have suggested that the number of infections associated with CIED is increasing.<sup>10,11</sup> We sought to analyze the historical trends for CIED infection in the United States over <u>sixteen</u> years and evaluate the implications of these trends.

Deleted: the past 15

#### Methods

The Nationwide Inpatient Sample (NIS) discharge records were queried to identify demographic (e.g., age, gender), health profile/risk (incidence and severity of comorbidities, mortality) and health economic (length of stay, procedural costs and charges) data for PM and ICD patients between 1993-2008 using the 9th Revision of the International Classification of Diseases (ICD-9-CM). Specifically, procedures were identified by the ICD-9-CM codes that identified both primary and revision CIED procedures: Primary PM: 37.80-83, 00.50; Primary ICD: 37.94, 37.96, 00.51; PM Removal: 37.79, 37.85-87, 37.89, 00.53; ICD Removal: 37.98, 00.54. Revision procedures include pulse generator replacement as well as device upgrades to either dual chamber or cardiac resynchronization therapy (CRT) devices. During this time period, the ICD-9-CM codes for these procedures have been consistent, thereby allowing the analysis of longitudinal trends in the data for prevalence of device implantation.

Patients with a CIED-related infection, either pocket infection or systemic infection including lead-associated endocarditis, were identified in one of two ways: (1) an ICD-9 diagnosis code for device related infection (996.61) along with any CIED primary procedure or removal code, or (2) a CIED device removal code (37.77, 37.7, 37.89, 37.99) along with evidence of systemic infection such as sepsis (038 or 785.59), bacteremia (790.7) or fever (780.6). Patient health profile was evaluated by coding for renal failure, heart failure, respiratory failure, and diabetes mellitus.

The CIED infection burden was calculated by dividing the number of device related infections by the corresponding number of primary or revision procedures. Analyses of the NIS records with the relevant surgical codes were conducted using SAS (Version 9.2). The sampling weights and the stratified sampling design of the NIS were taken into consideration when computing summary statistics and standard errors of these estimates. The number of surgeries performed for a particular demographic group is a positive integer and is assumed to follow a Poisson distribution. A regression model was used to estimate the surgery and infection rate, and normalized by the size of the population, and evaluation of the calendar year trend. The surgery rate was adjusted by age, sex, race, and census regions to accommodate differences in the prevalence among demographic subpopulations. The infection burden and patient health profile were calculated for each year and linear regression was used to test for changes over time.

#### Results

#### **Trends in CIED infection**

Between 1993 and 2008, over 4.2 million primary implantations of PM (3,204,700 records) and ICD (1,124,000 records) were identified using ICD-9-CM procedure codes. We found that the incidence of CIED implantation increased an average of 4.7% annually and the overall CIED implantation increased by 96% from 1993-2008 (Figure 1). The majority of this increase was due to the large increase in ICD implantation (504%) as pacemaker implantation increased by 45% during this time period. By 2008, ICDs represented 35% of all CIED implantations (Figure 2).

During the study period (1993-2008), approximately 69,000 patients were treated for CIED infection (incidence= 1.61%). The incidence of infection increased by 210% from 2660 cases in 1993 to 8230 cases in 2008. The annual rate of infections remained fairly constant until 2004 when a marked increase was observed. The rate of infection increased significantly from 1.53% in 2004 to 2.41% in 2008 (p<0.001) (Figure 3).

The rates of CIED infection from 1993-2008, categorized by patient demographics (age, gender, race), showed that the highest infection rates occurred in white (82%), male (67%) patients over the age of 65 (64%). (Figure 4).

#### Role of comorbidities in CIED infection

The incidence of four major comorbidities (renal failure, respiratory failure, heart failure and diabetes) in patients with CIED infection remained fairly constant from 1993 through 2004 when a marked increase was observed (Figure 5). In addition, the risk of mortality significantly increased in patients with respiratory failure (odds ratio = 13.58; 95% CI 12.88-14.3), renal failure (odds ratio = 4.28; 95% CI 4.04-4.53), heart failure (odds ratio = 2.71; 95% CI 2.54-2.88) but decreased slightly in patients with diabetes (odds ratio = 0.91; 95% CI 0.86-0.96) (p<0.001).

#### Financial burden and mortality rates associated with CIED infection

In 1993, in-hospital charges for CIED infection were approximately \$75,000 and increased to over \$146,000 by 2008, an increase of 47% per decade (Figure 6). In-patient mortality associated with CIED infection averaged 4.39%, but increased from 2.91% in 1993 to 4.69% in 2008, representing an increase of 1% per decade. During the study period hospitalization remained constant and averaged 13.8 days.

# Discussion

An analysis of the Nationwide Inpatient Sample (NIS) demonstrates that during the study period 1993-2008, the national CIED infection burden has increased. Specifically, there has been an increase in the incidence of CIED infection along with an increase in inpatient mortality. Current patients have a high number of clinical comorbidities associated with prolonged hospital stays and an increase in the utilization of medical resources reflected by an increase in hospital charges.

Over the past 20 years, the number of patients with CIEDs has dramatically increased.<sup>1,2,3</sup> Among Medicare beneficiaries, the rate of cardiac device implantation increased by 42% between 1990-1999.<sup>10</sup> Our initial analysis of NIS data<sup>1</sup>, which included patients with all types of insurance coverage, showed a 30% increase in the primary pacemaker implantation rate of 50.0 per 100,000 persons of population in 1993 to 65.1 per 100,000 in 2006. ICD implantation rate increased more rapidly during this period of time. In 1993, the ICD implantation rate was 6.1 per 100,000 persons of population and rose dramatically to 46.2 per 100,000 persons of population by 2006, an increase of over 500%.

The present study demonstrates that overall CIED implantation increased by 96%. Most of this increase was due to the marked increase in ICD utilization. By the end of the study period, ICDs represented 35% of all devices. It is estimated that CIED device utilization will continue to grow over the next several years due to expanded Medicare coverage for these devices.<sup>12</sup> Complications of CIED implantation are an important consideration in patient selection for CIED implantation. Interestingly, immediate post procedural complications related to CIED implantation have decreased. Al-Khatib and co-workers reported that the rate of post procedural complications in CIED recipients fell between 2002-2005.<sup>13</sup> The fall in procedure related complications may be due to operator experience, improved device technology, and patient selection.<sup>13,14,15</sup> Unfortunately, the corresponding risk of device-related infection has not changed during the same period of time.<sup>8,11, 11,16,17</sup>

Previous studies have attempted to define the burden of CIED infection. Voigt and colleagues analyzed records from the National Hospital Discharge Survey (NHDS) between 1996 and 2006.<sup>11</sup> Analysis of the NHDS database showed that device related infection in CIED patients increased out of proportion to the overall increase in device utilization during the period 1996-2006. The NIS database, analyzed in the present study, samples about 25 times more

discharge records than the NHDS and tends to sample data from larger institutions. This may account for some variation between the two surveys.

Results from our analysis of the NIS show that the annual incidence of CIED infection increased by 210% to 2.41% in 2008 (p<0.0001). Our data may more accurately reflect the true CIED infection burden since previous studies may have included patients with infections of other cardiac devices such as prosthetic heart valves and not CIED alone. We required a CIED procedure code along with the 996.61 code to define a CIED infection. We also defined a device related infection as evidence of sepsis or bacteremia along with a CIED removal code. This study and others highlight the disturbing trend of increasing CIED infection. <sup>10,11,10,11,16,16</sup>

CIED infection is associated with high patient morbidity and a mortality rate of up to 18%.<sup>Error! Bookmark not defined.9,8,9,16</sup> The financial burden of CIED infection is reflected by the 47% per decade increase in hospital charges related to CIED infection. By 2008, hospital charges were over \$146,000. These expenditures do not include the additional costs of prolonged recovery and rehabilitation following treatment of the infection. Therefore, CIED infection has enormous economic implications.<sup>18</sup>

The reason for the increasing rate of CIED infection despite a decrease in overall device related complications is not clear. One possibility for this observation includes the increasing numbers of ICD and CRT devices whose longevity is significantly lower than PM. It is estimated that over 70% of ICD recipients will require device replacement surgery.<sup>19</sup> Device replacement surgery is associated with an increased risk of infection.<sup>19,20</sup> There may be an increasing burden of device replacements in the overall CIED population since ICDs now represent 35% of all implantations.

Patient characteristics, in addition to replacement burden, likely contribute to the increasing infection burden. It is well known that patients with chronic renal insufficiency and diabetes mellitus are at particular risk for CIED infection.<sup>21</sup> Our analysis showed that there is an increasing incidence of these risk factors in CIED patients. While the present study shows that the incidence of comorbidities in patients who present with CIED infection is increasing it does not address the important question of what specific risk factors predict CIED infection or what factors might mitigate this issue. However, it does appear that patients with multiple comorbidities are at particular risk. Further study of these critical issues is important.

In summary, the infection burden associated with CIED implantation is increasing over time. This is likely due to expanding ICD indications and the increasing comorbidities in the CIED population. Infection is associated with prolonged hospital stays and high financial costs. Further investigation into the risk factors for CIED infection is warranted.



Figure 1. Annual Number of PM and ICD Implantations: 1993-2008





Figure 3. Rate of CIED infection.





Figure 4. Distribution of CIED infection based on patient age.



Figure 5. Incidence of comorbidities in patients with CIED infection



# Figure 6. In-hospital charges associated with CIED infection (inflation adjusted to \$2009)

<sup>1</sup> Kurtz SM, Ochoa JA, Lau E, et al: Implantation trends and patient profiles for pacemakers and implantable cardioverter defibrillators in the United States: 1993-2006. Pacing Clin Electrophysiol 2010;33:705-711

<sup>2</sup> Mond HG, Irwin M, Ector H, Proclemer A. The world survey of cardiac pacing and cardioverter defibrillators: Calendar year and Electrophysiology 2005- an International Cardiac Pacing and Electrophysiology Society (ICPES) project. Pacing Clin Electrophysiol 2008;31:1202-1212

<sup>3</sup> Uslan DZ, Tleyjeh IM, Baddour LM, St Sauver JL, Hayes DL: Temporal trends in permanent pacemaker implantation: A population study. Am Heart J: 2008;155:896-903

<sup>4</sup> Myerburg RJ: Implantable cardioverter-defibrillators after myocardial infarction. N Eng J Med 2008;359:2245-2253

<sup>5</sup> Bardy GH, Lee KL, Mark DB, et al.: Sudden cardiac death in heart failure trial (SCD-HeFT). Amiodarone or an implantable defibrillator for congestive heart failure. N Eng J Med 2005;352:225-237

<sup>6</sup> Moss AJ, Zareba W, Hall WJ et al.: Multicenter Automatic Defibrillator Implantation Trial II Investigators. Prophylactic implantation of a defibrillator in patients with myocardial infarction and reduced ejection fraction. N Eng J Med 2002;346:877-883

<sup>7</sup> Darouiche RO: Treatment of infections associated with surgical implants. N Eng J Med 2004;350:1422-1429

<sup>8</sup> Sohail MR, Uslan DZ, Khan AH, et al.: Management and outcome of permanent pacemaker and implantable cardioverter-defibrillator infections. J Am Coll Cardiol 2007;49:1851-1859

<sup>9</sup> Baddour LM, Epstein AE, Erickson CC, et al.: Update on cardiovascular implantable electronic device infections and their management. A scientific statement from the American Heart Association. Circulation 2010;121:458-477

<sup>10</sup> Cabell CH, Heidenreich PA, Chu VH, et al: Increasing rates of cardiac device infections among Medicare beneficiaries:1990-1999. Am Heart J 2004;147:582-586

<sup>11</sup> Voigt A, Shalaby A, Saba S: Continued rise in rates of cardiovascular implantable device infection in the United States: Temporal trends and causative insights. Pacing Clin Electrophysiol 2010;33:414-419

<sup>12</sup> McClellan MB, Tunis SR: Medicare coverage of ICDs. N Eng J Med 2005;352:222-224

<sup>13</sup> Al-Khatib SM, Lucas Jollis JG, Malenka DJ, Wennberg DE: The relation between patients' outcome and the volume of cardioverter-defibrillator implantation procedures performed by physicians treating Medicare beneficiaries. J Am Coll Cardiol 2005;46:1536-1540

<sup>14</sup> Peterson PN, Daugherty SL, Wand Y, et al.: Gender differences in procedure-related adverse events in patients receiving implantable cardioverter-defibrillator therapy. Circulation 2009;119:1078-1084

<sup>15</sup> Curtis JP, Luebbert JJ, Wang Y, et al.: Association of physician certification and outcomes among patients receiving an implantable cardioverter-defibrillator. JAMA 2009;301:1661-1670

<sup>16</sup> Nery PB, Fernandes B, Nair GM, et al.: Device-related infection among patients with pacemakers and implantable defibrillators: Incidence, risk factors, and consequences. J Cardiovasc Electrophysiol 2010;21:786-790

<sup>17</sup> Uslan DZ, Sohail MR, St Sauver JL, et al.: Permanent pacemaker and implantable cardioverter-defibrillator infection. A population-based study. Arch Int Med 2007;167:669-675

<sup>18</sup> Ferguson TB Jr., Ferguson CL, Crites K, Crimmins-Reda P: The additional hospital costs generated in the management of complications of pacemaker and defibrillator implantations. J Thorac Cardiovasc Surg 1996;111:742-75

<sup>19</sup> Borleffs CJM, Thijssen J, De Bie MK, et al.: Recurrent implantable cardioverter-defibrillator replacement is associated with an increasing risk of pocket-related complications. Pacing Clin Electrophysiol 2010;33:1013-1019

<sup>20</sup> Poole JE, Gleva MJ, Mela T, et al. for the REPLACE Registry Investigators: Complication rates associated with pacemaker or implantable cardioverter-defibrillator replacements and upgrade procedures: Results from the REPLACE registry. Circulation 2010;122:1553-1561

<sup>21</sup> Dasgupta A, Montalvo J, Medendorp S, et al.: Increased complication rates of cardiac rhythm management devices in ESRD patients. Am J Kidney Dis 2007;49:656-663