

Figure 3

| Parameter | | N | Events | Univariate Hazard Ratio (95% CI) | P Value | Multivariate Hazard Ratio (95% CI) | P Value |
|-------------------------|---------------------------|-----|--------|----------------------------------|---------|------------------------------------|---------|
| T Stage | T1/2 | 102 | 14 | | | referent | --- |
| | T3 | 136 | 31 | 1.73 (0.92-3.25) | 0.089 | 1.75 (0.89-3.43) | 0.106 |
| | T4 | 19 | 4 | 1.64 (0.54-4.99) | 0.383 | 2.35 (0.67-8.28) | 0.185 |
| Grade | Poor/Unknown | 40 | 5 | | | referent | --- |
| | Well | 19 | 3 | 1.54 (0.61-3.89) | 0.365 | 0.86 (0.2-3.74) | 0.839 |
| | Moderate | 198 | 41 | 1.07 (0.25-4.46) | 0.931 | 1.1 (0.42-2.86) | 0.850 |
| Location | Rectal | 35 | 6 | | | referent | --- |
| | Right | 108 | 17 | 0.95 (0.38-2.42) | 0.919 | 1.09 (0.40-3.03) | 0.861 |
| | Left | 17 | 4 | 1.63 (0.46-5.80) | 0.448 | 1.52 (0.40-5.86) | 0.541 |
| | Sigmoid | 97 | 22 | 1.55 (0.63-3.83) | 0.343 | 1.81 (0.71-4.60) | 0.215 |
| Lymphovascular Invasion | No | 204 | 43 | | | referent | --- |
| | Yes | 53 | 6 | 0.67 (0.28-1.57) | 0.356 | 0.51 (0.20-1.32) | 0.166 |
| Nodes Harvested | <12 | 45 | 13 | | | referent | --- |
| | >=12 | 212 | 36 | 1.30 (0.70-2.41) | 0.415 | 0.61 (0.31-1.21) | 0.158 |
| Treatment | Surgery | 200 | 35 | | | referent | --- |
| | Surgery + Chemotherapy | 57 | 14 | 0.58 (0.31-1.09) | 0.089 | 1.22 (0.61-2.41) | 0.574 |
| | Occult Metastases | | | | | | |
| | pN0(mol-) | 32 | 2 | | | referent | --- |
| | pN0(mol+) | 225 | 47 | 4.09 (0.99-16.85) | 0.051 | 4.66 (1.11-19.57) | 0.035 |

