

March 2006

Cell-signalling dynamics in time and space

Boris N. Kholodenko PhD, DSci
Thomas Jefferson University

Follow this and additional works at: <https://jdc.jefferson.edu/pacbfp>



Part of the [Medical Cell Biology Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Kholodenko, Boris N. PhD, DSci, "Cell-signalling dynamics in time and space" (2006). *Department of Pathology, Anatomy, and Cell Biology Faculty Papers*. Paper 5.
<https://jdc.jefferson.edu/pacbfp/5>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Pathology, Anatomy, and Cell Biology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

CELL SIGNALLING DYNAMICS IN TIME AND SPACE

Boris N. Kholodenko

Department of Pathology, Anatomy and Cell Biology, Thomas Jefferson University, 1020 Locust St., Philadelphia, PA 19107, USA. E-mail: Boris.Kholodenko@jefferson.edu.

PREFACE.

The specificity of cellular responses to receptor stimulation is encoded by the spatial and temporal dynamics of downstream signalling networks. Computational models provide insights into the intricate relationships between stimuli and responses and reveal mechanisms that enable networks to amplify signals, reduce noise and generate discontinuous bistable dynamics or oscillations. These temporal dynamics are coupled to precipitous spatial gradients of signalling activities, which guide pivotal intracellular processes, but also necessitate mechanisms to facilitate signal propagation across a cell.

INTRODUCTION

Cells respond to a multitude of external cues using a limited number of signalling pathways activated by plasma membrane receptors, such as G protein-coupled receptors (GPCRs) and receptor tyrosine kinases (RTKs). These pathways do not simply transmit, but process, encode and integrate internal and external signals. In recent years, it has become apparent that distinct spatio-temporal activation profiles of the same repertoire of signalling proteins result in different gene activation patterns and diverse physiological responses¹⁻³. Thus, pivotal cellular decisions, such as cytoskeletal reorganization, cell cycle checkpoints and apoptosis (active cell death), depend on the precise temporal control and relative spatial distribution of activated signal-transducers.

Signalling by RTKs has long been in the limelight of scientific interest owing to its central role in the regulation of embryogenesis, cell survival, motility, proliferation, differentiation, glucose metabolism, and apoptosis⁴⁻⁶. Malfunction of RTK signalling is a leading cause of major human diseases that range from developmental defects to cancer, chronic inflammatory syndromes and diabetes⁶⁻⁸. Upon stimulation, RTKs undergo dimerization (for example, the epidermal growth factor (EGF) receptor) or allosteric transitions (insulin receptor) that results in activation of the intrinsic tyrosine kinase^{4,9}. Subsequent phosphorylation of multiple tyrosine residues transmits a biochemical message to a number of cytoplasmic proteins, triggering their mobilization to the cell surface^{4,10}. The

resulting cellular responses occur through complex biochemical circuits of protein interactions and covalent-modification cascades.

An emerging picture of interconnected networks has replaced the earlier view of discrete linear pathways that relate extracellular signals to specific genes, raising questions about the specificity of signal-response events. In fact, the protein complement that mediates signal transduction is similar for all RTK pathways¹¹. Both GPCRs and RTKs activate kinase/phosphatase cascades, such as mitogen-activated protein kinase (MAPK) cascades, that turn on nuclear transcription factors. For any individual receptor pathway, there is no single protein or gene responsible for signalling specificity. Rather, specificity is determined by the temporal and spatial dynamics of downstream signalling components. The classical example is the distinct biological outcome of PC12 cell stimulation with EGF and nerve growth factor (NGF). EGF-induced *transient* MAPK activation results in proliferation, whereas a *sustained* MAPK activation by NGF changes the cell fate and induces differentiation^{1,2}. However, the factors controlling the kinetics of MAPK cascades are intricate. MAPK cascades can generate bistable dynamics (where two stable “On” and “Off” steady states coexist), abrupt switches, and oscillations¹²⁻¹⁴, and their responses depend dramatically on subcellular localization or recruitment to scaffolds^{15,16}.

The purpose of this review is to survey dynamic and spatial aspects of intracellular communication. Wherever possible, I outline general principles by which chemical transformations and Brownian motion of myriad signalling molecules create coordinated behaviour in time and space and generate stimulus-specific responses. I explain how the timing, amplitude and duration of signalling responses are elucidated by exploiting mechanistic systems-level models that help unravel crucial interactions and kinetic factors. Extremely complex dynamic behaviours are shown to arise from simple basic modules, adding to the repertoire of specific signalling outcomes. A number of excellent reviews have focused on computational functions of signalling networks, offering an intriguing glimpse into the parallels between biological and human-made control systems¹⁷⁻²⁰. However, there are important distinctions between electronic and living cell circuitry, which are illustrated by examining the spatial dynamics of intracellular communication. These studies have led to unexpected predictions about the spatial control of intracellular gradients of signalling activities^{21,22} and to the recognition that diffusion alone cannot account for effective propagation of phosphorylation signals that are terminated by phosphatases^{23,24}. The transfer of information over *intracellular* distances of more than a few micrometers requires facilitated transport mechanisms, including movement of phosphorylated kinases on scaffolds and endosomes driven by molecular motors and travelling waves of phosphoproteins^{23,25-28}.

TEMPORAL DYNAMICS OF SIGNALLING NETWORKS.

Mechanistic models can reveal crucial regulations. Since the 1990s, modelling has emerged as a novel tool to handle the rapidly growing information on the molecular parts list and the overwhelmingly complex interaction circuitry of signalling networks²⁹⁻⁴¹. These mechanistic models aspired to create *in silico* replicas of cellular networks with the initial purpose of understanding the temporal dynamics of signalling responses. General principles of model building are illustrated exploiting models of the EGF receptor (EGFR) network (BOX 1). Importantly, EGFR is not only the best studied RTK, but together with other members of the ErbB family plays a pivotal role in carcinogenesis^{7,8,42,43}. Phosphorylation of a number of tyrosine residues on EGFR and binding to, and activation of, EGFR adapter and target proteins (Supplementary Table S1) initiates signal propagation through multiple interacting branches including the phospholipase C- γ (PLC γ), phosphatidylinositol 3-kinase (PI3K)/Akt and extracellular-signal-regulated kinase (ERK)/MAPK pathways (see EGF network diagrams in reference⁴⁴ and at: <http://www.grt.kyushu-u.ac.jp/spad/pathway/egf.html>, <http://www.biocarta.com/pathfiles/egfPathway.asp>). Merely qualitative arguments fall short of providing insights into the complex temporal responses of a variety of downstream EGFR targets, and reliable and testable computational models are required to predict signalling dynamics⁴⁵⁻⁴⁷.

The first mechanistic model of the EGFR network was published in 1999 and explained the temporal dynamics of signalling responses in liver cells stimulated with EGF³⁰. Interrogation of this model generated a number of hypotheses and counterintuitive predictions (BOX 1). A particularly surprising prediction was that EGFR-mediated phosphorylation of the Src homology and collagen domain protein (Shc) would decrease its binding affinity and facilitate Shc dissociation from the receptor. EGFR phosphorylates Shc on Tyr317 located within the central collagen-homology linker region, distant from the N- and C-terminal domains that mediate binding to EGFR. The modular structure underlying protein interactions¹⁰ might imply that phosphorylation of residues outside the Shc binding domains should not influence the affinity. However, molecular dynamics simulations revealed that Tyr317 phosphorylation significantly affects collective motions of Shc domains, increases structural rigidity of the linker region and decreases the flexibility of the binding domains, significantly reducing their capacity to interact with EGFR⁴⁸. These findings corroborated the prediction of the kinetic model³⁰ and favoured a broader view that the affinities of many RTK-binding partners (for example, the p85-subunit of PI3K) might decrease following RTK-mediated phosphorylation. In recent years, a number of EGFR pathway models with predictive and explanatory

power have been developed. These models addressed various aspects of EGFR-mediated signalling, including (1) transient versus sustained responses of the MAPK cascade “gatekeepers” (small GTPases Ras and Rap1) to various growth factors^{32,41,49}, (2) the non-linear dependences of the amplitude of MAPK activation on the EGFR numbers³⁴, (3) autocrine positive-feedback loops⁵⁰, (4) cross-talk between the MAPK and Akt pathways³⁷, and (5) integration of EGFR signalling from the plasma membrane and the endosomes⁵¹.

Challenges in mechanistic modelling. Perhaps, the most significant challenges that face mechanistic modelling are (i) the lack of quantitative kinetic data and (ii) the *combinatorial* increase in the number of emerging distinct species and states of the protein network being simulated^{38,52}. The first challenge is beginning to be addressed by nascent quantitative proteomics of posttranslational modification^{53,54}. The second challenge arises because RTKs and many signalling proteins possess multiple docking sites, serving as *scaffolds* that generate a variety of heterogeneous multi-protein complexes, each involved in multiple parallel reactions. Even initial steps in signal transduction can generate hundreds of thousands of distinct states³⁸, referred to as “micro-states” of a network⁵⁵. Because of the exceedingly high numbers of micro-states, previous models merely ignored this combinatorial variety and simulated only a small part of feasible states and reactions. Several methods of handling this problem have been proposed, all based on specifying rules that automatically generate species and reactions. Programs implementing these methods include StochSim⁵², BioNetGen^{56,57}, and Molecularizer⁵⁸. The entire micro-state network can either be generated in advance for deterministic simulations^{56,57}, or the species and reactions can be generated as needed during a stochastic simulation^{52,57,58}.

An alternative “domain-oriented” approach rigorously simplifies or approximates a mechanistic micro-state picture in terms of “macro-states”, such as the phosphorylation levels and the fractions occupied by binding partners^{55,59}. A necessary prerequisite is the presence of domains/sites that do not allosterically influence each other. This domain-oriented framework drastically reduces the number of states and differential equations to be solved and, therefore, the computational cost of both deterministic and stochastic simulations.

Cycle and cascade motifs. A universal motif found in cellular networks is the cycle that is formed by two or more interconvertible forms of a signalling protein. The protein forms are modified by two opposing enzymes, such as a kinase and phosphatase for phosphoproteins, or a guanine nucleotide exchange factor (GEF) and GTPase-activating protein (GAP) for small G-proteins (FIG.1). Cascades of such cycles form the backbone of most signalling pathways that propagate external stimuli from the membrane to the nucleus or other distant targets. The well-known property of these cycles is

“ultrasensitivity” to input signals, which occurs when the converting enzymes operate near saturation⁶⁰. Depending on the degree of saturation, the response of either interconvertible form ranges from a merely hyperbolic to an extremely steep sigmoidal curve. Sequestration of a signalling protein by converting enzymes significantly decreases sigmoidicity of responses. Likewise, ultrasensitivity can disappear if converting enzymes are inhibited or saturated by their products⁶¹. By contrast, multi-site phosphorylation (following a distributive, multi-collision mechanism⁶²) was shown to increase output-input sensitivity dramatically, resisting the sequestration effect and leading to switch-like responses⁶²⁻⁶⁵. Multi-site protein modification as a variation of the basic cycle motif is repeatedly used in nature, and this has pivotal ramifications for signalling dynamics⁶²⁻⁶⁵.

Feedback loops induce complex dynamics. An increase in the number of interconnecting cycles in a cascade^{66,67} or positive feedback further increases the sensitivity of the target to the input signal. The notion of feedback is one of the most fundamental concepts in biological control. Positive feedback amplifies the signal, whereas negative feedback attenuates it. However, feedback loops not only change steady-state responses, but also favour the occurrence of instabilities. When a steady state becomes unstable, a system can jump to another stable state, start to oscillate or exhibit chaotic behaviour. Positive feedback can cause bistability¹⁴, but also either alone or in combination with negative feedback, it can trigger oscillations, for example, the Ca^{2+} oscillations arising from Ca^{2+} -induced Ca^{2+} release⁴⁶ and the cell cycle oscillations^{68,69}. Such positive-feedback oscillations generally do not have sinusoidal shapes and are referred to as relaxation oscillations, operating in a pulsatory manner: a part of a dynamic system is bistable, and there is a slow process that periodically forces the system to jump between “Off” and “On” states, generating oscillations (BOX 2).

While positive feedback endows signalling cascades with the potential for bistability and relaxation oscillations, negative feedback can bring about adaptation and robustness to parameter variations within the feedback loop (for instance, caused by genetic variability)^{20,70}. Although negative feedback can stabilize the cascade output when demand fluctuates, above certain threshold strength, this feedback induces damped or sustained oscillations. These oscillations are caused by the time delay within the negative feedback loop and require some degree of ultrasensitivity of individual cascade cycles¹². Notably, relaxation oscillations and negative-only feedback oscillations differ in their robustness to noise⁷¹ and generally exhibit different shapes and control of the amplitude and period.

Intricate dynamic properties have been traditionally associated with cascades of cycles^{69,72}, yet even single cycles can exhibit complex dynamics, such as bistability and relaxation oscillations (BOX 2). For instance, multi-site protein modification not only increases ultrasensitivity, but

potentially leads to bistability⁶⁵. The reported kinetic data allow us to suggest that a single MAPK cascade level, e.g., the dual phosphorylation ERK cycle, can exhibit bistability and hysteresis within a certain parameter range⁶⁵; this prediction is awaiting experimental verification. A simple one-site modification cycle can turn into a bistable switch by four different regulatory mechanisms, in which one of the protein forms stimulates its own production or inhibits its consumption, thereby creating a destabilizing control loop (BOX2 and Supplementary Table S3). An extra (stabilizing) feedback loop that affects the rate of synthesis or degradation of a converting enzyme can render this bistable switch into a relaxation oscillator (the resulting 32 distinct feedback designs that can give rise to oscillations are shown in FIG.2 and Supplementary FIG.S1).

Cascade dynamics govern cellular functions. Following stimulation, signalling proteins become involved in collective dynamic behaviour that none of the individual molecules can exhibit in isolation. Inherently complex dynamics of universal signalling motifs allows a cascade of these motifs to generate even large spectrum of temporal patterns that contribute to the signal-response specificity. A multitude of negative and positive feedback loops enables cascades to generate gradual and ultrasensitive responses, multi-stability and oscillations^{12,18,20,46,47,72-74}. For instance, the same basic architecture allows MAPK cascades to operate as negative feedback amplifiers that reduce noise, as ultrasensitive or discontinuous switches, or flexible integration modules; these theoretical predictions were verified experimentally^{13,16,72}.

The signalling dynamics can become multi-stable, when two or more bistable cycles form a cascade, such as MAPK cascade⁶⁵. The biological outcome of multi-stability is the ability to control multiple irreversible transitions, for instance, sequential transitions in the cell cycle. Central components of the cell cycle machinery are cyclin-dependent kinases (such as, CDK1/Cdc2), which sequential activation/inactivation governs cell-cycle transitions. CDK1/Cdc2 activity is low (OFF) in G1-phase (resting state) and has to be high (ON) for entry into mitosis (M-phase). Recently, hysteresis and bistability were shown to occur in the activation/inactivation of CDK1/Cdc2, confirming a theoretical prediction made by Novak and Tyson a decade ago⁷⁵. Bistability in the CDK1/Cdc2 cycle arises from positive and double-negative feedback loops in the reactions, where CDK1/Cdc2 activates its activator (the phosphatase Cdc25) and inactivates its inhibitor (the kinases Wee1 and Myt1). Negative feedback from the anaphase promoting complex (APC) renders the CDK1/Cdc2 bistable switch into a relaxation oscillator that drives the cell cycle^{68,69}. Intriguingly, Cdc25 and Wee1 themselves can be phosphorylated on multiple sites and therefore can potentially exhibit bistability, implying that the entire CDK/cyclin system can display multiple steady states⁶⁵ (this prediction is awaiting experimental verification). Sequential bifurcations of multiple steady

states provide more flexibility in the control of the cell fate and allow for a number of check points in the cell cycle.

SPATIAL DIMENSION OF SIGNALLING NETWORKS

Activation of cell-surface receptors and their downstream targets leads to the spatial relocation of multiple proteins within the cell. During evolution, cells have developed not only means to control the temporal dynamics of signalling networks, but also mechanisms for precise spatial sensing of the relative localization of signalling proteins. The regulation of signalling within the cellular space is pivotal for a number of physiological processes, such as cell division, motility and migration. Here I show how basic principles of the control of reaction rates, diffusive movement and directed transport underlie sophisticated mechanisms that activate signalling routes by the membrane recruitment of binding partners, provide spatial cues for cell division and transmit signals to distant cellular targets.

Regulation of signalling by membrane recruitment. Receptor stimulation triggers the mobilization of cytosolic adaptor proteins and enzymes to cellular membranes. Subsequent phosphorylation results in the assembly of signalling complexes on receptors, scaffolds and cytoskeletal elements⁷⁶. These spatial relocations are effective control mechanisms of switching-on signalling pathways⁷⁷. The classical example is the control of the Ras/MAPK cascade by membrane recruitment of SOS and RasGAP (GEF and GAP for the small GTPase Ras, respectively), mediated by RTKs (for example, by EGFR) and membrane-bound scaffolds. It has been previously proposed that the role of this recruitment is to increase diffusion-limited (first-encounter) rates, but it was recently shown that the function of membrane localization is to amplify the number of complexes that are formed between the signalling partners⁷⁶⁻⁷⁸. SOS and RasGAP bound to EGFR are confined to a small volume near the membrane that results in a $10^2 - 10^3$ -fold increase in the apparent affinity of these catalysts for Ras. Simulations corroborate the theory, demonstrating that in the absence of the membrane recruitment, the cytoplasmic concentrations of SOS and RasGAP would have to increase $10^2 - 10^3$ -fold to account for the observed rates of Ras activation/deactivation⁴⁹. We conclude that the spatial organization of the Ras circuit is crucial for the effective control of Ras activity.

Location determines signalling outputs. The localization of signalling proteins to distinct subcellular regions, such as internal membranes and membrane microenvironments (including lipid rafts) modulates signalling outputs^{76,79}. Specific anchoring subunits direct catalytic subunits of kinases and phosphatases, such as cAMP-dependent protein kinase, protein kinase C and serine/threonine protein phosphatases PP1, PP2A and PP2B to different cellular regions⁸⁰. The general mechanism is to orient broad-specificity enzymes towards specific targets and physically

separate them from undesirable substrates. Discrete subcellular distribution enhances the specificity and fidelity of phosphorylation and dephosphorylation catalyzed by these kinases and phosphatases.

Qualitatively different patterns of signalling are generated by receptors and downstream effectors associated with endosomes or the plasma membrane⁸¹. Likewise, the same protein cascades operate in surprisingly dissimilar ways when localized to different cellular compartments. The input-output sensitivity of MAPK cascade is different for signalling from the plasma membrane, the Golgi apparatus and endosomes^{16,82}. Computational models have yet to take into full account the ramifications of subcellular localization on signalling outcomes.

Spatial gradients of signalling activities. In the late 1990s, the novel concept of protein activity gradients within a cell was proposed^{21,83-85}. This concept has matured in recent years, when fluorescence resonance energy transfer-based biosensors enabled discoveries of intracellular gradients of the active form of the small GTPase Ran⁸⁶ and the phosphorylated form of stathmin-oncoprotein 18 (Op18/stathmin) that regulates the microtubule polymerization⁸⁷. Spatial gradients of protein activities organize signalling around cellular structures, such as membranes, chromosomes and scaffolds, and provide positional cues for key processes, including cell division. During mitosis, the microtubule network changes from the radial architecture emanating from the centrosome to a bipolar spindle. How this remarkable rearrangement occurs is not completely understood. Spatial gradients of several molecules that influence microtubule dynamics, including Op18/stathmin and RanGTP which interacts with the nuclear-transport receptor importin- β , were recently suggested to guide microtubule-kinetochore positioning during the mitotic-spindle assembly⁸⁸⁻⁹¹.

The basic prerequisite for signalling gradients is the spatial segregation of opposing reactions (for instance, kinase and phosphatase) in a universal protein-modification cycle (FIG.3). For a protein phosphorylated by a membrane-bound kinase and dephosphorylated by a cytosolic phosphatase, Brown & Kholodenko predicted that there can be a gradient of the phosphorylated protein (BOX 3) - high concentration close to the membrane and low concentration within the cell²¹. Given measured values of protein diffusivity and kinase and phosphatase activities, it was estimated that phosphoprotein gradients might be large within the intracellular space (Fig. 3C). Even in small bacterial cells, spatial gradients of this kind were recently suggested for chemotaxis proteins^{22,92}. Interestingly, the existence of intracellular cAMP-gradients generated by the membrane-associated adenylate cyclase and cytosolic phosphodiesterase was conjectured theoretically in 1980⁹³.

For a simple cycle of two opposing enzymes, one confined to a cellular structure and the other in the cytoplasm, the characteristic size of the gradient is determined predominantly by the protein

diffusivity and the activity of the cytoplasmic enzyme (BOX 3). If the enzyme in the cytoplasm is saturated, the gradient length also depends on the activity of the enzyme confined to the structure²³. Cells employ additional means to control the shape and the extent of spatial gradients. For instance, if the input activity changes gradually in space, ultrasensitive responses would generate precipitous, short length gradients, whereas linear responses can generate shallow output gradients. Kinase and GTPase cascades can expand gradients over large spatial regions (BOX 4). How far gradients of the active form reach into the cytoplasm can also be controlled by the association with an adaptor protein that protects against the deactivating enzyme. In fact, the complex of RanGTP with importin- β generates more extended gradients than RanGTP alone because the GTP hydrolysis by RanGAP is prevented during the life-time of the RanGTP-importin- β complex⁹¹. Alternatively, binding an adaptor that enhances the deactivation rate will decrease the length of the gradient.

Temporal signalling dynamics can induce spatial gradients. The time-course of signalling responses and the formation of spatial gradients of signalling activities are directly related. The spatial segregation of opposing enzymes is often initiated by specific signals. For instance, the cyclin-dependent kinase CDK1/Cdc2 phosphorylates the nuclear localization signal (NLS) of RCC1, which is a GEF for Ran⁹⁴. In the G1-phase CDK1/Cdc2 activity is low and, therefore, the NLS phosphorylation level is low. CDK1/Cdc2 becomes active to drive cells into mitosis. Owing to NLS phosphorylation by CDK1/Cdc2, RCC1 binds to the chromosomes and catalyzes the conversion of inactive RanGDP into active RanGTP. Since the opposing RanGAP activity is predominantly cytoplasmic, spatial gradients of active Ran emerge with high RanGTP concentration near mitotic chromosomes and low in the surrounding area. Therefore, the RanGTP-related gradients that guide the mitotic-spindle self-organization⁹¹ are driven by the temporal dynamics of CDK1/Cdc2.

FACILITATED COMMUNICATION WITHIN CELLS.

Phosphoprotein gradients in MAPK cascades. Phosphoprotein gradients are hallmarks of kinase/phosphatase cascades, including MAPK cascades. MAPK cascades contain three interconnected cycles of MAPK, MAPK kinase (MAPKK) and MAPKK kinase (MAPKKK). In the MAPK/ERK cascade (the most well-characterized biochemically), these kinases are ERK, MEK and Raf. Upon RTK stimulation and Ras activation, the cytosolic Raf is recruited to the cell membrane, where it binds to and phosphorylates MEK on two serine residues. Phosphorylated MEK drifts into the cell interior, where it phosphorylates ERK on threonine and tyrosine residues. Because MEK is dephosphorylated in the cytoplasm, spatial gradients of phosphorylated MEK, and subsequently

phosphorylated ERK might occur. Calculations show that these gradients can be precipitous^{23,24}, decreasing the strength of the phosphorylation signal to the nucleus. Instructively, the phosphorylation signal reaches further into the cell if the cascade has more levels, and this might be one of the reasons that cascades exist (BOX 4). The cascades found in eukaryotes tend to have more levels than the cascades in prokaryotes; this can be related to larger distances of signal propagation in eukaryotes.

Endocytosis and signalling: a marriage of convenience. Many cytosolic proteins that are phosphorylated at the plasma membrane travel into the cell interior to interact with the targets, which are confined to mitochondria, cytoskeleton, the Golgi, and the nucleus. The membrane confinement of kinase activity and the cytosolic localization of phosphatases can result in unfavorable gradients of phosphorylated signal-transducers provided they spread solely by diffusion; this would impede information transfer^{21,23}. In view of this problem, alternative mechanisms to relay stimuli from the surface to distant targets were proposed including trafficking of phosphorylated kinases with endosomes (“signalling endosome”) or non-vesicular signalling complexes driven by molecular motors^{23,26,28,95,96}. Motor-mediated movement of the endosomes and kinase complexes along microtubules is remarkably distinct from chaotic diffusive motion and is able to prevent the formation of precipitous reaction-diffusion gradients^{23,26}. Although in the past, endocytosis was thought to be a mechanism to attenuate signalling, a dual role of endocytosis is now emerging: a robust, immediate signal transducer on a short time-scale and a downregulator of receptor signalling on longer times^{23,95}. Distinct endocytic compartments, including clathrin, caveolae, and Rab domains, can deliver differential sets of proteins to diverse cellular targets, generating specific signalling outputs^{95,97}.

Retrograde transport and phosphoprotein waves. An interesting puzzle in neurobiology concerns the mechanisms used by neurons to transfer signals over long distances. The survival of developing neurons depends on neurotrophins, such as the nerve growth factor (NGF) and its receptor, TrkA. NGF is produced by peripheral tissues and binds to TrkA on distal axons that are located as far as one meter away from the neuronal soma. How do survival signals reach the cell body in a physiologically relevant span of time? Diffusion is ruled out as a mechanism of long-range signalling, because it would be prohibitively slow. In fact, we have seen that diffusion may be insufficient even for spreading signals across the cytoplasm of large cells, such as *Xenopus* eggs^{23,26}.

Retrograde transport of endosomes containing the NGF-TrkA complexes is critical for neuronal survival^{96,98}. Yet, recent evidence indicates that survival signals can also be transmitted by NGF-independent mechanisms⁹⁹. These might include lateral waves of receptor activation propagating

along the axon membrane²⁵⁻²⁷ and movement of a signalling complex of phosphorylated ERK with intermediate filament vimentin and importin, driven by the molecular motor dynein²⁸. However, lateral propagation of TrkA activation can be excluded, as nearly complete inhibition of TrkA in the cell bodies/proximal axons did not affect survival, whereas TrkA inhibition at distal axons induced apoptosis¹⁰⁰. Although transport of phosphorylated kinases driven by molecular motors is a robust mechanism of retrograde signalling, it cannot account for the initial burst of tyrosine phosphorylation, which reaches neuron bodies as early as several minutes after NGF stimulation¹⁰⁰. This initial rapid signal cannot be carried out by molecular motors that move at 1 to 10 $\mu\text{m}/\text{sec}$ ^{96,101}. It is feasible that the first survival signals are transmitted by waves of protein phosphorylation emerging from kinase/phosphatase cascades, such as MAPK or PI3K cascades, or GEF/GAP cascades of G-protein activation (BOX 4).

Outlook/Future directions

Quantitative models that generate novel experimentally testable hypotheses will have an increasingly important role in postgenomic biology. Future models will integrate data on the distinct spatio-temporal dynamics of signalling from different cellular compartments and provide new insight into the connection between external stimuli and the signalling outcome in terms of gene expression responses. Challenges of the combinatorial complexity of signalling networks and experimental uncertainty in parameter values will be addressed by modular approaches, stochastic and pattern-oriented modelling. The goal of the pattern-oriented approach is to predict and explain dynamic patterns of cellular responses to a multitude of external cues and perturbations. An exceedingly large number of quantitative and also qualitative data patterns will facilitate the verification of the proposed molecular mechanisms and exclude models that are too simplistic and uncertain¹⁰². These systems-level, data-driven models will generate new knowledge and provide strategies for the regulation of the cellular machinery. Understanding mechanisms underlying signalling network function will provide breakthroughs in the identification of critical controlling factors that will be targets for pharmacological interventions in the treatment of major human diseases.

BOX 1. Mechanistic models: keeping track of molecular processes.

The temporal dynamics of signalling networks is described by ordinary differential equations, which are known as chemical kinetics equations¹⁰³ and are derived similarly to the Michaelis-Menten equation, familiar to any biologist. The derivation begins with listing all chemical transformations thereby providing a kinetic scheme of a pathway. Figure, part A shows a simplified scheme of the signalling routes emanating from the epidermal growth factor receptor (EGFR), including the Shc, Grb2-SOS, GAP and phospholipase C- γ (PLC γ) signalling branches, and the RasGDP/RasGTP circuit (phosphorylated proteins are indicated by added “P” after protein name, R is EGFR, molecules within complexes are abbreviated, and the designations are given in Supplementary Table S2). The scheme is translated into differential equations, one for each time dependent molecular species. The rate of the concentration change is the sum of the reaction rates producing a given species minus the sum of consuming rates. Numerical integration (simulations) gives the time course of the concentrations (see figure, parts B and C where EGF-induced responses were simulated using the scheme shown in part A^{49,104}).

Comparison of simulations with data helps generate novel hypotheses and often instigates an overhaul of a model. Data obtained from isolated hepatocytes (black squares and red triangles, see figure, parts B and C) demonstrate that despite the constant level of EGF (10nM), phosphorylation of EGFR and PLC γ is markedly transient (with the peaks reached within the first 15 seconds and the low pseudo-stationary levels within few minutes), whereas phosphorylation of Shc increased almost monotonically^{30,105}. The mechanistic model elucidates that the transient time-course of EGFR phosphorylation arises from the protection of phosphotyrosine residues against phosphatases, whilst these residues are occupied by an adaptor/target protein. Transient patterns of tyrosine phosphorylation of PLC γ are explained by the slow dissociation of the PLC γ -phosphatase complex^{30,104}. In fact, the existence of such complexes was reported¹⁰⁶. Hypotheses generated by computational models have a certainty and precision, furthering our understanding of signalling dynamics. A variety of software tools can assist in quantitative modelling^{56,58,107 109}, and several databases of biological models have been developed, offering an interesting environment to generate and test novel hypotheses by using a computer keyboard¹¹⁰⁻¹¹².

BOX 2. Complex temporal dynamics in a nutshell.

Exotic dynamics emerges from simple, basic signalling motifs. Known examples include bistability arising from multi-site phosphorylation in a single protein cycle or from positive or double-negative feedback in a two-cycle cascade, and negative-feedback oscillations in a cascade with at least three cycles^{12,18,65}. Here I show two additional basic signalling modules that bring about bistable and oscillatory dynamics. A single-site phosphorylation cycle generates only ultrasensitive, but not discontinuous switches. Yet, positive feedback from the phosphorylated form (M_p) to its kinase can render this cycle into a bistable switch. Assuming the kinase (v_{kin}) and phosphatase (v_{phos}) rates follow Michaelis-Menten kinetics (including the activation term), this system is described by a remarkably simple equation,

$$\frac{dM_p}{dt} = v_{kin} - v_{phos} = \frac{k_{kin}^{cat} E_{kin} M}{(K_{m1} + M)} \cdot \frac{(1 + AM_p / K_a)}{(1 + M_p / K_a)} - \frac{k_{phos}^{cat} E_{phos} M_p}{(K_{m2} + M_p)}, \quad M = M^{tot} - M_p \quad (1)$$

Here, the products of the catalytic constants and enzyme concentrations $k_{kin}^{cat} E_{kin}$ and $k_{phos}^{cat} E_{phos}$ are the maximal rates of the kinase and phosphatase, K_{m1}, K_{m2}, K_a and A are kinetic constants, M^{tot} is the protein abundance. In a wide parameter range, there are three distinct solutions to the steady-state relationship $v_{kin} = v_{phos}$. The low and high M_p concentrations correspond to stable ‘‘Off’’ and ‘‘On’’ states, whereas the intermediate state is unstable. The steady-state M_p dependence on the input kinase (phosphatase) activity (known as one-dimensional bifurcation diagram) displays hysteresis, the hallmark of bistability (figure, part A). Likewise, phosphatase rate inhibition by M_p , phosphatase rate activation or kinase rate inhibition by M can produce a similar bistable switch (Supplementary Table S3).

If, in addition, the phosphorylated form M_p inhibits transcription/translation of the kinase protein or promotes its degradation (Supplementary Eqs.S2-S3 and Table S3), thereby creating negative feedback, the universal cycle becomes a relaxation oscillator (figure, part B),

$$\frac{dE_{kin}}{dt} = v_{kin}^{synth} - v_{kin}^{deg} = V_{kin}^0 \frac{(1 + M_p / K_I)}{(1 + I \cdot M_p / K_I)} - k_{kin}^{deg} E_{kin}, \quad I > 1. \quad (2)$$

Likewise, alternative negative feedback design where M_p activates the phosphatase protein transcription/translation (or inhibits its degradation) also generates relaxation oscillations (figure, part C),

$$\frac{dE_{phos}}{dt} = v_{phos}^{synth} - v_{phos}^{deg} = V_{phos}^0 \frac{(1 + A_{dp} M_p / K_d)}{(1 + M_p / K_d)} - k_{phos}^{deg} E_{phos}, \quad A_{dp} > 1 \quad (3)$$

This negative feedback might result from changes in mRNA or protein turnover, immediate-early gene expression, or de novo synthesis of transcription factors regulating protein levels. Different scenarios will correspond to different time scales and affect the period and shape of oscillations.

BOX 3. When do the spatial activity gradients occur?

Here I show how intracellular signalling gradients arise from chemical transformation and diffusion. For a spatially-confined kinase and homogeneously-distributed phosphatase (or a similar enzyme pair, GEF and GAP), the spatio-temporal dynamics of the phosphorylated form c_p of the interconvertible protein is governed by the reaction-diffusion equation,

$$\frac{\partial c_p}{\partial t} = D\Delta c_p - v_p(c_p) \quad (1).$$

When the diffusivities D are equal for the phosphorylated c_p and unphosphorylated c_u forms, their total concentration is constant across the cell, $c_p + c_u = C_{tot}$ (which is untrue for different diffusivities⁸³).

The simplest one-dimensional geometry corresponds to a cylindrical (for instance, bacterial) cell of the length L with the kinase localized to one pole (surface-reaction with rate v_{kin}^{mem} at $x=0$) and the cytoplasmic phosphatase (rate v_p). The steady-state spatial profile $c_p(x)$ is determined by letting the time-derivative in Eq.1 equal zero and imposing the following boundary conditions (the diffusive flux equals v_{kin}^{mem} at the kinase pole and zero at the opposite pole),

$$D \frac{d^2 c_p}{dx^2} - v_p = 0, \quad -D \left. \frac{dc_p}{dx} \right|_{x=0} = v_{kin}^{mem}, \quad \left. \frac{dc_p}{dx} \right|_{x=L} = 0 \quad (2)$$

When the phosphatase is far from saturation, $v_p = k_p c_p$ ($k_p = V_{max}/K_m$ is the observed first-order rate constant), the analytical solution to Eq.2 reads,

$$c_p(x) = c_p(0) \left(\frac{e^{\alpha x} + e^{2L\alpha} e^{-\alpha x}}{1 + e^{2\alpha L}} \right), \quad \alpha^2 = \frac{k_p}{D} \quad (3)$$

When $\alpha L \ll 1$, the phospho-protein concentration decreases almost linearly, and when $\alpha L \geq 1$, it decreases nearly exponentially $c_p(x)/c_p(0) \approx e^{-\alpha x}$ with distance x from the membrane. This provides a simple, but powerful criterion⁸³ that large phospho-protein gradients exist when the

dephosphorylation time $1/k_p$ is smaller than the diffusion time L^2/D . The kinase activity only influences the concentration $c_p(0)$ near the membrane^{21,83}.

Spherical symmetry simplifies analysis of signalling in three dimensions. For a cell of the radius L with a kinase located on the cell surface and a phosphatase in the cytoplasm (FIG.3B), the steady-state phospho-protein concentration decreases from the membrane towards the cell centre nearly exponentially if $\alpha L \geq 1$, FIG. 3C²¹.

A similar exponential decrease in the phosphorylation signal $c_p(r)$ may occur when a kinase is bound to a supra-molecular structure (of radius s) and a phosphatase resides in the surrounding area (of the radius L , FIG.3A). Assuming spherical symmetry, the steady-state concentration $c_p(r)$ is determined by (Supplementary FIG.S2),

$$\begin{aligned} \frac{D}{r^2} \frac{d}{dr} \left(r^2 \frac{dc_p}{dr} \right) - k_p c_p &= 0, \quad -D \frac{dc_p}{dr} \Big|_{r=s} = v_{kin}^{mem}, \quad \frac{dc_p}{dr} \Big|_{r=L} = 0 \\ c_p(r) &= c_p(s) \frac{se^{-\alpha r}}{re^{-\alpha s}} \left(\frac{e^{2\alpha r}(\alpha L + 1) + e^{2\alpha L}(\alpha L - 1)}{e^{2\alpha s}(\alpha L + 1) + e^{2\alpha L}(\alpha L - 1)} \right), \quad \alpha^2 = \frac{k_p}{D} \end{aligned} \quad (4)$$

Considering how $c_p(r)$ decreases for different values of αL , we conclude that signalling gradients cannot be built merely by diffusion, but require the spatial segregation of opposing enzymes.

BOX 4. Facilitated communication through kinase cascades.

For a cascade where a kinase (M_i) at each level activates a kinase (M_{i+1}) at the subsequent downstream level, the gradients of the phosphorylated forms become shallower down the cascade²¹. For a spherical cell, where a membrane-bound kinase phosphorylates M_1 (rate v_{kin}^{mem}) and all other kinases and phosphatases diffuse in the cytoplasm, the spatio-temporal dynamics of a three-level cascade (Fig. 1C) is described by the following equations,

$$\begin{aligned} \frac{\partial M_1 P}{\partial t} &= \frac{D}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial M_1 P}{\partial r} \right) - v_p^1(M_1 P), \quad \frac{\partial M_i P}{\partial t} = \frac{D}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial M_i P}{\partial r} \right) + M_{i-1} P \cdot w_{kin}^i(M_i P) - v_p^i(M_i P), \quad i = 2, 3 \\ D \frac{\partial M_1 P}{\partial r} \Big|_{x=L} &= v_{kin}^{mem}, \quad \frac{\partial M_i P}{\partial r} \Big|_{x=0} = 0, \quad i = 1, 2, 3 \quad M_i + M_i P = M_i^{Tot} = const \end{aligned} \quad (1)$$

Here $M_{i-1} P \cdot w_{kin}^i(M_i P)$ is the rate of phosphorylation of M_i (proportional to the concentration $M_{i-1} P$ of the active kinase upstream) and $v_p^i(M_i P)$ is the rate of the i -th phosphatases (Supplementary

Table S4). The calculated steady-state spatial profiles (figure, part A) show that descending down the cascade, phosphorylated kinases reach further into the cell.

Although the existence of more levels in a cascade facilitates signal transfer over several micrometers, signalling over longer distances (for instance, from the plasma membrane to the nucleus in large cells, such as *Xenopus* eggs) requires additional means, such as vesicular or non-vesicular transport of phosphorylated kinases along microtubules and travelling waves of protein phosphorylation. Such waves propagating through bistable protein modification cascades (Supplementary FIG.S3) were recently predicted²⁶. In fact, travelling waves in bistable systems are well-known in physics, chemistry and biology²⁷. Bistability intrinsic to the multisite activation/deactivation cycles in the cytoplasm⁶⁵ (such as the MAPKK or MAPK cycles) gives rise to travelling waves that propagate binary phosphorylation signals to distant targets. Figure, part B shows the travelling phosphorylation wave that propagates through a three-level cascade following a 30-second pulse of activity of the input membrane-bound kinase (V_{kin}^{mem} , Supplementary Table S5). Additional positive feedback in the cytoplasm may enable phosphorylation waves to propagate with high velocity over exceedingly long distances, possibly solving a long-standing enigma of survival signalling in neurons.

References

1. Marshall, C. J. Specificity of receptor tyrosine kinase signaling: transient versus sustained extracellular signal-regulated kinase activation. *Cell* **80**, 179-185 (1995).
2. Murphy, L. O., Smith, S., Chen, R. H., Fingar, D. C. & Blenis, J. Molecular interpretation of ERK signal duration by immediate early gene products. *Nat. Cell Biol.* **4**, 556-564 (2002).
3. Hoffmann, A., Levchenko, A., Scott, M. L. & Baltimore, D. The I κ B-NF- κ B signaling module: temporal control and selective gene activation. *Science* **298**, 1241-1245 (2002).
4. Schlessinger, J. Cell signaling by receptor tyrosine kinases. *Cell* **103**, 211-225. (2000).
5. Hunter, T. Signaling--2000 and beyond. *Cell* **100**, 113-127. (2000).
6. Gray, S. G., Stenfeldt Mathiasen, I. & De Meyts, P. The insulin-like growth factors and insulin-signalling systems: an appealing target for breast cancer therapy? *Horm. Metab. Res.* **35**, 857-871 (2003).
Presented the domain organisation of a variety of receptor tyrosine kinases, including those implicated in human malignancies.
7. Yarden, Y. & Sliwkowski, M. X. Untangling the ErbB signalling network. *Nat. Rev. Mol. Cell Biol.* **2**, 127-137 (2001).
8. Fischer, O. M., Hart, S., Gschwind, A. & Ullrich, A. EGFR signal transactivation in cancer cells. *Biochem. Soc. Trans.* **31**, 1203-1208 (2003).
9. De Meyts, P. & Whittaker, J. Structural biology of insulin and IGF1 receptors: implications for drug design. *Nat. Rev. Drug Discov.* **1**, 769-783 (2002).
10. Pawson, T. & Nash, P. Assembly of cell regulatory systems through protein interaction domains. *Science* **300**, 445-452 (2003).
11. Schlessinger, J. Common and distinct elements in cellular signaling via EGF and FGF receptors. *Science* **306**, 1506-1507 (2004).
12. Kholodenko, B. N. Negative feedback and ultrasensitivity can bring about oscillations in the mitogen-activated protein kinase cascades. *Eur. J. Biochem.* **267**, 1583-1588 (2000).
13. Bhalla, U. S., Ram, P. T. & Iyengar, R. MAP kinase phosphatase as a locus of flexibility in a mitogen-activated protein kinase signaling network. *Science* **297**, 1018-1023. (2002).
14. Xiong, W. & Ferrell, J. E., Jr. A positive-feedback-based bistable 'memory module' that governs a cell fate decision. *Nature* **426**, 460-465 (2003).
15. Whitehurst, A., Cobb, M. H. & White, M. A. Stimulus-coupled spatial restriction of extracellular signal-regulated kinase 1/2 activity contributes to the specificity of signal-response pathways. *Mol. Cell. Biol.* **24**, 10145-10150 (2004).
16. Harding, A., Tian, T., Westbury, E., Frische, E. & Hancock, J. F. Subcellular localization determines MAP kinase signal output. *Curr. Biol.* **15**, 869-873 (2005).
Elegant experimental demonstration that in mammalian cells the MAPK cascade can operate as a switch with different sensitivity to the input signals from the plasma membrane or from the cytoplasm. Positive feedback loops that might encircle the MEK/ERK module in different cell types are not required for the switch-like behavior.
17. Bray, D. Protein molecules as computational elements in living cells. *Nature* **376**, 307-312 (1995).
18. Tyson, J. J., Chen, K. C. & Novak, B. Sniffers, buzzers, toggles and blinkers: dynamics of regulatory and signaling pathways in the cell. *Current Opinion in Cell Biology* **15**, 221-231 (2003).
This thought-provoking and influential review provides memorable examples of elementary signalling modules that generate linear, hyperbolic and sigmoidal responses and complex dynamic behaviours.

19. Wolf, D. M. & Arkin, A. P. Motifs, modules and games in bacteria. *Curr. Opin. Microbiol.* **6**, 125-134 (2003).
20. Sauro, H. M. & Kholodenko, B. N. Quantitative analysis of signaling networks. *Prog. Biophys. Mol. Biol.* **86**, 5-43 (2004).
21. Brown, G. C. & Kholodenko, B. N. Spatial gradients of cellular phospho-proteins. *FEBS Lett.* **457**, 452-454 (1999).
First theoretical analysis of protein activity gradients arising from the spatial separation of opposing enzymes in a protein-modification cycle. Together with references 23 and 83, criteria for the existence of these gradients within a cell and estimates of their characteristic sizes were formulated.
22. Lipkow, K., Andrews, S. S. & Bray, D. Simulated diffusion of phosphorylated CheY through the cytoplasm of Escherichia coli. *J. Bacteriol.* **187**, 45-53 (2005).
23. Kholodenko, B. N. MAP kinase cascade signaling and endocytic trafficking: a marriage of convenience? *Trends Cell Biol.* **12**, 173-177. (2002).
Demonstrated that the propagation of phosphorylation signals solely by diffusion can be terminated by cytoplasmic phosphatases. Motor-driven trafficking of endosomes and scaffolds carrying phosphorylated kinases/signalling complexes can be required for the transfer of signals, especially across large cells.
24. Maly, I. V., Wiley, H. S. & Lauffenburger, D. A. Self-organization of polarized cell signaling via autocrine circuits: computational model analysis. *Biophys. J.* **86**, 10-22 (2004).
25. Reynolds, A. R., Tischer, C., Vermeer, P. J., Rocks, O. & Bastiaens, P. I. EGFR activation coupled to inhibition of tyrosine phosphatases causes lateral signal propagation. *Nat. Cell Biol.* **5**, 447-453 (2003).
Using single cell live monitoring, the lateral propagation of EGFR activation was demonstrated and explained by a membrane wave of receptor phosphorylation triggered by bistability emerging in the EGFR activation/deactivation network. Bistability is thought to arise from inhibition of protein phosphatases by EGFR-mediated production of reactive oxygen species (reviewed in reference 27).
26. Kholodenko, B. N. Four-dimensional organization of protein kinase signaling cascades: the roles of diffusion, endocytosis and molecular motors. *J. Exp. Biol.* **206**, 2073-2082 (2003).
27. Tischer, C. & Bastiaens, P. I. Lateral phosphorylation propagation: an aspect of feedback signalling? *Nat. Rev. Mol. Cell Biol.* **4**, 971-974 (2003).
28. Perlson, E. et al. Vimentin-dependent spatial translocation of an activated MAP kinase in injured nerve. *Neuron* **45**, 715-726 (2005).
First experimental demonstration that axonal transport of phosphorylated kinases over long distances may not necessarily involve endocytic vesicles. Rather in lesioned nerve, injury-signalling complexes containing ppErk, vimentin and importin, are driven by the molecular motor dynein along the axon.
29. Bhalla, U. S. & Iyengar, R. Emergent properties of networks of biological signaling pathways. *Science* **283**, 381-387 (1999).
Pioneering paper that presented simplified dynamic models for a number of signalling pathways and demonstrated that they can exhibit collective dynamic behaviours that result in signal integration, modulation and amplification, bistability and hysteresis.
30. Kholodenko, B. N., Demin, O. V., Moehren, G. & Hoek, J. B. Quantification of short term signaling by the epidermal growth factor receptor. *J. Biol. Chem.* **274**, 30169-30181 (1999).
31. Haugh, J. M., Wells, A. & Lauffenburger, D. A. Mathematical modeling of epidermal growth factor receptor signaling through the phospholipase C pathway: mechanistic insights and predictions for molecular interventions. *Biotechnol. Bioeng.* **70**, 225-238. (2000).

32. Brightman, F. A. & Fell, D. A. Differential feedback regulation of the MAPK cascade underlies the quantitative differences in EGF and NGF signalling in PC12 cells. *FEBS Lett.* **482**, 169-174 (2000).
33. Asthagiri, A. R. & Lauffenburger, D. A. A computational study of feedback effects on signal dynamics in a mitogen-activated protein kinase (mapk) pathway model. *Biotechnol. Prog.* **17**, 227-239 (2001).
34. Schoeberl, B., Eichler-Jonsson, C., Gilles, E. D. & Muller, G. Computational modeling of the dynamics of the MAP kinase cascade activated by surface and internalized EGF receptors. *Nat. Biotechnol.* **20**, 370-375 (2002).
35. Heinrich, R., Neel, B. G. & Rapoport, T. A. Mathematical models of protein kinase signal transduction. *Mol. Cell.* **9**, 957-970 (2002).
36. Woolf, P. J. & Linderman, J. J. Untangling ligand induced activation and desensitization of G-protein-coupled receptors. *Biophys. J.* **84**, 3-13 (2003).
37. Hatakeyama, M. et al. A computational model on the modulation of mitogen-activated protein kinase (MAPK) and Akt pathways in heregulin-induced ErbB signalling. *Biochem. J.* **373**, 451-463 (2003).
38. Goldstein, B., Faeder, J. R. & Hlavacek, W. S. Mathematical and computational models of immune-receptor signalling. *Nat. Rev. Immunol.* **4**, 445-456 (2004).
An insightful account of intricacies involved in computational modelling of immune-receptor signalling including the combinatorial complexity of states and processes in highly interconnected networks.
39. Bornheimer, S. J., Maurya, M. R., Farquhar, M. G. & Subramaniam, S. Computational modeling reveals how interplay between components of a GTPase-cycle module regulates signal transduction. *Proc. Natl. Acad. Sci. U S A* **101**, 15899-15904 (2004).
40. Hornberg, J. J. et al. Principles behind the multifarious control of signal transduction. ERK phosphorylation and kinase/phosphatase control. *Febs J.* **272**, 244-258 (2005).
41. Sasagawa, S., Ozaki, Y., Fujita, K. & Kuroda, S. Prediction and validation of the distinct dynamics of transient and sustained ERK activation. *Nat. Cell Biol.* **7**, 365-373 (2005).
42. Wiley, H. S., Shvartsman, S. Y. & Lauffenburger, D. A. Computational modeling of the EGF-receptor system: a paradigm for systems biology. *Trends Cell Biol.* **13**, 43-50. (2003).
43. Hendriks, B. S., Orr, G., Wells, A., Wiley, H. S. & Lauffenburger, D. A. Parsing ERK activation reveals quantitatively equivalent contributions from epidermal growth factor receptor and HER2 in human mammary epithelial cells. *J. Biol. Chem.* **280**, 6157-6169 (2005).
44. Oda, K., Matsuoka, Y., Funahashi, A. & Kitano, H. A comprehensive pathway map of epidermal growth factor receptor signaling. *Mol. Syst. Biol.* doi: **10.1038/msb4100014** (2005).
45. Bray, D. Reductionism for biochemists: how to survive the protein jungle. *Trends Biochem. Sci.* **22**, 325-326 (1997).
Short and revealing comment on reductionism in biology, arguing that only computational approaches may enable us to see “the forest for the trees.”
46. Goldbeter, A. Computational approaches to cellular rhythms. *Nature* **420**, 238-245 (2002).
47. Wolkenhauer, O., Sreenath, S. N., Wellstead, P., Ullah, M. & Cho, K. H. A systems- and signal-oriented approach to intracellular dynamics. *Biochem. Soc. Trans.* **33**, 507-515 (2005).
48. Suenaga, A. et al. Tyr-317 phosphorylation increases shc structural rigidity and reduces coupling of domain motions remote from the phosphorylation site as revealed by molecular dynamics simulations. *J. Biol. Chem.* **279**, 4657-4662 (2004).
49. Markevich, N. I. et al. Signal processing at the Ras circuit: What shapes Ras activation patterns? *IEE Systems Biology* **1**, 104-113 (2004).

50. Shvartsman, S. Y., Muratov, C. B. & Lauffenburger, D. A. Modeling and computational analysis of EGF receptor-mediated cell communication in *Drosophila* oogenesis. *Development* **129**, 2577-2589 (2002).
51. Resat, H., Ewald, J. A., Dixon, D. A. & Wiley, H. S. An integrated model of epidermal growth factor receptor trafficking and signal transduction. *Biophys. J.* **85**, 730-743 (2003).
52. Morton-Firth, C. J. & Bray, D. Predicting temporal fluctuations in an intracellular signalling pathway. *J. Theor. Biol.* **192**, 117-128 (1998).
53. Kratchmarova, I., Blagoev, B., Haack-Sorensen, M., Kassem, M. & Mann, M. Mechanism of divergent growth factor effects in mesenchymal stem cell differentiation. *Science* **308**, 1472-1477 (2005).
54. Kim, J. E., Tannenbaum, S. R. & White, F. M. Global phosphoproteome of HT-29 human colon adenocarcinoma cells. *J. Proteome Res.* **4**, 1339-1346 (2005).
55. Borisov, N. M., Markevich, N. I., Hoek, J. B. & Kholodenko, B. N. Signaling through receptors and scaffolds: Independent interactions reduce combinatorial complexity. *Biophys. J.* **89**, 951-966 (2005).
56. Blinov, M. L., Faeder, J. R., Goldstein, B. & Hlavacek, W. S. BioNetGen: software for rule-based modeling of signal transduction based on the interactions of molecular domains. *Bioinformatics* **20**, 3289-3291 (2004).
57. Faeder, J. R., Blinov, M. L., Goldstein, B. & Hlavacek, W. S. Rule-based modeling of biochemical networks. *Complexity* **10**, 22-41 (2005).
58. Lok, L. & Brent, R. Automatic generation of cellular reaction networks with Molecuizer 1.0. *Nat. Biotechnol.* **23**, 131-136 (2005).
59. Conzelmann, H., Saez-Rodriguez, J., Sauter, T., Kholodenko, B. N. & Gilles, E. D. A domain-oriented approach to the reduction of combinatorial complexity in signal transduction networks. *BMC Bioinformatics* 2006, 7:34 doi:10.1186/1471-2105-7-34.
60. Goldbeter, A. & Koshland, D. E., Jr. An amplified sensitivity arising from covalent modification in biological systems. *Proc. Natl. Acad. Sci. U S A* **78**, 6840-6844 (1981).
Almost 25 years after publication, still an instructive theoretical exploration of input-output responses of a universal signalling cycle; the term "ultrasensitivity" was coined.
61. Ortega, F., Acerenza, L., Westerhoff, H. V., Mas, F. & Cascante, M. Product dependence and bifunctionality compromise the ultrasensitivity of signal transduction cascades. *Proc. Natl. Acad. Sci. U S A* **99**, 1170-1175 (2002).
62. Ferrell, J. E., Jr. & Bhatt, R. R. Mechanistic studies of the dual phosphorylation of mitogen-activated protein kinase. *J. Biol. Chem.* **272**, 19008-19016 (1997).
63. Kholodenko, B. N., Hoek, J. B., Brown, G. C. & Westerhoff, H. V. in *BioThermoKinetics in the Post Genomic Era* (eds. C., L., I.-L., P. & L., G.) 102-107 (Güterborg, 1998).
64. Salazar, C. & Hofer, T. Allosteric regulation of the transcription factor NFAT1 by multiple phosphorylation sites: a mathematical analysis. *J. Mol. Biol.* **327**, 31-45 (2003).
65. Markevich, N. I., Hoek, J. B. & Kholodenko, B. N. Signaling switches and bistability arising from multisite phosphorylation in protein kinase cascades. *J. Cell. Biol.* **164**, 353-359 (2004).
66. Kholodenko, B. N., Hoek, J. B., Westerhoff, H. V. & Brown, G. C. Quantification of information transfer via cellular signal transduction pathways. *FEBS Lett.* **414**, 430-434 (1997).
67. Ferrell, J. E., Jr. How responses get more switch-like as you move down a protein kinase cascade. *Trends Biochem. Sci.* **22**, 288-289 (1997).
68. Sha, W. et al. Hysteresis drives cell-cycle transitions in *Xenopus laevis* egg extracts. *Proc. Natl. Acad. Sci. U S A* **100**, 975-980 (2003).
Together with reference 69, provided the first experimental demonstration that hysteresis and bistability in Cdc2 activation drives the cell-cycle oscillator.

69. Pomerening, J. R., Sontag, E. D. & Ferrell, J. E. Building a cell cycle oscillator: hysteresis and bistability in the activation of Cdc2. *Nat. Cell. Biol.* **5**, 346-351 (2003).
70. Stelling, J., Sauer, U., Szallasi, Z., Doyle, F. J., 3rd & Doyle, J. Robustness of cellular functions. *Cell* **118**, 675-685 (2004).
71. Vilar, J. M., Kueh, H. Y., Barkai, N. & Leibler, S. Mechanisms of noise-resistance in genetic oscillators. *Proc. Natl. Acad. Sci. U S A* **99**, 5988-5992 (2002).
72. Ferrell, J. E., Jr. Self-perpetuating states in signal transduction: positive feedback, double-negative feedback and bistability. *Curr. Opin. Cell Biol.* **14**, 140-148. (2002).
73. Levchenko, A., Bruck, J. & Sternberg, P. W. Scaffold proteins may biphasically affect the levels of mitogen-activated protein kinase signaling and reduce its threshold properties. *Proc. Natl. Acad. Sci. U S A* **97**, 5818-5823 (2000).
74. Kolch, W., Calder, M. & Gilbert, D. When kinases meet mathematics: the systems biology of MAPK signalling. *FEBS Lett.* **579**, 1891-1895 (2005).
Surveys the complexity of a multitude of feedback circuits and other regulations in the MAPK/ERK cascade and argues that their understanding requires mathematical modelling approaches.
75. Novak, B. & Tyson, J. J. Numerical analysis of a comprehensive model of M-phase control in *Xenopus* oocyte extracts and intact embryos. *J. Cell. Sci.* **106**, 1153-1168 (1993).
76. Bray, D. Signaling complexes: biophysical constraints on intracellular communication. *Annu. Rev. Biophys. Biomol. Struct.* **27**, 59-75 (1998).
A profound survey of the effects of the spatial organization and biophysical constraints imposed by macromolecular crowding, diffusion and membrane compartments on the control and evolution of cell communication.
77. Kholodenko, B. N., Hoek, J. B. & Westerhoff, H. V. Why cytoplasmic signalling proteins should be recruited to cell membranes. *Trends Cell Biol.* **10**, 173-178 (2000).
78. Haugh, J. M. & Lauffenburger, D. A. Physical modulation of intracellular signaling processes by locational regulation. *Biophys. J.* **72**, 2014-2031 (1997).
79. Simons, K. & Toomre, D. Lipid rafts and signal transduction. *Nat. Rev. Mol. Cell Biol.* **1**, 31-39 (2000).
80. Bauman, A. L. & Scott, J. D. Kinase- and phosphatase-anchoring proteins: harnessing the dynamic duo. *Nat. Cell Biol.* **4**, E203-206 (2002).
81. Sorkin, A. & Von Zastrow, M. Signal transduction and endocytosis: close encounters of many kinds. *Nat. Rev. Mol. Cell Biol.* **3**, 600-614 (2002).
82. Bivona, T. G. et al. Phospholipase Cgamma activates Ras on the Golgi apparatus by means of RasGRP1. *Nature* **424**, 694-698 (2003).
83. Kholodenko, B. N., Brown, G. C. & Hoek, J. B. Diffusion control of protein phosphorylation in signal transduction pathways. *Biochem. J.* **350**, 901-907 (2000).
84. Andersen, S. S. et al. Mitotic chromatin regulates phosphorylation of Stathmin/Op18. *Nature* **389**, 640-643 (1997).
85. Carazo-Salas, R. E. et al. Generation of GTP-bound Ran by RCC1 is required for chromatin-induced mitotic spindle formation. *Nature* **400**, 178-181 (1999).
86. Kalab, P., Weis, K. & Heald, R. Visualization of a Ran-GTP gradient in interphase and mitotic *Xenopus* egg extracts. *Science* **295**, 2452-2456 (2002).
Together with references 87 and 91, reported the first experimental observations of gradients of signalling proteins that control and maintain the assembly of the mitotic spindle around chromosomes.
87. Niethammer, P., Bastiaens, P. & Karsenti, E. Stathmin-tubulin interaction gradients in motile and mitotic cells. *Science* **303**, 1862-1866 (2004).

First experimental demonstration of spatial gradients of a phospho-protein (the microtubule-binding protein stathmin) within living cells.

88. Sprague, B. L. et al. Mechanisms of microtubule-based kinetochore positioning in the yeast metaphase spindle. *Biophys. J.* **84**, 3529-3246 (2003).
89. Gorlich, D., Seewald, M. J. & Ribbeck, K. Characterization of Ran-driven cargo transport and the RanGTPase system by kinetic measurements and computer simulation. *Embo J.* **22**, 1088-1100 (2003).
90. Wollman, R. et al. Efficient chromosome capture requires a bias in the 'search-and-capture' process during mitotic-spindle assembly. *Curr. Biol.* **15**, 828-832 (2005).
91. Caudron, M., Bunt, G., Bastiaens, P. & Karsenti, E. Spatial coordination of spindle assembly by chromosome-mediated signaling gradients. *Science* **309**, 1373-1376 (2005).
92. Rao, C. V., Kirby, J. R. & Arkin, A. P. Phosphatase localization in bacterial chemotaxis: divergent mechanism, convergent principles. *Phys. Biol.* **2**, 148-158 (2005).
93. Fell, D. A. Theoretical analyses of the functioning of the high- and low-K_m cyclic nucleotide phosphodiesterases in the regulation of the concentration of adenosine 3',5'-cyclic monophosphate in animal cells. *J. Theor. Biol.* **84**, 361-385 (1980).
94. Li, H. Y. & Zheng, Y. Phosphorylation of RCC1 in mitosis is essential for producing a high RanGTP concentration on chromosomes and for spindle assembly in mammalian cells. *Genes Dev.* **18**, 512-527 (2004).
95. Miaczynska, M., Pelkmans, L. & Zerial, M. Not just a sink: endosomes in control of signal transduction. *Curr. Opin. Cell Biol.* **16**, 400-406 (2004).
- Provides insight into how different types of endosomes, a variety of membrane microdomains and their protein complement control signal specificity in time and space.**
96. Howe, C. L. & Mobley, W. C. Signaling endosome hypothesis: A cellular mechanism for long distance communication. *J. Neurobiol.* **58**, 207-216 (2004).
97. Sorkin, A. Cargo recognition during clathrin-mediated endocytosis: a team effort. *Curr. Opin. Cell Biol.* **16**, 392-399 (2004).
98. Ginty, D. D. & Segal, R. A. Retrograde neurotrophin signaling: Trk-ing along the axon. *Curr. Opin. Neurobiol.* **12**, 268-274. (2002).
99. Campenot, R. B. & MacInnis, B. L. Retrograde transport of neurotrophins: fact and function. *J. Neurobiol.* **58**, 217-229 (2004).
100. MacInnis, B. L., Senger, D. L. & Campenot, R. B. Spatial requirements for TrkA kinase activity in the support of neuronal survival and axon growth in rat sympathetic neurons. *Neuropharmacology* **45**, 995-1010 (2003).
101. Hill, D. B., Plaza, M. J., Bonin, K. & Holzwarth, G. Fast vesicle transport in PC12 neurites: velocities and forces. *Eur. Biophys. J.* **33**, 623-32 (2004).
102. Grimm, V. et al. Pattern-oriented modeling of agent-based complex systems: lessons from ecology. *Science* **310**, 987-991 (2005).
103. Tyson, J. J., Chen, K. & Novak, B. Network dynamics and cell physiology. *Nat. Rev. Mol. Cell Biol.* **2**, 908-916. (2001).
104. Moehren, G. et al. Temperature dependence of the epidermal growth factor receptor signaling network can be accounted for by a kinetic model. *Biochemistry* **41**, 306-320 (2002).
105. Di Guglielmo, G. M., Baass, P. C., Ou, W. J., Posner, B. I. & Bergeron, J. J. Compartmentalization of SHC, GRB2 and mSOS, and hyperphosphorylation of Raf-1 by EGF but not insulin in liver parenchyma. *Embo J.* **13**, 4269-4277. (1994).
106. Machide, M., Kamitori, K. & Kohsaka, S. Hepatocyte growth factor-induced differential activation of phospholipase cgamma 1 and phosphatidylinositol 3-kinase is regulated by tyrosine phosphatase SHP-1 in astrocytes. *J. Biol. Chem.* **275**, 31392-31398. (2000).

107. Goryanin, I., Hodgman, T. C. & Selkov, E. Mathematical simulation and analysis of cellular metabolism and regulation. *Bioinformatics* **15**, 749-758 (1999).
108. Sauro, H. M. et al. Next generation simulation tools: the Systems Biology Workbench and BioSPICE integration. *Omics* **7**, 355-372 (2003).
109. Slepchenko, B. M., Schaff, J. C., Macara, I. & Loew, L. M. Quantitative cell biology with the Virtual Cell. *Trends Cell. Biol.* **13**, 570-576 (2003).
110. Sivakumaran, S., Hariharaputran, S., Mishra, J. & Bhalla, U. S. The Database of Quantitative Cellular Signaling: management and analysis of chemical kinetic models of signaling networks. *Bioinformatics* **19**, 408-415 (2003).
111. Campagne, F. et al. Quantitative information management for the biochemical computation of cellular networks. *Sci. STKE* **2004**, p111 (2004).
112. Olivier, B. G. & Snoep, J. L. Web-based kinetic modelling using JWS Online. *Bioinformatics* **20**, 2143-2144 (2004).

Acknowledgement.

I am grateful to Jan Hoek, Marc Birtwistle, Anatoly Kiyatkin, Nick Markevich, John Pastorino and Michael Tsyganov for stimulating discussions and help with illustrative materials. I apologize that it was not possible to cite all of the many valuable contributions to the field. This work is supported by the NIH grant GM59570.

Online links




The International Consortium "Systems Biology of RTK signaling" <http://www.rtkconsort.org/>
 The Systems Biology Markup Language (SBML) page <http://sbml.org/index.psp> lists over 90 software systems and databases of biological models, including the following used by the author:
 BioModels Database: <http://www.ebi.ac.uk/biomodels>
 BioNetGen: <http://cellsignaling.lanl.gov/bionetgen/>
 DOQCS <http://doqcs.ncbs.res.in/>
 DBsolve: http://biosim.genebee.msu.edu/dbsdownload_en.php
 Gepasi: <http://www.gepasi.org/>
 Systems Biology Workbench: www.sys-bio.org
 Silicon Cell: <http://jjj.biochem.sun.ac.za/>
 SigPath: <http://www.sigpath.org/>
 Virtual Cell: <http://www.nrcam.uchc.edu/>
 XPPAUT: <http://www.math.pitt.edu/~bard/xpp/xpp.html>

Figure legends

FIG. 1. Universal motifs of cellular signalling networks. **A.** One-site phosphorylation cycle. The protein M is phosphorylated by a kinase to yield the phosphorylated form M_p , which is dephosphorylated by an opposing phosphatase. **B.** A cycle of a small GTPase (Ran). A guanine nucleotide exchange factor (GEF) catalyzes the transformation of an inactive guanosine diphosphate (GDP)-bound form (Ran-GDP) into an active guanosine triphosphate (GTP)-bound form (Ran-GTP). A GTPase-activating protein (GAP) is the opposing enzyme that catalyzes the reverse transformation. **C.** A cascade of cycles. Negative feedback provides robustness to noise, increasing resistance to disturbances inside the feedback loop, but brings about oscillations if it is too strong and the cascade is ultrasensitive^{12,20}. Positive feedback greatly increases the sensitivity of the target to the signal and may also lead to bistability and relaxation oscillations^{12,18,46,72}.

FIG. 2. Feedback designs that can turn a universal signalling cycle into a bistable switch and relaxation oscillator. Simple cycle can turn bistable in four distinct ways; either M_p or M stimulates its own production (positive feedback) by product activation or substrate inhibition of the kinase or phosphatase reactions. Each of the four rows of feedback designs correspond to a different bistable switch, provided that the kinase and phosphatase abundances are assumed constant and only single feedback (within the M cycle) is present. Sixteen relaxation oscillation designs are generated by extra negative feedback brought about by negative or positive regulation of the synthesis or degradation rates of the kinase protein or phosphatase protein by M_p or M . Designs **A*-H*** are mirror images of designs **A-H**. Although synthesis and degradation reactions are shown for both the kinase and phosphatase proteins, the protein concentration that is not controlled by feedback from the M cycle is considered constant, resulting in only two differential equations for each diagram. All feedback regulations are described by simple Michaelis-Menten type expressions (BOX 2 and Supplementary Table S3). The remaining sixteen relaxation oscillation designs are shown in Supplementary FIG.S1 and can require some degree of cooperativity within feedback loops.

FIG. 3. Spatial segregation of two opposing enzymes in a protein-modification cycle generates intracellular gradients. Kinases localize to (A) supra-molecular structures (sphere) or (B) the cell membrane, whereas phosphatases are homogeneously distributed in the cytoplasm. The concentration gradients are shown by colour intensity. **C.** Stationary phosphorylation levels c_p decline with the distance d from the cell membrane toward the centre [Brown, 1999 #61 (see panel B)]. The steepness of the gradient (reciprocal of the characteristic length) is determined by the parameter α ($\alpha^2 = k_p/D$ is the ratio of the phosphatase activity k_p and the protein diffusivity D , BOX 3).

A**B****C****Fig. 1**




Fig 2.

A




Fig 3A

B




Fig. 3B

C




Fig. 3C






Figure Box 1. Parts A, B and C




Figure Box 2. Parts A, B and C

A




Figure Box 4, Part A

B




Figure Box 4, Part B