

10-26-2020

## Nurse Education and Enteral Feedings in Mechanically Ventilated Patients

Georgina Campbell

*Thomas Jefferson University, [georgina.campbell@jefferson.edu](mailto:georgina.campbell@jefferson.edu)*

John Durison

*Thomas Jefferson University, [john.durison@jefferson.edu](mailto:john.durison@jefferson.edu)*

Daniel Russo

*Thomas Jefferson University, [daniel.russo@jefferson.edu](mailto:daniel.russo@jefferson.edu)*

Follow this and additional works at: <https://jdc.jefferson.edu/nurseresidencyposters>



Part of the [Medical Sciences Commons](#), and the [Nursing Commons](#)

**[Let us know how access to this document benefits you](#)**

---

### Recommended Citation

Campbell, Georgina; Durison, John; and Russo, Daniel, "Nurse Education and Enteral Feedings in Mechanically Ventilated Patients" (2020). *Abington Jefferson Health Nurse Residency Posters*. 2. <https://jdc.jefferson.edu/nurseresidencyposters/2>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Abington Jefferson Health Nurse Residency Posters by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

## Introduction

### Purpose

- To measure the effect of educating nurses in the ICU on the latest research on early initiation of enteral feedings in mechanically ventilated patients.

### Problem Statement

- Research has shown that early initiation of enteral feeding in mechanically ventilated patients is associated with shorter lengths of stay in the ICU. (Schub, 2018). Some members of ICU care teams are not aware of this association.

### PICO question

- In mechanically ventilated ICU patients, is exposure of the nursing staff to the most recent research on early initiation of enteral feeding beneficial to its implementation compared to patients whose care team had not been exposed to that research?

## Evidence

### Literature Review

- Artinian, DiGiovine, and Krayem found that starting enteral feeding within 48 hours of being placed on mechanical ventilator reduced mortality rates, infection rates, but also increased rates of VAP.

- Hsu, Huang, and Kang found that earlier feeding was associated with increased length of stay in the more ill patients, and also an increase in GI complications. Severity of illness, however, was linked for to mortality than feeding.

- Charney, Compber, Hise, Kattelmann, Russell, and Stokes found that early enteral feeding (24-48h) was associated with reduced incidence of grade 1 infectious complications and may be associated with decreased length of stay. However, no statistically significant correlation between feeding timing and mortality was noted.

- Yip, Rai, & Wong found that an escalating energy deficit was more tied to mortality rather than early enteral feeding. Additionally, delays in feeding were hard to overcome, especially in critically ill patients.

- Schub & Karakashian found that Initiation of nutritional support in ICU patients within 48 hours of admission to the ICU is associated with improved clinical outcomes, lower infection rates, and a reduced length of hospital stay. Additionally, early enteral feeding at a slow rate is more beneficial to ICU patients than other feeding approaches.

## Methods

- Nursing staff in the Medical ICU and Trauma/Neuro ICUs were provided with a review of recent literature on the benefits of early enteral feeding in mechanically ventilated patients.

- A screening tool was distributed along with the literature review to gather data on timing of feed initiation, enteral access, consults and length of stay.

- Timing of enteral feed initiation after distribution of this literature was compared with timing of enteral feed initiation in a randomly chosen sample of previous MICU/STU/NCC patients in the three months before distribution of this literature. Sample sizes were similar (50 patients vs 43 patients). This was done through chart review.

## Results

### Control Group

- Of the patients chosen randomly in the months before distribution of literature review & screening tool (n=50), 56% (28 pts) had enteral feeds initiated within 48 hours of intubation, with 32% (16 pts) starting later than 48 hours and 12% (6 pts) with no enteral feeds started.

- 80% of this population (40 pts) had enteral access established simultaneously with or on the same day as intubation, with 16% (8 pts) having access established 1 day or more post intubation and 4% (2 pts) having no enteral access established.

- A nutrition consult was placed for 86% of these patients.

-Among the 28 patients in the early enteral feeding group and not taking into consideration patients that expired in the ICU, (8), ICU length of stay was on average 9.4 days.

- Among the 16 patients in the late enteral feeding group and not taking into consideration patients that expired in the ICU (5), ICU length of stay was on average 13.2 days.

-Among the 6 patients that did not receive enteral feeding in the ICU and not taking into consideration patients that expired in the ICU, (2), length of stay was on average 5 days.

### Experimental Group

- Of the patients considered after distribution of literature review & screening tool (n=43), 51% (22 pts) had enteral feeds initiated within 48 hours of intubation, with 37% (16 pts) starting later than 48 hours and 12% (5 pts) with no enteral feeds started.

- 86% of this population (37 pts) had enteral access established simultaneously with or on the same day as intubation, with 12% having access established 1 day or more post intubation and 2% (1 pt) having no enteral access established.

- A nutrition consult was placed for 88% of these patients.

-Among the 22 patients in the early enteral feeding group and not taking into consideration patients that expired in the ICU, (6), ICU length of stay was on average 11.8 days.

- Among the 16 patients in the late enteral feeding group and not taking into consideration patients that expired in the ICU, (4), ICU length of stay was on average 9.5 days.

- Among the 5 patients that did not receive enteral feeding in the ICU and not taking into consideration patients that expired in the ICU, (3), length of stay was on average 3.5 days.

## Next Steps

We recommend that the clinical staff and in particular the policymakers in these units stay informed on the latest research on the benefits of early initiation of enteral feeding in mechanically ventilated adults in the ICU.

## Conclusions

- Based solely on the data collected, it would appear that exposure to the review of literature had a net negative effect on the early initiation of enteral feedings. Average length of stay in the ICU flipped from longer in the late-fed control group to shorter in the early-fed experimental group.

- That being said, there are a substantial number of confounding factors in the data (see previous two slides). Additionally, the sample size(s) for these groups were small (n=50 and n=43).

- We recommend that data continue to be collected and analyzed on timing of enteral feed initiation in mechanically ventilated adults in the ICU.

## References

1. Artinian V, Krayem H, DiGiovine B. Effects of early enteral feeding on the outcome of critically ill mechanically ventilated medical patients. *Chest* . 2006;129(4):960-967.doi:10.1378/chest.129.4.960

2. Huang, H., Hsu, C., Kang, S. et al. Association between illness severity and timing of initial enteral feeding in critically ill patients: a retrospective observational study. *Nutr J* 11, 30 (2012).

3. Kattelmann KK, Hise M, Russell M, Charney P, Stokes M, Compber C. Preliminary evidence for a medical nutrition therapy protocol: enteral feedings for critically ill patients. *J Am Diet Assoc* . 2006;106(8):1226-1241.doi:10.1016/j.jada.2006.05.320

4. Yip, K.F., Rai, V. & Wong, K.K. Evaluation of delivery of enteral nutrition in mechanically ventilated Malaysian ICU patients. *BMC Anesthesiol* 14, 127 (2014).

5. Schub, T. B., & Karakashian, A. R. B. (2018). Feeding Practices: Critically Ill Patients. *CINAHL Nursing Guide*.