2-2021

Timing of Preoperative Surgical Antibiotic Prophylaxis Prior to One-to-Three Level Elective Lumbar Fusion

Akash Singh

Thomas Jefferson University, akash.singh@students.jefferson.edu

Gregory Schroeder, MD

Thomas Jefferson University, Gregory.Schroeder@jefferson.edu

Jose Canseco, MD

Thomas Jefferson University, jose.canseco@jefferson.edu

Alexander Vaccaro, MD

Thomas Jefferson University, Alex.Vaccaro@rothmaninstitute.com

Parthik Patel, MD

Thomas Jefferson University, parthik.patel@jefferson.edu

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2023_phase1

Part of the [Orthopedics Commons](#), and the [Translational Medical Research Commons](#)

See next page for additional authors

[Let us know how access to this document benefits you](#)

Recommended Citation

Singh, Akash; Schroeder, MD, Gregory; Canseco, MD, Jose; Vaccaro, MD, Alexander; Patel, MD, Parthik; and Reyes, MD, Ariana, "Timing of Preoperative Surgical Antibiotic Prophylaxis Prior to One-to-Three Level Elective Lumbar Fusion" (2021). *Phase 1*. Paper 108.

https://jdc.jefferson.edu/si_ctr_2023_phase1/108

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Akash Singh; Gregory Schroeder, MD; Jose Canseco, MD; Alexander Vaccaro, MD; Parthik Patel, MD; and Ariana Reyes, MD

Timing of Preoperative Surgical Antibiotics Prior to One-to-Three Level Lumbar Fusions

Singh, Akash; Schroeder, Gregory, MD*; Canseco, Jose, MD; Vaccaro, Alexander, MD; Patel, Parthik, MD; Reyes, Ariana, MD

Disclosures and Acknowledgements

- No conflicts of interests
- Acknowledgements
 - Dr. Gregory Schroeder – Project Advisor
 - Dept. of Orthopedic Surgery at Rothman Orthopedic Institute

Introduction & Objectives

- Reducing Surgical Site Infections (SSIs) in Lumbar Fusion patients
- Preoperative
 - Medical clearances for significant comorbidities¹
 - Cardiac stress, HbA1c, albumin
 - Nasal swabs for MRSA
 - Stopping anticoagulation
- Intraoperative
 - Thoroughly supervising the OR
- Antibiotic Prophylaxis before lumbar fusions
 - Cefazolin² – Pre vs. Post Op
 - Vancomycin/Gentamycin³ for MRSA
 - Greater than 120 min before incision and after incision → higher SSIs⁴
- Cefuroxime given 59-30 mins before → lower SSIs⁵

Objectives & Hypothesis

- Research Question
 - What is the optimal timing of antibiotic prophylaxis in One-to-Three Level Lumbar Fusion Patients?
- Hypothesis
 - SSIs are greater in patients who receive antibiotic prophylaxis greater than 60 minutes prior to surgical incision compared with patients who receive antibiotic prophylaxis within 60 minutes.

Methods

- Retrospective Cohort
- Patients status post elective, 1-to-3 level lumbar fusion at Rothman Orthopedics between January 2013- December 2018
- Primary analysis: Timing of antibiotic before incision
 - Group A: 0-15 min Group C: 31-45 min Group E: 61+
 - Group B: 16-30 min Group D: 46-60 min
- Secondary analysis: Type of antibiotic
 - Cefazolin
 - Vancomycin
- Primary Outcome: SSI
- Secondary Outcomes
 - 90 day readmissions
 - Revision surgeries

Results

- 27 SSI out of 1,131 patients (2.4%)
- Group E compared to Group A
 - 11x more likely to get SSI
 - ($p < 0.001$)
 - 7.2x more likely to get readmitted within 90 days
- Groups B-D: no significantly higher odds of SSI vs. Group A

Group Letter	Timing of Antibiotic Admin.	SSIs (Rates)
Group A N=571	0-15 min	11 (1.9%)
Group B N=294	16-30 min	5 (1.7%)
Group C N=124	31-45 min	2 (1.6%)
Group D N=69	46-60 min	1 (1.4%)
Group E N=73	61+ min	8 (11.0%)

- By antibiotic type
 - Group E still highest rates of SSI ($p < 0.001$)

Vancomycin

Group Letter	Timing of Antibiotic	SSIs (Rates)
Group A N=23	0-15 min	0 (0%)
Group B N=17	16-30 min	0 (0%)
Group C N=30	31-45 min	1 (3.3%)
Group D N=58	46-60 min	0 (0%)
Group E N=71	61+ min	6 (8.5%)

Cefazolin

Group Letter	Timing of Antibiotic	SSIs (Rates)
Group A N=548	0-15 min	11 (2.0%)
Group B N=277	16-30 min	5 (1.8%)
Group C N=94	31-45 min	1 (1.1%)
Group D N=11	46-60 min	1 (9.0%)
Group E N=2	61+ min	2 (100%)

Results

Probability of SSI by Timing

Stratified by Antibiotic

Conclusions

- Preoperative antibiotic administration beyond an hour → 11x increased likelihood of SSI
 - One-to-Three Level Lumbar Fusion patients
- Other surgeries:
 - Cervical & Lumbar⁶: 3x increased likelihood if given >1 hour
 - Hip & Knee, Colorectal, Vascular, Gyn⁷: Higher rates of SSI if given > 1 hour
 - Cefuroxime use⁸: 30-59 mins has lower SSI than 0-30 mins
- Further research needed to better identify optimal timing between 0-60 minutes before incision

1. Epstein N. Preoperative measures to prevent/minimize risk of surgical site infection in spinal surgery. *Surg Neurol Int.* 2018; 9:251.
2. Marimuthu C, Abraham VT, Ravichandran M, Achimuthu R. Antimicrobial prophylaxis in instrumented spinal fusion surgery: A comparative analysis of 24-Hour and 72-Hour dosages. *Asian Spine J.* 2016;10:1018–22
3. Park HY, Sheppard W, Smith R, Xiao J, Gatto J, Bowen R, et al. The combined administration of vancomycin IV, standard prophylactic antibiotics, and vancomycin powder in spinal instrumentation surgery: Does the routine use affect infection rates and bacterial resistance? *J Spine Surg.* 2018;4:173–9.
4. Willem de Jonge S, Gans S, Ateema J, Solomkin J, Dellinger P, Boermeester M. Timing of preoperative antibiotic prophylaxis in 54,552 patients and the risk of surgical site infection - A systematic review and meta-analysis. *Medicine (Baltimore).* 2017; 96 (29): e6903.
5. Weber WP, Marti WR, Zwahlen M, Misteli H, Rosenthal R, Reck S, Fueglistaler P, Bolli M, Trampuz A, Oertli D, Widmer AF. The timing of surgical antimicrobial prophylaxis. *Ann Surg.* 2008; 247(6): 918-26.
6. Bratzler DW, Dellinger EP, Olsen KM, Perl TM, Auwaerter PG, Bolon MK, Fish DN, Napolitano LM, Sawyer RG, Slain D, Steinberg JP, Weinstein RA; American Society of Health-System Pharmacists; Infectious Disease Society of America; Surgical Infection Society; Society for Healthcare Epidemiology of America. Clinical practice guidelines for antimicrobial prophylaxis in surgery. *Am J Health Syst Pharm.* 2013 Feb 1;70(3):195-283. doi: 10.2146/ajhp120568. PMID: 23327981.
7. Hawn, M. T. et al. Timing of surgical antibiotic prophylaxis and the risk of surgical site infection. *JAMA Surg.* 148, 649–657 (2013).
8. Weber, W. P. et al. The timing of surgical antimicrobial prophylaxis. *Ann. Surg.* 247, 918–926 (2008).