

8-1-2010

## Phospholemman: a novel cardiac stress protein.

Joseph Y Cheung  
*Thomas Jefferson University*

Xue-Qian Zhang  
*Thomas Jefferson University*


Jianliang Song  
*Thomas Jefferson University*

Erhe Gao  
*Thomas Jefferson University*

Joseph E Rabinowitz  
*Thomas Jefferson University*

Follow this and additional works at: <https://jdc.jefferson.edu/transmedfp>

See next page for additional authors

 Part of the [Cardiology Commons](#)

[Let us know how access to this document benefits you](#)

---

### Recommended Citation

Cheung, Joseph Y; Zhang, Xue-Qian; Song, Jianliang; Gao, Erhe; Rabinowitz, Joseph E; Chan, Tung O; and Wang, Jufang, "Phospholemman: a novel cardiac stress protein." (2010). *Center for Translational Medicine Faculty Papers*. Paper 4.

<https://jdc.jefferson.edu/transmedfp/4>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Center for Translational Medicine Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

---

**Authors**

Joseph Y Cheung, Xue-Qian Zhang, Jianliang Song, Erhe Gao, Joseph E Rabinowitz, Tung O Chan, and Jufang Wang

**As submitted to:**

***Clinical and translational science***

**And later published as:**

**“Phospholemman: A Novel Cardiac Stress Protein”**

**Volume 3, Number 4, August 2010 , pp. 189-196(8)**

**DOI: 10.1111/j.1752-8062.2010.00213.x**

Joseph Y. Cheung, M.D., Ph.D., Xue-Qian Zhang, M.D., Jianliang Song, M.D., Ph.D., Erhe Gao, M.D., Ph.D., Joseph E. Rabinowitz, Ph.D., Tung O. Chan, Ph.D., and JuFang Wang, M.D.

Division of Nephrology and Center of Translational Medicine, Department of Medicine,  
Jefferson Medical College, Thomas Jefferson University, Philadelphia, PA 19107.

Address correspondence to:

Joseph Y. Cheung, M.D., Ph.D.

Division of Nephrology

Jefferson Medical College

833 Chestnut Street, Suite 700

Philadelphia, PA 19107.

Tel. 215-503-6830

Tel. 215-503-4099

Email. [joseph.cheung@jefferson.edu](mailto:joseph.cheung@jefferson.edu)

## **Abstract**

Phospholemman (PLM), a member of the FXYD family of regulators of ion transport, is a major sarcolemmal substrate for protein kinases A and C in cardiac and skeletal muscle. In the heart, PLM co-localizes and co-immunoprecipitates with Na<sup>+</sup>-K<sup>+</sup>-ATPase, Na<sup>+</sup>/Ca<sup>2+</sup> exchanger and L-type Ca<sup>2+</sup> channel. Functionally, when phosphorylated at serine<sup>68</sup>, PLM stimulates Na<sup>+</sup>-K<sup>+</sup>-ATPase but inhibits Na<sup>+</sup>/Ca<sup>2+</sup> exchanger in cardiac myocytes. In heterologous expression systems, PLM modulates the gating of cardiac L-type Ca<sup>2+</sup> channel. Therefore, PLM occupies a key modulatory role in intracellular Na<sup>+</sup> and Ca<sup>2+</sup> homeostasis and is intimately involved in regulation of excitation-contraction (EC) coupling. Genetic ablation of PLM results in a slight increase in baseline cardiac contractility and prolongation of action potential duration. When hearts are subjected to catecholamine stress, PLM minimizes the risks of arrhythmogenesis by reducing Na<sup>+</sup> overload and simultaneously preserves inotropy by inhibiting Na<sup>+</sup>/Ca<sup>2+</sup> exchanger. In heart failure, both expression and phosphorylation state of PLM are altered and may partly account for abnormalities in EC coupling. The unique role of PLM in regulation of Na<sup>+</sup>-K<sup>+</sup>-ATPase, Na<sup>+</sup>/Ca<sup>2+</sup> exchanger and potentially L-type Ca<sup>2+</sup> channel in the heart, together with the

changes in its expression and phosphorylation in heart failure, make PLM a rational and novel target for development of drugs in our armamentarium against heart failure.

## **Introduction**

Phospholemman (PLM) was initially identified by Larry Jones in 1985 (58) as a 15-kDa sarcolemmal (SL) protein that is phosphorylated in response to isoproterenol and is distinct from phospholamban (PLB)(58). Follow-up studies indicated that this 15-kDa SL protein is also phosphorylated by protein kinase (PK) C (59) and  $\alpha$ -adrenergic agonists (38). In 1991, this 15-kDa SL phosphoprotein was purified, the complete protein sequence determined by Edman degradation, the cDNA cloned, and the name “phospholemman” was coined (53). In 1997, the human PLM gene is localized to chromosome 19q13.1 (11).

PLM is synthesized as a 92 amino acid peptide containing at its N-terminus a 20 amino acid signal peptide which is cleaved off during processing. The mature protein contains 72 amino acid residues with a calculated molecular weight of 8409, but a mobility of ~15 kDa in SDS-PAGE gels. The first 17 amino acid residues lie in the extracellular domain. The transmembrane (TM) region contains 20 amino acids (residues 18-37) while the remaining 35 amino acids (residues 38-72) at the C-terminus are in the cytoplasm. Palmer et al. (53) also noted sequence homology between PLM and  $\gamma$ -subunit of  $\text{Na}^+$ - $\text{K}^+$ -ATPase, as well as a short region of sequence

similarity between PLM (at its C-terminus facing the cytoplasm) and PLB (at its N-terminus also facing the cytoplasm). This region of sequence similarity (RSSIRRLST<sup>69</sup> in PLM and RSAIRRAST<sup>17</sup> in PLB) contains serines and threonines that are potential phosphorylation sites. Indeed, serine<sup>68</sup> in PLM and serine<sup>16</sup> in PLB are phosphorylated by PKA (67, 79).

The extracellular N-terminus of PLM contains a FXYD motif, and the cytoplasmic tail of dog, human and rat PLM contains 3 serines (at residues 62, 63 and 68) and 1 threonine (at residue 69) but threonine<sup>69</sup> is replaced by serine in mouse PLM. By nuclear magnetic resonance (NMR) (23) and infrared spectroscopy (2), the TM domain of PLM reconstituted in liposomes is an  $\alpha$ -helix with a maximum tilt of 15-17<sup>0</sup>. Specifically, NMR spectroscopic studies of highly purified PLM in model micelles indicate that the molecule consists of 4  $\alpha$ -helices: H1 (residues 12-17) is in the extracellular N-terminus, H2 (residues 22-38) is the main TM helix followed by the short H3 (residues 39-45), and H4 (residues 60-68) in the C-terminus is connected to H3 by a flexible linker (Fig. 1)(76). In vivo, PKA phosphorylates serine<sup>68</sup> while PKC phosphorylates serine<sup>63</sup> and serine<sup>68</sup> of PLM (79). In vitro studies using PLM fragments suggest that PKA also phosphorylates serine<sup>63</sup> while PKC phosphorylates threonine<sup>69</sup> (25). In adult rat myocytes, ~46% of serine<sup>68</sup> and ~16% of serine<sup>63</sup> are estimated to be phosphorylated in the resting state (69). Using phospho-specific anti-PLM antibodies (25, 62), ~30-40% of PLM in adult rat myocytes (25, 89) and ~25% of PLM in guinea pig myocytes (66) are phosphorylated under basal conditions. In transfected HEK293 cells, ~30-45 % of exogenous PLM is phosphorylated under resting conditions (92).

Based on observations on *Xenopus* oocytes in which PLM is overexpressed, Randall Moorman suggested that PLM is a hyperpolarization-activated anion-selective channel (46). When reconstituted in lipid bilayers, PLM forms a channel that is highly selective for taurine

(12) and is thought to be involved in regulation of cell volume in non-cardiac tissues (15, 47).

The function of PLM in the heart remains unknown until the dawning of the 21<sup>st</sup> century.

### **Phospholemman: Founding Member of the FXYD Family of Regulators of Ion Transport**

In 2000, Kathy Sweadner described the FXYD family of regulators of ion transport (73), of which PLM is the first cloned member (FXYD1). At present, there are at least 12 known FXYD proteins, including  $\gamma$ -subunit of  $\text{Na}^+\text{-K}^+\text{-ATPase}$  (FXYD2), mammary associated tumor 8 kDa (MAT-8 or FXYD3), channel inducing factor (CHIF or FXYD4), dysadherin (FXYD5; also known as related to ion channel RIC), phosphohippolin (FXYD6), FXYD7, and PLM-S (FXYD10; the shark homolog of PLM). As a family, FXYD proteins are found predominantly in tissues involved in solute and fluid transport (kidney, colon, pancreas, mammary gland, liver, lung, prostate and placenta) or are electrically excitable (heart, skeletal and neural tissues). All FXYD members have the signature FXYD motif in the N-terminus and a single TM domain. Except for  $\gamma$ -subunit of  $\text{Na}^+\text{-K}^+\text{-ATPase}$ , all other known members of the FXYD gene family has at least one serine or threonine within the cytoplasmic tail, indicating potential phosphorylation sites. PLM is unique among FXYD proteins in that it has consensus sequence for phosphorylation by PKA (RRXS), PKC (RXXSXR) and never-in-mitosis aspergillus (NIMA) kinase (FRXS/T). PLM is also a substrate for myotonic dystrophy protein kinase (48). The  $\gamma$ -subunit of  $\text{Na}^+\text{-K}^+\text{-ATPase}$  is the only member in the FXYD family boasting two alternative splice variants (FXYD2a and FXYD2b).

### **Phospholemman: Regulator of Cardiac $\text{Na}^+\text{-K}^+\text{-ATPase}$**

In 2002, Kaethi Geering demonstrated that PLM co-immunoprecipitates with  $\alpha$ -subunits of  $\text{Na}^+\text{-K}^+\text{-ATPase}$  in bovine sarcolemma (14). In isolated adult rat cardiac myocytes, PLM co-localizes with  $\alpha$ -subunits of  $\text{Na}^+\text{-K}^+\text{-ATPase}$  (Fig. 2). When co-expressed with  $\alpha$ - and  $\beta$ -subunits

of Na<sup>+</sup>-K<sup>+</sup>-ATPase in *Xenopus* oocytes, PLM modulates Na<sup>+</sup>-K<sup>+</sup>-ATPase activity primarily by decreasing K<sub>m</sub> for Na<sup>+</sup> and K<sup>+</sup> without affecting V<sub>max</sub> (14). Data obtained from cardiac myocytes or homogenates indicate that PLM inhibits Na<sup>+</sup>-K<sup>+</sup>-ATPase by reducing its apparent affinities for intracellular Na<sup>+</sup> (9, 17) and external K<sup>+</sup> (29) or decreasing V<sub>max</sub> (3, 24, 25, 28, 55, 66, 70, 80, 89). When phosphorylated at serine<sup>68</sup>, PLM relieves its inhibition on Na<sup>+</sup>-K<sup>+</sup>-ATPase by decreasing its apparent K<sub>m</sub> for Na<sup>+</sup> (17, 80) but not for K<sup>+</sup> (29) and increasing V<sub>max</sub> (24, 66, 80).

In terms of molecular interactions between PLM and Na<sup>+</sup>-K<sup>+</sup>-ATPase, mutational analysis suggests that FXYD proteins (FXYD2, 4 and 7) interact with TM9 segment of Na<sup>+</sup>-K<sup>+</sup>-ATPase (36). Co-immunoprecipitation and covalent cross-linking studies demonstrate the TM segment of PLM is close to TM2 segment of Na<sup>+</sup>-K<sup>+</sup>-ATPase (39). Molecular modeling based on Ca<sup>2+</sup>-ATPase crystal structure in the E<sub>1</sub>ATP-bound conformation suggests that the single TM segment of FXYD proteins docks into the groove between TM segments 2, 6 and 9 of the α-subunit of Na<sup>+</sup>-K<sup>+</sup>-ATPase (39). High resolution crystal structure (2.4 Angstrom) of shark rectal gland Na<sup>+</sup>-K<sup>+</sup>-ATPase in the E2.2K<sup>+</sup>.P<sub>i</sub> state indicates that FXYD proteins interact almost exclusively with the outside of TM9 of the α-subunit (65). The role of the signature FXYD(Y) motif is to stabilize interactions between α- and β-subunits of Na<sup>+</sup>-K<sup>+</sup>-ATPase and residue D (in the FXYD motif) caps the helix and defines the membrane border (65). In transfected HEK293 cells, PLM interacts with either α1- or α2-subunit of Na<sup>+</sup>-K<sup>+</sup>-ATPase in a 1:1 stoichiometry (9). Phosphorylation of PLM-S causes it to dissociate from shark Na<sup>+</sup>-K<sup>+</sup>-ATPase (42). However, phosphorylation of PLM did not cause it to dissociate from the α-subunit of Na<sup>+</sup>-K<sup>+</sup>-ATPase (9, 66). Despite NMR studies of PLM in micelles showing no major conformational changes on phosphorylation of serine<sup>68</sup> (77), in intact cells examined with fluorescence resonance energy



transfer (FRET), interaction between PLM and Na<sup>+</sup>-K<sup>+</sup>-ATPase is decreased on phosphorylation of PLM (9, 10).

There are 4 isoforms of the catalytic  $\alpha$ -subunits of Na<sup>+</sup>-K<sup>+</sup>-ATPase and expression of a particular  $\alpha$ -isoform is both tissue- and species-dependent (7). Human (43, 85) and rabbit (8) hearts are known to express  $\alpha$ 1-,  $\alpha$ 2- and  $\alpha$ -3 isoforms while rodent hearts express only  $\alpha$ 1- (ouabain-resistant) and  $\alpha$ 2-isoforms of Na<sup>+</sup>-K<sup>+</sup>-ATPase (4, 66, 72, 80). In both adult rat and mouse ventricles, the ouabain-sensitive  $\alpha$ 2-subunit is preferentially localized to the t-tubules (4, 74) and its activity represents <25% of total Na<sup>+</sup>-K<sup>+</sup>-ATPase activities (4, 72, 80). PLM co-immunoprecipitates all 3  $\alpha$ -subunits of Na<sup>+</sup>-K<sup>+</sup>-ATPase in human and rabbit (8),  $\alpha$ 1- and  $\alpha$ 2-subunits in mouse (80) and bovine (14), but only  $\alpha$ 1-subunit in rat (24) and guinea pig (66) hearts. In both wild-type (WT) mouse (80) and guinea pig (66) ventricular myocytes, PLM regulates the activity of  $\alpha$ 1- but not  $\alpha$ 2-isoform of Na<sup>+</sup>-K<sup>+</sup>-ATPase. This conclusion must be tempered with the recent finding that in “SWAP” mouse (21) in which the ouabain affinities of the  $\alpha$ -subunits are reversed, PLM regulates the apparent affinities for Na<sup>+</sup> of both  $\alpha$ 1- and  $\alpha$ 2-subunits of Na<sup>+</sup>-K<sup>+</sup>-ATPase (9). Together with observations made on *Xenopus* oocytes heterologously expressing PLM and Na<sup>+</sup>-K<sup>+</sup>-ATPase (6, 14), it is likely the PLM regulates both  $\alpha$ 1- and  $\alpha$ 2-subunits of Na<sup>+</sup>-K<sup>+</sup>-ATPase.

### **Cardiac Excitation-Contraction Coupling**

Both  $\alpha$ 1- (21) and  $\alpha$ 2-subunits (74) of Na<sup>+</sup>-K<sup>+</sup>-ATPase have been implicated in the control of cardiac contractility. Therefore, modulation of Na<sup>+</sup>-K<sup>+</sup>-ATPase activity suggests an important role for PLM in regulation of inotropy. We will give a brief overview of cardiac excitation-contraction coupling (Fig. 3) which has previously been reviewed by Don Bers in detail (5). During the upstroke of the action potential, Na<sup>+</sup> enters via Na<sup>+</sup> channels and further

depolarizes the sarcolemma. Depolarization activates the voltage-dependent L-type  $\text{Ca}^{2+}$  channels, allowing extracellular  $\text{Ca}^{2+}$  to enter as an inward current ( $I_{\text{Ca}}$ ), which contributes to the plateau phase of the action potential. Some  $\text{Ca}^{2+}$  also enters via the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (NCX1) operating in the reverse mode (3  $\text{Na}^+$  out: 1  $\text{Ca}^{2+}$  in) during this phase of the action potential, although the amount and duration of  $\text{Ca}^{2+}$  influx via reverse  $\text{Na}^+/\text{Ca}^{2+}$  exchange vary among species.  $\text{Ca}^{2+}$  entry triggers release of  $\sim 2/3$  of  $\text{Ca}^{2+}$  stored in the sarcoplasmic reticulum (SR) via the ryanodine receptor (RyR2). The combination of  $\text{Ca}^{2+}$  influx and SR  $\text{Ca}^{2+}$  release abruptly raises the free intracellular  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ), allowing  $\text{Ca}^{2+}$  to bind to troponin C and activate the contractile apparatus. Relaxation requires termination of SR  $\text{Ca}^{2+}$  release, and  $[\text{Ca}^{2+}]_i$  to decline so that  $\text{Ca}^{2+}$  can dissociate from troponin C. About 70-92% of myoplasmic  $\text{Ca}^{2+}$  is re-sequestered into the SR by sarco(endo)plasmic reticulum  $\text{Ca}^{2+}$ -ATPase (SERCA2) which is under the control of PLB. To maintain steady-state  $\text{Ca}^{2+}$  balance,  $\text{Ca}^{2+}$  that has entered during systole is mainly extruded by NCX1 operating in the forward mode (3  $\text{Na}^+$  in: 1  $\text{Ca}^{2+}$  out), with the sarcolemmal  $\text{Ca}^{2+}$ -ATPase playing a smaller role. Likewise, the small amount of  $\text{Na}^+$  that has entered during depolarization is extruded by  $\text{Na}^+/\text{K}^+$ -ATPase during diastole. In this way, the cardiac myocyte maintains beat-to-beat  $\text{Ca}^{2+}$  and  $\text{Na}^+$  balance. Outward  $\text{K}^+$  currents contribute to repolarization phase of the action potential.

Among the many transporters and ion channels involved in cardiac  $\text{Ca}^{2+}$  fluxes, NCX1 is unique in that during an action potential, it participates in  $\text{Ca}^{2+}$  influx,  $[\text{Ca}^{2+}]_i$  transient buffering and  $\text{Ca}^{2+}$  efflux (5). The direction of  $\text{Ca}^{2+}$  flux (in or out) depends on the thermodynamic driving force determined by the membrane potential ( $E_m$ ) and the concentrations of  $\text{Na}^+$  and  $\text{Ca}^{2+}$  ions sensed by NCX1. Other than the determinants of its thermodynamic driving force, remarkably little is known about the regulation of NCX1.

In the heart, inhibition of Na<sup>+</sup>-K<sup>+</sup>-ATPase by PLM is expected to raise intracellular Na<sup>+</sup> concentration ([Na<sup>+</sup>]<sub>i</sub>), thereby decreasing the thermodynamic driving force for forward Na<sup>+</sup>/Ca<sup>2+</sup> exchange (Ca<sup>2+</sup> efflux) and increasing the driving force for reverse Na<sup>+</sup>/Ca<sup>2+</sup> exchange (Ca<sup>2+</sup> influx). Both these actions are expected to increase [Ca<sup>2+</sup>]<sub>i</sub> and SR Ca<sup>2+</sup> load, thereby enhancing cardiac contractility. Indeed, inhibition of Na<sup>+</sup>-K<sup>+</sup>-ATPase with secondary effects on NCX1 has long been proposed to be the mechanism of positive inotropy of digitalis glycosides (27).

### **Phospholemman: First Endogenous Regulator of Cardiac Na<sup>+</sup>/Ca<sup>2+</sup> Exchanger**

When PLM is overexpressed (1.4 to 3.5-fold) in adult rat left ventricular (LV) myocytes by adenovirus-mediated gene transfer (68, 89, 90), expression of SERCA2, α1- and α2-subunits of Na<sup>+</sup>-K<sup>+</sup>-ATPase, NCX1 and calsequestrin remains unchanged. As expected, Na<sup>+</sup>-K<sup>+</sup>-ATPase current (I<sub>pump</sub>) is decreased in rat myocytes overexpressing PLM, primarily as a result of decrease in V<sub>max</sub> rather than changes in apparent K<sub>m</sub> for Na<sup>+</sup> and K<sup>+</sup> (89). A totally unexpected finding is that both contraction and [Ca<sup>2+</sup>]<sub>i</sub> transient amplitudes (5.0 mM [Ca<sup>2+</sup>]<sub>o</sub>, 1 Hz, 37<sup>0</sup>C) in myocytes overexpressing PLM are lower, rather than higher, when compared to control rat myocytes overexpressing green fluorescent protein (GFP)(68). This serendipitous but critical observation is inconsistent with the theoretical prediction that inhibition of Na<sup>+</sup>-K<sup>+</sup>-ATPase by PLM leads to enhanced cardiac contractility. Because the contractile phenotype of myocytes overexpressing PLM is similar to that observed in myocytes in which NCX1 is downregulated (75), and opposite to that in which NCX1 is overexpressed (91), we were the first to propose in 2002 that PLM directly regulates NCX1 activity, independent of its effects on Na<sup>+</sup>-K<sup>+</sup>-ATPase (68). Follow-up studies in adult rat myocytes demonstrate that PLM co-localizes with NCX1 to the sarcolemma and t-tubules (90), that PLM co-immunoprecipitates with NCX1 (1, 45), that overexpression of PLM inhibits Na<sup>+</sup>/Ca<sup>2+</sup> exchange current (I<sub>NaCa</sub>)(69, 90), and that downregulation of PLM by

antisense increases  $I_{NaCa}$  (45). Using HEK293 cells that are devoid of NCX1 and PLM, we demonstrated that cells transfected with NCX1 display the characteristic  $I_{NaCa}$ , and that cells co-transfected with PLM demonstrate inhibition of  $I_{NaCa}$  as well as  $Na^+$ -dependent  $Ca^{2+}$  uptake (1). In addition, PLM co-immunoprecipitates with NCX1 in transfected HEK293 cells (1), pig sarcolemmal vesicles (1) and guinea pig ventricular myocyte membranes (82). In cardiac myocytes isolated from PLM-null mice,  $I_{NaCa}$  was higher in PLM-null myocytes (87) despite no differences in NCX1 protein levels (78). The cumulative evidence obtained from 3 model systems: adult rat myocytes, HEK293 cells and PLM-null mice are all consistent with our hypothesis that PLM directly regulates NCX1 (13).

NCX1 is a 938 amino acid (939 amino acid in the rat) peptide consisting of an extracellular N-terminal domain comprising the 1<sup>st</sup> 5 TM segments, a large intracellular loop (residues 218-764), and an intracellular C-terminal domain consisting of the last 4 TM segments (52, 56). The  $\alpha$ -repeats of TM segments 2, 3 and 7 of NCX1 are important in ion transport activity (31, 51) while the large intracellular loop contains the regulatory domains of the exchanger (35, 37, 40). Using glutathione S-transferase (GST) pulldown assay, we demonstrated that neither the N- nor the C-terminal TM domains of NCX1 associates with PLM (81). Rather, the cytoplasmic tail of PLM both physically and functionally interacts with the intracellular loop (residues 218-358) of NCX1 (81). Using overlapping NCX1 loop deletion mutants, we further showed that PLM interacts with NCX1 at 2 distinct regions encompassing residues 238-270 and 300-328 (92).

There are significant differences between the mechanisms by which PLM regulates the activities of NCX1 and  $Na^+K^+$ -ATPase. First, phosphorylation of PLM at serine<sup>68</sup> relieves its inhibition of  $Na^+K^+$ -ATPase (17, 24, 66). By contrast, PLM phosphorylated at serine<sup>68</sup> is the

active species that inhibits NCX1 (69, 87). Second, the TM segment of FXYD proteins (and by inference PLM) interacts with TM segments 2, 6 and 9 of  $\alpha$ -subunit of  $\text{Na}^+$ - $\text{K}^+$ -ATPase (36, 39, 65). By contrast, TM43, a PLM mutant with its cytoplasmic tail truncated, targets correctly to the sarcolemma (69) but does not co-immunoprecipitate NCX1 (81) and has no effect on myocyte contractility (69), suggesting little-to-no association between the TM domains of PLM and NCX1.

### **Phospholemman: Regulator of Cardiac L-type Calcium Channel**

In guinea pig cardiac myocytes, Blaise Peterson recently demonstrated that PLM co-immunoprecipitates not only NCX1 but also L-type  $\text{Ca}^{2+}$  channels ( $\text{Ca}_v1.2$ )(82). In transfected HEK293 cells and using  $\text{Ba}^{2+}$  as charge carrier, PLM modulates gating kinetics of  $\text{Ca}_v1.2$  but not  $\text{Ca}_v2.1$  (P/Q-type) or  $\text{Ca}_v2.2$  (N-type)  $\text{Ca}^{2+}$  channels (82). Specifically, PLM was found to modulate 4 important gating processes of  $\text{Ca}_v1.2$  channels: (i) activation kinetics were slowed at voltages near the threshold for channel activation; (ii) deactivation kinetics were slowed following voltage steps mimicking human cardiac action potential; (iii) voltage-dependent inactivation was enhanced at voltages corresponding to the plateau phase of the cardiac action potential; and (iv) increased number of channels enter a deep inactivated state from which recovery is slow. When a human cardiac action potential is imposed on HEK293 cells transfected with  $\text{Ca}_v1.2$  channels, PLM increases  $\text{Ca}^{2+}$  influx during the repolarization phase of the cardiac action potential (82). The role of PLM phosphorylation in the regulation of  $\text{Ca}_v1.2$  gating kinetics remains to be elucidated.

The possibility that in cardiac myocytes, PLM may potentially modulate gating kinetics of L-type  $\text{Ca}^{2+}$  channels, in addition to its known effects on  $\text{Na}^+$ - $\text{K}^+$ -ATPase and  $\text{Na}^+$ / $\text{Ca}^{2+}$  exchanger, renders the interpretation of the effects of PLM expression/phosphorylation on

cardiac contractility extremely complex. However, heterologous expression systems often do not reproduce a protein's native milieu and may distort the stoichiometry of interaction between proteins. A good example is the potentiation of  $I_{Ca}$  by adrenergic agonists, so readily observed in cardiac myocytes (63, 88), has yet to be reproduced in heterologous expression systems (34). In addition, using  $Ca^{2+}$  as charge carrier, we did not detect any differences in maximal  $I_{Ca}$  amplitude, fast and slow inactivation time constants, slope conductance and test potential at which maximal  $I_{Ca}$  occurs between WT and PLM-null myocytes (78). Therefore, the physiological significance of regulation of  $Ca_v1.2$  by PLM, while intriguing, remains to be established in cardiac myocytes.

### **Regulation of Single Myocyte Contraction by Phospholemman: $Na^+$ - $K^+$ -ATPase vs. NCX1**

In cultured myocytes isolated from PLM-null mice and expressing the phosphomimetic PLM S68E mutant,  $I_{NaCa}$  but not  $I_{pump}$  is inhibited (70). This is associated with decreased  $[Ca^{2+}]_i$  transient and contraction amplitudes (1 Hz, 37°C) measured at 5.0 but not at 1.8 mM  $[Ca^{2+}]_o$ : the phenotype that we observed when NCX1 is downregulated (75) or when PLM is overexpressed (68). By contrast, when cultured PLM-null myocytes overexpress the non-phosphorylatable PLM S68A mutant,  $I_{pump}$  but not  $I_{NaCa}$  is inhibited (70). This is associated with no changes in  $[Ca^{2+}]_i$  transient and contraction amplitudes at both  $[Ca^{2+}]_o$ . Therefore, under conditions in which  $[Ca^{2+}]_o$  is varied to manipulate the thermodynamic driving force for NCX1, regulation of single cardiac myocyte contractility by PLM is mediated by its inhibitory effects on NCX1 rather than  $Na^+$ - $K^+$ -ATPase.

When myocytes are subjected to rapid pacing (2 Hz) and isoproterenol (1  $\mu$ M) stimulation,  $[Na^+]_i$  initially increases but then starts to decline in WT but not in PLM-null myocytes (18, 80).  $[Ca^{2+}]_i$  transient and contraction amplitudes follow the time course of  $[Na^+]_i$ :

initially increase followed by decline in WT but not PLM-null myocytes. When pacing was slowed to 0.5 Hz to minimize the steep rise in  $[\text{Na}^+]_i$ , both  $[\text{Ca}^{2+}]_i$  transient and contraction amplitudes increase to a lower steady-state level without any time-dependent decline in both WT and PLM-null myocytes (80). These observations suggest that under conditions of high  $[\text{Na}^+]_i$ , phosphorylated PLM activates  $\text{Na}^+\text{-K}^+\text{-ATPase}$  to limit intracellular  $\text{Na}^+$  overload at the expense of reduced inotropy. Therefore, at the level of a single myocyte, PLM can be shown to regulate  $\text{Na}^+$  and  $\text{Ca}^{2+}$  fluxes (and hence  $[\text{Ca}^{2+}]_i$  transients and contractility), by either NCX1 or  $\text{Na}^+\text{-K}^+\text{-ATPase}$ , depending on experimental manipulations.

### **Regulation of In Vivo Contractility by Phospholemman: Studies with PLM-null Mice**

In 2005, Amy Tucker made a major contribution to the understanding of PLM and cardiac function by successfully engineering the PLM-null mouse (32). There is mild cardiac hypertrophy (3, 32, 80), at least partly due to increased fibrosis in PLM-null hearts (80) since neither LV myocyte length and width (78) nor whole cell membrane capacitance (a measure of cell surface area)(17, 78) is different between WT and PLM-null myocytes. There are no differences in protein levels of  $\alpha 1$ -,  $\alpha 2$ -,  $\beta 1$ - and  $\beta 2$ -subunits of  $\text{Na}^+\text{-K}^+\text{-ATPase}$ , SERCA2, PLB, NCX1 and calsequestrin between wild-type and congenic PLM-null hearts (3, 78). The majority of proteins that are differentially expressed between WT and PLM-null hearts are involved in cell metabolism (3).  $\text{Na}^+\text{-K}^+\text{-ATPase}$  enzymatic activity (3),  $I_{\text{pump}}$  (17, 70) and  $I_{\text{NaCa}}$  (70, 87) are higher in PLM-null hearts, as expected from the relief of inhibition of  $\text{Na}^+\text{-K}^+\text{-ATPase}$  and NCX1. There are no changes in  $I_{\text{Ca}}$  amplitudes but action potential duration is prolonged in PLM-null myocytes (78).

The effects of PLM on cardiac contractility in vivo are complicated and controversial. Using magnetic resonance imaging, Amy Tucker (32) initially reported cardiac hypertrophy and

increased ejection fraction in PLM-null hearts of mice with mixed genetic background (C57BL/6 and 129/SvJ). By contrast, in vivo hemodynamic measurements made by Mike Shattock with conductance catheter introduced by LV puncture in open-chest mice demonstrate no significant differences in measured cardiac indices between WT and PLM-null mice of congenic (C57BL/6) background (3). Our own data in closed-chest catheterized mice show slightly increased baseline +dP/dt in congenic PLM-null hearts (80). Different genetic backgrounds, non-invasive imaging vs. invasive catheterization, different anesthesia, blood loss associated with opening the chest and LV puncture, and heat dissipation in an open-chest mice, may account for these discrepancies. The weight of current evidence, however, indicates that PLM-null hearts contract just as well, if not better, than WT hearts. This is inconsistent with the expectation that with relief of inhibition of Na<sup>+</sup>-K<sup>+</sup>-ATPase, PLM-null hearts should exhibit lower contractility when compared to WT hearts.

### **Phospholemman: a Novel Cardiac Stress Protein**

When PLM is overexpressed in adult rat LV myocytes, contractility and [Ca<sup>2+</sup>]<sub>i</sub> transient amplitudes measured under physiological conditions (1.8 mM [Ca<sup>2+</sup>]<sub>o</sub>, 1 Hz, 37<sup>0</sup>C) are only slightly less than those measured in control myocytes expressing GFP (68). Likewise, contractility and [Ca<sup>2+</sup>]<sub>i</sub> transient amplitudes (1.8 mM [Ca<sup>2+</sup>]<sub>o</sub>, 1 Hz, 37<sup>0</sup>C) are similar between WT and PLM-null myocytes (78). Only when the thermodynamic driving force for NCX1 is altered by varying [Ca<sup>2+</sup>]<sub>o</sub> (0.6 or 5.0 mM) are the effects of PLM on myocyte contractility and [Ca<sup>2+</sup>]<sub>i</sub> transients evident (45, 68, 69, 78, 90). The effects of PLM on Na<sup>+</sup>-K<sup>+</sup>-ATPase are also not apparent in myocytes under resting conditions: basal [Na<sup>+</sup>]<sub>i</sub> is similar between wild-type and PLM-null myocytes (17, 80). Therefore, under resting conditions, PLM is functionally quiescent.



In the intact heart,  $\beta$ -adrenergic stimulation increases chronotropy leading to more frequent depolarizations and increased  $\text{Na}^+$  entry. In addition,  $I_{\text{Ca}}$  and SERCA2 activity are also increased in response to  $\beta$ -adrenergic stimulation, resulting in increased  $\text{Ca}^{2+}$  entry and SR  $\text{Ca}^{2+}$  loading. Elevated SR  $\text{Ca}^{2+}$  content available for release largely accounts for the enhanced inotropy associated with  $\beta$ -adrenergic stimulation. Increased  $\text{Ca}^{2+}$  entry must be balanced by greater  $\text{Ca}^{2+}$  efflux mediated by forward NCX1, thereby bringing more  $\text{Na}^+$  into the myocyte. This, if unchecked, will lead to cellular  $\text{Na}^+$  and  $\text{Ca}^{2+}$  overload. Don Bers (18) hypothesized that  $\beta$ -adrenergic agonists increase PLM phosphorylation at serine<sup>68</sup>, thereby activating  $\text{Na}^+$ - $\text{K}^+$ -ATPase and resulting in lower  $[\text{Na}^+]_i$ . The lower  $[\text{Na}^+]_i$  promotes  $\text{Ca}^{2+}$  efflux via NCX1, resulting in lower  $[\text{Ca}^{2+}]_i$  transient and contraction amplitudes (18, 80). Indeed, when hearts in vivo are stressed with maximal doses of isoproterenol, inotropy (+dP/dt) rises to a peak within 2 minutes followed by decline in WT but not PLM-null hearts (Fig. 4). Therefore, when hearts are under duress, one of the major functions of PLM is to limit  $\text{Na}^+$  and  $\text{Ca}^{2+}$  overload, thereby minimizing the risks of arrhythmogenesis apparently at the expense of reduced inotropy.

On the other hand, reduced cardiac contractility under conditions of fight or flight is clearly not in the best interests of the organism. In 2006 we proposed a coordinated paradigm in which PLM, upon phosphorylation at serine<sup>68</sup>, enhances  $\text{Na}^+$ - $\text{K}^+$ -ATPase activity to minimize risks of arrhythmogenesis but inhibits  $\text{Na}^+$ / $\text{Ca}^{2+}$  exchanger to preserve inotropy during stress (87). Our recent experiments provide support for this hypothesis. In PLM-null hearts in which isoproterenol has little-to-no effects on  $\text{Na}^+$ - $\text{K}^+$ -ATPase (17, 80), injection of recombinant adeno-associated virus (serotype 9) expressing the phosphomimetic PLM S68E mutant (rAAV9-S68E) directly into the LV resulted in expression of the mutant protein after 4-5 weeks. PLM S68E mutant inhibits NCX1 but not  $\text{Na}^+$ - $\text{K}^+$ -ATPase (70). Isoproterenol stimulation resulted in similar

increases in  $[Na^+]_i$  but higher  $+dP/dt$  in PLM-null hearts expressing PLM S68E mutant when compared to PLM-null hearts expressing GFP (Fig. 5). Therefore, inhibition of NCX1 by phosphorylated PLM preserves cardiac contractility under stressful situations.

### **FXYP Proteins in Aging, Exercise and Disease**

PLM expression is 2-fold higher in neonatal rabbit ventricular membranes and declines within 10 days to the level observed in adults (71). The decrease in PLM expression with postnatal maturation is concurrent with reduction in  $Na^+-K^+-ATPase$  and  $Na^+/Ca^{2+}$  exchanger (71) and suggests tight coordination of PLM with the 2 ion transporters. With aging, expression of PLM in sedentary rat skeletal muscle is not altered but the level of  $\alpha 1$ -subunit of  $Na^+-K^+-ATPase$  that co-immunoprecipitates with PLM increases 3-fold (61). There are no detectable changes of association of  $\alpha 2$ -subunit of  $Na^+-K^+-ATPase$  with PLM with aging (61).

Acute exercise (treadmill running) in rats increases sarcolemmal PLM in skeletal muscle by 200-350% due to translocation, but phosphorylation at serine<sup>68</sup> appears not to be altered (60). When senescent rats (26 months-old) are subjected to endurance treadmill running for 13-14 weeks, PLM in skeletal muscle is increased by 150% when compared to sedentary senescent rats (61). In addition, increased association of PLM with  $\alpha 1$ -subunit of  $Na^+-K^+-ATPase$  in skeletal muscle of senescent rats (as compared to young rats) is decreased with endurance treadmill running (61).

In 2002, using cDNA microarrays containing 86 known genes and 989 unknown cDNAs, Sehl et al. (64) are the first to report that PLM is 1 of only 19 genes to increase after myocardial infarction (MI) in the rat. We confirmed that PLM protein levels increased 2.4- and 4-fold at 3 and 7 days post-MI, respectively, in the rat (89). PLM overexpression may very well explain the depression in both  $Na^+-K^+-ATPase$  (20) and  $Na^+/Ca^{2+}$  exchanger (19, 86) activities observed in

the post-MI rat model. In rat hearts subjected to acute ischemia, PLM is phosphorylated which leads to profound activation of sarcolemmal Na<sup>+</sup>-K<sup>+</sup>-ATPase (24). In isolated perfused mouse hearts subjected to ischemia/reperfusion, protection against infarction by sildenafil is associated with increased PLM phosphorylation at serine<sup>69</sup> which enhances Na<sup>+</sup>-K<sup>+</sup>-ATPase activity during reperfusion (41). Increased Na<sup>+</sup>-K<sup>+</sup>-ATPase activity during acute ischemia ± reperfusion is critical in maintaining [Na<sup>+</sup>]<sub>i</sub> homeostasis in order to minimize the adverse effects of elevated [Na<sup>+</sup>]<sub>i</sub> on contractility and arrhythmogenesis. In human heart failure, protein levels of PLM in LV homogenates are reduced by 24% (8). In a rabbit model of volume overload heart failure which is prone to arrhythmias (57), expression of PLM is reduced by 42-48% but phosphorylation at serine<sup>68</sup> is dramatically increased (8). Thus, both altered expression and phosphorylation of PLM have been observed in various cardiac disease models. In this context, it is very relevant to note that the 2 classes of drugs that have been clinically proven to be efficacious in human heart failure, β-adrenergic blockers (lowering PKA activity) and angiotensin converting enzyme inhibitors (reducing PKC activity), both have PLM as a common target.

Sepsis is a major clinical problem which is characterized by profound hypotension, systemic vasodilatation and depression in cardiac contractility. A wide range of inflammatory cascades is activated during systemic sepsis (54). Increased nitric oxide (NO) by inducible NO synthase (iNOS) has been suggested to cause depressed cardiac contractility in sepsis (26). In this light, Helge Rasmussen demonstrated that NO stimulates Na<sup>+</sup>-K<sup>+</sup>-ATPase (22, 83) and Mike Shattock reported (in abstract form) that this is dependent on PLM. Acceleration of Na<sup>+</sup>-K<sup>+</sup>-ATPase activity by PLM may account for hyperpolarization and relaxation of vascular smooth muscle (vasodilatation) in addition to depression of cardiac contractility.

PLM has also been implicated in other diseases. For example, PLM is downregulated in layer II/III stellate neurons in patients with schizophrenia (30). Rett syndrome, an X-linked neuro-development disorder that ranks as the second most prevalent cause of mental retardation in girls (49), is due to heterozygous de novo mutations in the methyl-CpG-binding protein 2 (MeCP2) gene. MeCP2 normally represses PLM transcription through direct interactions with sequences in the PLM promoter. In patients with Rett syndrome and MeCP2-null mice, PLM is elevated in neurons in the frontal cortex and cerebellum (16, 33). Increasing neuronal PLM expression is sufficient to reduce dendritic arborization and spine formation, hallmarks of neuropathology in patients with Rett syndrome.

Dominant-negative mutation in FXYD2 ( $\gamma$ -subunit of  $\text{Na}^+$ - $\text{K}^+$ -ATPase) causes defective routing to the plasma membrane and is the cause of primary renal hypomagnesemia (44). Increased MAT-8 (FXYD3) expression is associated with tumor progression in human breast, prostate and colorectal cancers (84). Likewise, dysadherin (FXYD5) expression is altered in a wide variety of human cancers, including but not limited to breast, gastrointestinal stromal, head and neck, papillary thyroid, colorectal, non-small cell lung, and testicular cancers, and also epithelioid sarcoma and malignant melanoma (50).

### **Future Directions**

The physiological role of PLM on regulation of L-type  $\text{Ca}^{2+}$  channel needs to be established in its natural environment. The stoichiometry of interaction between PLM and  $\text{Na}^+$ - $\text{K}^+$ -ATPase, PLM and  $\text{Na}^+$ / $\text{Ca}^{2+}$  exchanger, and PLM and L-type  $\text{Ca}^{2+}$  channel, remains to be determined in cardiac myocytes. The role of PLM in regulating cardiac contractility in vivo, both in health and disease states, needs further investigation. This will likely require development of novel genetic models. The effects of oxidative stress and NO on both PLM and

Na<sup>+</sup>-K<sup>+</sup>-ATPase are just beginning to be addressed. For effective but specific drug targeting, the precise molecular interactions between PLM and Na<sup>+</sup>-K<sup>+</sup>-ATPase, and PLM and Na<sup>+</sup>/Ca<sup>2+</sup> exchanger need to be mapped out.

## **Conclusion**

FXYP proteins are emerging not only as novel endogenous regulators of ion transport but also as important targets in many human diseases including neurological, cardiac and renal diseases and a wide variety of cancers. Phospholemman (FXYP1) regulates Na<sup>+</sup>-K<sup>+</sup>-ATPase, Na<sup>+</sup>/Ca<sup>2+</sup> exchanger and possibly L-type Ca<sup>2+</sup> channel in the heart. Its effects on in vivo cardiac contractility are complex and remain to be clarified. When hearts are subjected to stress, phospholemman minimizes risks of arrhythmogenesis and preserves inotropy. Elucidating the mechanisms by which alterations or mutations of FXYP proteins are involved in human diseases will undoubtedly provide novel and rational therapeutic targets.

## **Acknowledgement**

This work was supported in part by National Institutes of Health Grants RO1-HL58672 and RO1-HL74854 (JYC), RO1-HL91096 (JER), and by an American Heart Association Scientist Development Grant F64702 (TOC).

## References

1. **Ahlers BA, Zhang XQ, Moorman JR, Rothblum LI, Carl LL, Song J, Wang J, Geddis LM, Tucker AL, Mounsey JP, and Cheung JY.** Identification of an endogenous inhibitor of the cardiac  $\text{Na}^+/\text{Ca}^{2+}$  exchanger, phospholemman. *J Biol Chem* 280: 19875-19882, 2005.
2. **Beevers AJ and Kukol A.** Secondary structure, orientation, and oligomerization of phospholemman, a cardiac transmembrane protein. *Protein Sci* 15: 1127-1132, 2006.
3. **Bell JR, Kennington E, Fuller W, Dighe K, Donoghue P, Clark JE, Jia LG, Tucker AL, Moorman JR, Marber MS, Eaton P, Dunn MJ and Shattock MJ.** Characterisation of the phospholemman knockout mouse heart: depressed left ventricular function with increased Na/K ATPase activity. *Am J Physiol* 294: H613-H621, 2008.
4. **Berry RG, Despa S, Fuller W, Bers DM, and Shattock MJ.** Differential distribution and regulation of mouse cardiac  $\text{Na}^+/\text{K}^+$ -ATPase alpha1 and alpha2 subunits in T-tubule and surface sarcolemmal membranes. *Cardiovasc Res* 73: 92-100, 2007.

5. **Bers DM.** Cardiac excitation-contraction coupling. *Nature* 415: 198-205, 2002.
6. **Bibert S, Roy S, Schaer D, Horisberger JD, and Geering K.** Phosphorylation of phospholemman (FXVD1) by protein kinases A and C modulates distinct Na,K-ATPase isozymes. *J Biol Chem* 283: 476-486, 2008.
7. **Blanco G and Mercer RW.** Isozymes of the Na-K-ATPase: heterogeneity in structure, diversity in function. *Am J Physiol* 275: F633-650, 1998.
8. **Bossuyt J, Ai X, Moorman JR, Pogwizd SM, and Bers DM.** Expression and phosphorylation of the Na-pump regulatory subunit phospholemman in heart failure. *Circ Res* 97: 558-565, 2005.
9. **Bossuyt J, Despa S, Han F, Hou Z, Robia SL, Lingrel JB, and Bers DM.** Isoform-specificity of the Na/K-ATPase association and regulation by phospholemman. *J Biol Chem* 284: 26749-26757, 2009.
10. **Bossuyt J, Despa S, Martin JL, and Bers DM.** Phospholemman phosphorylation alters its fluorescence resonance energy transfer with the Na/K-ATPase pump. *J Biol Chem* 281: 32765-32773, 2006.
11. **Chen LS, Lo CF, Numann R, and Cuddy M.** Characterization of the human and rat phospholemman (PLM) cDNAs and localization of the human PLM gene to chromosome 19q13.1. *Genomics* 41: 435-443, 1997.
12. **Chen Z, Jones LR, O'Brian JJ, Moorman JR, and Cala SE.** Structural domains in phospholemman: a possible role for the carboxyl terminus in channel inactivation. *Circ Res* 82: 367-374, 1998.

13. **Cheung JY, Rothblum LI, Moorman JR, Tucker AL, Song J, Ahlers BA, Carl LL, Wang J, and Zhang XQ.** Regulation of cardiac Na<sup>+</sup>/Ca<sup>2+</sup> exchanger by phospholemman. *Ann NY Acad Sci* 1099: 119-134, 2007.
14. **Crambert G, Fuzesi M, Garty H, Karlisch S, and Geering K.** Phospholemman (FXVD1) associates with Na,K-ATPase and regulates its transport properties. *Proc Natl Acad Sci U S A* 99: 11476-11481, 2002.
15. **Davis CE, Patel MK, Miller JR, John JE, 3rd, Jones LR, Tucker AL, Mounsey JP, and Moorman JR.** Effects of phospholemman expression on swelling-activated ion currents and volume regulation in embryonic kidney cells. *Neurochem Res* 29: 177-187, 2004.
16. **Deng V, Matagne V, Banine F, Frerking M, Ohliger P, Budden S, Pevsner J, Dissen GA, Sherman LS, and Ojeda SR.** FXVD1 is an MeCP2 target gene overexpressed in the brains of Rett syndrome patients and Mecp2-null mice. *Human molecular genetics* 16: 640-650, 2007.
17. **Despa S, Bossuyt J, Han F, Ginsburg KS, Jia LG, Kutchai H, Tucker AL, and Bers DM.** Phospholemman-phosphorylation mediates the beta-adrenergic effects on Na/K pump function in cardiac myocytes. *Circ Res* 97: 252-259, 2005.
18. **Despa S, Tucker AL, and Bers DM.** PLM-mediated activation of Na/K-ATPase limits [Na]<sub>i</sub> and inotropic state during β-adrenergic stimulation in mouse ventricular myocytes. *Circulation* 117: 1849-1855, 2008.
19. **Dixon I, Hata T, and Dhalla N.** Sarcolemmal calcium transport in congestive heart failure due to myocardial infarction in rats. *Am J Physiol Heart Circ Physiol* 262: H1387-H1394, 1992.



20. **Dixon I, Hata T, and Dhalla N.** Sarcolemmal Na<sup>+</sup>-K<sup>+</sup>-ATPase activity in congestive heart failure due to myocardial infarction. *Am Journal of Physiol Cell Physiol* 262: C664-C671, 1992.
21. **Dostanic I, Schultz Jel J, Lorenz JN, and Lingrel JB.** The alpha 1 isoform of Na,K-ATPase regulates cardiac contractility and functionally interacts and co-localizes with the Na/Ca exchanger in heart. *J Biol Chem* 279: 54053-54061, 2004.
22. **Figtree GA, Liu CC, Bibert S, Hamilton EJ, Garcia A, White CN, Chia KK, Cornelius F, Geering K, and Rasmussen HH.** Reversible oxidative modification: a key mechanism of Na<sup>+</sup>-K<sup>+</sup> pump regulation. *Circ Res* 105: 185-193, 2009.
23. **Franzin CM, Gong XM, Thai K, Yu J, and Marassi FM.** NMR of membrane proteins in micelles and bilayers: the FXYD family proteins. *Methods (San Diego, Calif)* 41: 398-408, 2007.
24. **Fuller W, Eaton P, Bell JR, and Shattock MJ.** Ischemia-induced phosphorylation of phospholemman directly activates rat cardiac Na/K-ATPase. *FASEB J* 18: 197-199, 2004.
25. **Fuller W, Howie J, McLatchie L, Weber R, Hastie CJ, Burness K, Pavlovic D, and Shattock MJ.** FXYD1 phosphorylation in vitro and in adult rat cardiac myocytes: threonine 69 is a novel substrate for protein kinase C. *Am J Physiol Cell Physiol* 296: C1346-C1355, 2009.
26. **Grocott-Mason RM and Shah AM.** Cardiac dysfunction in sepsis: new theories and clinical implications. *Intensive Care Medicine* 24: 286-295, 1998.
27. **Grupp I, Im W-B, Lee C, Lee S-W, Pecker M, and Schwartz A.** Regulation of sodium pump inhibition to positive inotropy at low concentrations of ouabain in rat heart muscle. *J Physiol (Lond)* 360: 149-160, 1985.

28. **Han F, Bossuyt J, Despa S, Tucker AL, and Bers DM.** Phospholemman phosphorylation mediates the protein kinase C-dependent effects on Na<sup>+</sup>-K<sup>+</sup> pump function in cardiac myocytes. *Circ Res* 99: 1376-1383, 2006.
29. **Han F, Tucker AL, Lingrel JB, Despa S, and Bers DM.** Extracellular potassium dependence of the Na<sup>+</sup>-K<sup>+</sup>-ATPase in cardiac myocytes: isoform specificity and effect of phospholemman. *Am J Physiol Cell Physiol* 297: C699-705, 2009.
30. **Hemby SE, Ginsberg SD, Brunk B, Arnold SE, Trojanowski JQ, and Eberwine JH.** Gene expression profile for schizophrenia: discrete neuron transcription patterns in the entorhinal cortex. *Arch Gen Psych* 59: 631-640, 2002.
31. **Iwamoto T, Uehara A, Imanaga I, and Shigekawa M.** The Na<sup>+</sup>/Ca<sup>2+</sup> exchanger NCX1 has oppositely oriented reentrant loop domains that contain conserved aspartic acids whose mutation alters its apparent Ca<sup>2+</sup> affinity. *J Biol Chem* 275: 38571-38580, 2000.
32. **Jia LG, Donnet C, Bogaev RC, Blatt RJ, McKinney CE, Day KH, Berr SS, Jones LR, Moorman JR, Sweadner KJ, and Tucker AL.** Hypertrophy, increased ejection fraction, and reduced Na-K-ATPase activity in phospholemman-deficient mice. *Am J Physiol Heart Circ Physiol* 288: H1982-1988, 2005.
33. **Jordan C, Li HH, Kwan HC, and Francke U.** Cerebellar gene expression profiles of mouse models for Rett syndrome reveal novel MeCP2 targets. *BMC Medical Genetics* 8: 36, 2007.
34. **Lemke T, Welling A, Christel CJ, Blaich A, Bernhard D, Lenhardt P, Hofmann F, and Moosmang S.** Unchanged β-adrenergic stimulation of cardiac L-type calcium channels in Ca<sub>v</sub>1.2 phosphorylation site S1928A mutant mice. *J Biol Chem* 283: 34738-34744, 2008.

35. **Levitsky DO, Nicoll DA, and Philipson KD.** Identification of the high affinity  $\text{Ca}^{2+}$ -binding domain of the cardiac  $\text{Na}^+/\text{Ca}^{2+}$  exchanger. *J Biol Chem* 269: 22847-22852, 1994.
36. **Li C, Grosdidier A, Crambert G, Horisberger JD, Michielin O, and Geering K.** Structural and functional interaction sites between Na,K-ATPase and FXYD proteins. *J Biol Chem* 279: 38895-38902, 2004.
37. **Li Z, Nicoll DA, Collins A, Hilgemann D, Filoteo A, Penniston J, Weiss J, Tomich J, and Philipson KD.** Identification of a peptide inhibitor of the cardiac sarcolemmal  $\text{Na}^+/\text{Ca}^{2+}$  exchanger. *J Biol Chem* 266: 1014-1020, 1991.
38. **Lindemann JP.**  $\alpha$ -adrenergic stimulation of sarcolemmal protein phosphorylation and slow responses in intact myocardium. *J Biol Chem* 261: 4860-4867, 1986.
39. **Lindzen M, Gottschalk KE, Fuzesi M, Garty H, and Karlish SJ.** Structural interactions between FXYD proteins and  $\text{Na}^+/\text{K}^+$ -ATPase:  $\alpha/\beta/\text{FXYD}$  subunit stoichiometry and cross-linking. *J Biol Chem* 281: 5947-5955, 2006.
40. **Maack C, Ganesan A, Sidor A, and O'Rourke B.** Cardiac sodium-calcium exchanger is regulated by allosteric calcium and exchanger inhibitory peptide at distinct sites. *Circ Res* 96: 91-99, 2005.
41. **Madhani M, Hall AR, Cuello F, Charles RL, Burgoyne JR, Fuller W, Hobbs AJ, Shattock MJ, and Eaton P.** Phospholemman Ser-69 phosphorylation contributes to sildenafil-induced cardioprotection against reperfusion injury. *Am J Physiol Heart Circ Physiol*, in press 2010.
42. **Mahmoud YA, Vorum H, and Cornelius F.** Purification of a phospholemman-like protein from shark rectal glands. *J Biol Chem* 275: 35969-35977, 2000.

43. **McDonough A, Zhang Y, Shin V, and Frank JS.** Subcellular distribution of sodium pump isoform subunits in mammalian cardiac myocytes. *Am J Physiol Cell Physiol* 270: C1221-C1227, 1996.
44. **Meij IC, Koenderink JB, van Bokhoven H, Assink KF, Groenestege WT, de Pont JJ, Bindels RJ, Monnens LA, van den Heuvel LP, and Knoers NV.** Dominant isolated renal magnesium loss is caused by misrouting of the Na<sup>+</sup>-K<sup>+</sup>-ATPase  $\gamma$ -subunit. *Nature Genetics* 26: 265-266, 2000.
45. **Mirza MA, Zhang XQ, Ahlers BA, Qureshi A, Carl LL, Song J, Tucker AL, Mounsey JP, Moorman JR, Rothblum LI, Zhang TS, and Cheung JY.** Effects of phospholemman downregulation on contractility and [Ca<sup>2+</sup>]<sub>i</sub> transients in adult rat cardiac myocytes. *Am J Physiol Heart Circ Physiol* 286: H1322-1330, 2004.
46. **Moorman JR, Ackerman SJ, Kowdley GC, Griffin M, Mounsey JP, Chen Z, Cala SE, O'Brian JJ, Szabo G, and Jones LR.** Unitary anion currents through phospholemman channel molecules. *Nature* 377: 737-740, 1995.
47. **Morales-Mulia M, Pasantes-Morales H, and Moran J.** Volume sensitive efflux of taurine in HEK 293 cells overexpressing phospholemman. *Biochem Biophys Acta* 1496: 252-260, 2000.
48. **Mounsey JP, John III J, Helmke S, Bush E, Gilbert J, Roses A, Perryman M, Jones LR, and Moorman JR.** Phospholemman is a substrate for myotonic dystrophy protein kinase. *J Biol Chem* 275: 23362-23367, 2000.
49. **Naidu S.** Rett syndrome: a disorder affecting early brain growth. *Ann Neurol* 42: 3-10, 1997.

50. **Nam JS, Hirohashi S, and Wakefield LM.** Dysadherin: a new player in cancer progression. *Cancer Letters* 255: 161-169, 2007.
51. **Nicoll DA, Hryshko LV, Matsuoka S, Frank JS, and Philipson KD.** Mutation of amino acid residues in the putative transmembrane segments of the cardiac sarcolemmal Na<sup>+</sup>/Ca<sup>2+</sup> exchanger. *J Biol Chem* 271: 13385-13391, 1996.
52. **Nicoll DA, Ottolia M, Lu L, Lu Y, and Philipson KD.** A new topological model of the cardiac sarcolemmal Na<sup>+</sup>/Ca<sup>2+</sup> exchanger. *J Biol Chem* 274: 910-917, 1999.
53. **Palmer CJ, Scott BT, and Jones LR.** Purification and complete sequence determination of the major plasma membrane substrate for cAMP-dependent protein kinase and protein kinase C in myocardium. *J Biol Chem* 266: 11126-11130, 1991.
54. **Parrillo JE.** Pathogenetic mechanisms of septic shock. *New Eng J Med* 328: 1471-1477, 1993.
55. **Pavlovic D, Fuller W, and Shattock MJ.** The intracellular region of FXYP1 is sufficient to regulate cardiac Na/K ATPase. *FASEB J* 21: 1539-1546, 2007.
56. **Philipson KD and Nicoll DA.** Sodium-calcium exchange: a molecular perspective. *Annu Rev Physiol* 62: 111-133, 2000.
57. **Pogwizd SM, Schlotthauer K, Li L, Yuan W, and Bers DM.** Arrhythmogenesis and contractile dysfunction in heart failure: Roles of sodium-calcium exchange, inward rectifier potassium current, and residual  $\beta$ -adrenergic responsiveness. *Circ Res* 88: 1159-1167, 2001.
58. **Presti CF, Jones LR, and Lindemann JP.** Isoproterenol-induced phosphorylation of a 15-kilodalton sarcolemmal protein in intact myocardium. *J Biol Chem* 260: 3860-3867, 1985.

59. **Presti CF, Scott BT, and Jones LR.** Identification of an endogenous protein kinase C activity and its intrinsic 15-kilodalton substrate in purified canine cardiac sarcolemmal vesicles. *J Biol Chem* 260: 13879-13889, 1985.
60. **Rasmussen MK, Kristensen M, and Juel C.** Exercise-induced regulation of phospholemman (FXVD1) in rat skeletal muscle: implications for Na<sup>+</sup>-K<sup>+</sup>-ATPase activity. *Acta Physiol (Oxford, England)* 194: 67-79, 2008.
61. **Reis J, Zhang L, Cala S, Jew KN, Mace LC, Chung L, Moore RL, and Ng YC.** Expression of phospholemman and its association with Na<sup>+</sup>-K<sup>+</sup>-ATPase in skeletal muscle: effects of aging and exercise training. *J Appl Physiol* 99: 1508-1515, 2005.
62. **Rembold CM, Ripley ML, Meeks MK, Geddis LM, Kutchai HC, Marassi FM, Cheung JY, and Moorman JR.** Serine68 phospholemman phosphorylation during forskolin-induced swine carotid artery relaxation. *J Vasc Res* 42: 483-491, 2005.
63. **Reuter H.** The dependence of slow inward current in Purkinje fibres on the extracellular calcium-concentration. *J Physiol* 192: 479-492, 1967.
64. **Sehl PD, Tai JT, Hillan KJ, Brown LA, Goddard A, Yang R, Jin H, and Lowe DG.** Application of cDNA microarrays in determining molecular phenotype in cardiac growth, development, and response to injury. *Circulation* 101: 1990-1999, 2000.
65. **Shinoda T, Ogawa H, Cornelius F, and Toyoshima C.** Crystal structure of the sodium-potassium pump at 2.4 Å resolution. *Nature* 459: 446-450, 2009.
66. **Silverman BZ, Fuller W, Eaton P, Deng J, Moorman JR, Cheung JY, James AF, and Shattock MJ.** Serine68 phosphorylation of phospholemman: acute isoform-specific activation of cardiac Na/K ATPase. *Cardiovasc Res* 65: 93-103, 2005.

67. **Simmerman HK and Jones LR.** Phospholamban: protein structure, mechanism of action, and role in cardiac function. *Physiol Rev* 78: 921-947, 1998.
68. **Song J, Zhang XQ, Carl LL, Qureshi A, Rothblum LI, and Cheung JY.** Overexpression of phospholemman alter contractility and  $[Ca^{2+}]_i$  transients in adult rat myocytes. *Am Journal of Physiol Heart Circ Physiol* 283: H576-H583, 2002.
69. **Song J, Zhang XQ, Ahlers BA, Carl LL, Wang J, Rothblum LI, Stahl RC, Mounsey JP, Tucker AL, Moorman JR, and Cheung JY.** Serine 68 of phospholemman is critical in modulation of contractility,  $[Ca^{2+}]_i$  transients, and  $Na^+/Ca^{2+}$  exchange in adult rat cardiac myocytes. *Am J Physiol Heart Circ Physiol* 288: H2342-2354, 2005.
70. **Song J, Zhang XQ, Wang J, Cheskis E, Chan TO, Feldman AM, Tucker AL, and Cheung JY.** Regulation of cardiac myocyte contractility by phospholemman:  $Na^+/Ca^{2+}$  exchanger vs.  $Na^+-K^+-ATPase$ . *Am J Physiol Heart Circ Physiol* 295: H1615-H1625, 2008.
71. **Srivastava S, Cala SE, Coetzee WA, and Artman M.** Phospholemman expression is high in the newborn rabbit heart and declines with postnatal maturation. *Biochem Biophys Res Comm* 355: 338-341, 2007.
72. **Sweadner KJ, Herrera V, Amato S, Moellmann A, Gibbons D, and Repke K.** Immunologic identification of  $Na^+$ ,  $K^+-ATPase$  isoforms in myocardium. *Circ Res* 74: 669-678, 1994.
73. **Sweadner KJ and Rael E.** The FXYD gene family of small ion transport regulators or channels: cDNA sequence, protein signature sequence, and expression. *Genomics* 68: 41-56, 2000.

74. **Swift F, Tovsrud N, Enger UH, Sjaastad I, and Sejersted OM.** The Na<sup>+</sup>-K<sup>+</sup>-ATPase  $\alpha$ 2-isoform regulates cardiac contractility in rat cardiomyocytes. *Cardiovasc Res* 75: 109-117, 2007.
75. **Tadros GM, Zhang XQ, Song J, Carl LL, Rothblum LI, Tian Q, Dunn J, Lytton J, and Cheung JY.** Effects of Na<sup>+</sup>/Ca<sup>2+</sup> exchanger downregulation on contractility and [Ca<sup>2+</sup>]<sub>i</sub> transients in adult rat myocytes. *Am J Physiol Heart Circ Physiol* 283: H1616-1626, 2002.
76. **Teriete P, Franzin CM, Choi J, and Marassi FM.** Structure of the Na,K-ATPase regulatory protein FXYD1 in micelles. *Biochemistry* 46: 6774-6783, 2007.
77. **Teriete P, Thai K, Choi J, and Marassi FM.** Effects of PKA phosphorylation on the conformation of the Na,K-ATPase regulatory protein FXYD1. *Biochim Biophys Acta* 1788:2462-2470, 2009.
78. **Tucker AL, Song J, Zhang XQ, Wang J, Ahlers BA, Carl LL, Mounsey JP, Moorman JR, Rothblum LI, and Cheung JY.** Altered contractility and [Ca<sup>2+</sup>]<sub>i</sub> homeostasis in phospholemman-deficient murine myocytes: Role of Na<sup>+</sup>/Ca<sup>2+</sup> exchange. *Am J Physiol Heart Circ Physiol* 291: H2199-H2209, 2006.
79. **Walaas SI, Czernik AJ, Olstad OK, Sletten K, and Walaas O.** Protein kinase C and cyclic AMP-dependent protein kinase phosphorylate phospholemman, an insulin and adrenaline-regulated membrane phosphoprotein, at specific sites in the carboxy terminal domain. *Biochem J* 304: 635-640, 1994.
80. **Wang J, Gao E, Song J, Zhang XQ, Li J, Koch WJ, Tucker AL, Philipson KD, Chan TO, Feldman AM, and Cheung JY.** Phospholemman and  $\beta$ -adrenergic stimulation in the heart. *Am J Physiol Heart Circ Physiol* 298: H807-H815, 2010.



81. **Wang J, Zhang XQ, Ahlers BA, Carl LL, Song J, Rothblum LI, Stahl RC, Carey DJ, and Cheung JY.** Cytoplasmic tail of phospholemman interacts with the intracellular loop of the cardiac  $\text{Na}^+/\text{Ca}^{2+}$  exchanger. *J Biol Chem* 281: 32004-32014, 2006.
82. **Wang X, Gao G, Guo K, Yarotsky V, Huang C, Elmslie KS, and Peterson BZ.** Phospholemman modulates the gating of cardiac L-type calcium channels. *Biophys J* 98: 1149-1159, 2010.
83. **William M, Vien J, Hamilton E, Garcia A, Bundgaard H, Clarke RJ, and Rasmussen HH.** The nitric oxide donor sodium nitroprusside stimulates the  $\text{Na}^+-\text{K}^+$  pump in isolated rabbit cardiac myocytes. *J Physiol* 565: 815-825, 2005.
84. **Yamamoto H, Okumura K, Toshima S, Mukaisho K, Sugihara H, Hattori T, Kato M, and Asano S.** FXDY3 protein involved in tumor cell proliferation is overproduced in human breast cancer tissues. *Biological Pharmaceutical Bull* 32: 1148-1154, 2009.
85. **Zahler R, Gilmore-Hebert M, Baldwin JC, Franco K, and Benz EJ, Jr.** Expression of alpha isoforms of the Na,K-ATPase in human heart. *Biochim Biophys Acta* 1149: 189-194, 1993.
86. **Zhang XQ, Tillotson DL, Moore RL, Zelis R, and Cheung JY.**  $\text{Na}^+/\text{Ca}^{2+}$  exchange currents and SR  $\text{Ca}^{2+}$  contents in postinfarction myocytes. *Am Journal of Physiol* 271: C1800-C1807, 1996.
87. **Zhang XQ, Ahlers BA, Tucker AL, Song J, Wang J, Moorman JR, Mounsey JP, Carl LL, Rothblum LI, and Cheung JY.** Phospholemman inhibition of the cardiac  $\text{Na}^+/\text{Ca}^{2+}$  exchanger. Role of phosphorylation. *J Biol Chem* 281: 7784-7792, 2006.
88. **Zhang XQ, Moore RL, Tillotson DL, and Cheung JY.** Calcium currents in postinfarction rat cardiac myocytes. *Am J Physiol* 269: C1464-1473, 1995.

89. **Zhang XQ, Moorman JR, Ahlers BA, Carl LL, Lake DE, Song J, Mounsey JP, Tucker AL, Chan YM, Rothblum LI, Stahl RC, Carey DJ, and Cheung JY.** Phospholemman overexpression inhibits Na<sup>+</sup>-K<sup>+</sup>-ATPase in adult rat cardiac myocytes: Relevance to decreased Na<sup>+</sup> pump activity in post-infarction myocytes. *J Appl Physiol* 100: 212-220, 2006.
90. **Zhang XQ, Qureshi A, Song J, Carl LL, Tian Q, Stahl RC, Carey DJ, Rothblum LI, and Cheung JY.** Phospholemman modulates Na<sup>+</sup>/Ca<sup>2+</sup> exchange in adult rat cardiac myocytes. *Am J Physiol Heart Circ Physiol* 284: H225-233, 2003.
91. **Zhang XQ, Song J, Rothblum LI, Lun M, Wang X, Ding F, Dunn J, Lytton J, McDermott PJ, and Cheung JY.** Overexpression of Na<sup>+</sup>/Ca<sup>2+</sup> exchanger alters contractility and SR Ca<sup>2+</sup> content in adult rat myocytes. *Am J Physiol Heart Circ Physiol* 281: H2079-2088, 2001.
92. **Zhang XQ, Wang J, Carl LL, Song J, Ahlers BA, and Cheung JY.** Phospholemman regulates cardiac Na<sup>+</sup>/Ca<sup>2+</sup> exchanger by interacting with the exchanger's proximal linker domain. *Am J Physiol Cell Physiol* 296: C911-921, 2009.

### **Figure Legends.**

**Figure 1. Molecular model of phospholemman.** Nuclear magnetic resonance studies of highly purified phospholemman in micelles revealed 4 helices of the protein with a single transmembrane domain (after Francesca Marassi)(23, 76). The FXYD motif is in the extracellular domain and the important serine<sup>63</sup> and serine<sup>68</sup> are in the cytoplasm.

**Figure 2. Phospholemman co-localizes with  $\alpha 1$ -subunit of  $\text{Na}^+\text{-K}^+\text{-ATPase}$ .** Indirect immunofluorescence of adult rat ventricular myocytes doubly labeled with mouse monoclonal antibody against  $\alpha 1$ -subunit of  $\text{Na}^+\text{-K}^+\text{-ATPase}$  (A) and rabbit polyclonal anti-PLM antibody (B) are shown. Primary antibodies are visualized with Alexa Fluor 488-labeled goat anti-mouse IgG (A) and Alexa Fluor 594- labeled goat anti-rabbit IgG (B). Note the orange color in the merged image (C), indicating co-localization of PLM and  $\text{Na}^+\text{-K}^+\text{-ATPase}$ . Bar = 5  $\mu\text{m}$ .

**Figure 3. Cardiac excitation-contraction coupling.** Membrane depolarization is initiated by opening of the  $\text{Na}^+$  channel (not shown) with  $\text{Na}^+$  entry. Extracellular  $\text{Ca}^{2+}$  enters via L-type  $\text{Ca}^{2+}$  channel ( $I_{\text{Ca}}$ ) and  $\text{Na}^+/\text{Ca}^{2+}$  exchanger (NCX1), causing  $\text{Ca}^{2+}$  release from the ryanodine receptor (RyR) in the sarcoplasmic reticulum (SR).  $\text{Ca}^{2+}$  binds to troponin and initiates myofilament contraction. During diastole,  $\text{Ca}^{2+}$  is pumped back to the SR by SR  $\text{Ca}^{2+}\text{-ATPase}$  (SERCA) under the control of phospholamban (PLB). A small amount of  $\text{Ca}^{2+}$  is also taken up by the mitochondrial  $\text{Ca}^{2+}$  uniporter. The amount of  $\text{Ca}^{2+}$  that has entered during systole is extruded by  $\text{Na}^+/\text{Ca}^{2+}$  exchanger and to a lesser extent, by sarcolemmal  $\text{Ca}^{2+}\text{-ATPase}$ .  $\text{Na}^+$  that has entered via  $\text{Na}^+$  channel and  $\text{Na}^+/\text{Ca}^{2+}$  exchanger is pumped out by  $\text{Na}^+\text{-K}^+\text{-ATPase}$ . Repolarization is mediated by opening of  $\text{K}^+$  channels (only the transient outward  $\text{K}^+$  current  $I_{\text{to}}$  responsible for early repolarization is shown). Phospholemman (PLM) associates with and is an endogenous regulator of  $\text{Na}^+/\text{Ca}^{2+}$  exchanger,  $\text{Na}^+\text{-K}^+\text{-ATPase}$  and possibly L-type  $\text{Ca}^{2+}$  channel.  $\text{Na}^+/\text{Ca}^{2+}$  exchanger is depicted as operating in the forward mode ( $\text{Ca}^{2+}$  efflux) in the sarcolemma and reverse mode ( $\text{Ca}^{2+}$  influx) in the t-tubules. Broken arrows point to ion transporters, ion channels and myofilaments that are altered after myocardial infarction.

**Figure 4. Effects of activation of  $\text{Na}^+\text{-K}^+\text{-ATPase}$  by phosphorylated phospholemman on  $\beta$ -adrenergic response in vivo.** Shown are normalized in vivo hemodynamics (+dP/dt) of anesthetized wild-type ( $\bullet$ ; n=9) and phospholemman-null ( $\square$ ; n=14) mice after stimulation with 25 ng of

isoproterenol. Note time-dependent decline of +dP/dt in wild-type but not phospholemman-null hearts. Phospholemman phosphorylated at serine<sup>68</sup> activates Na<sup>+</sup>-K<sup>+</sup>-ATPase (17, 80), leading to decreases in [Na<sup>+</sup>]<sub>i</sub> in wild-type but not phospholemman-null cardiac myocytes (18, 80).

**Figure 5. Effects of inhibition of Na<sup>+</sup>/Ca<sup>2+</sup> exchanger by phosphorylated phospholemman on β-adrenergic response in vivo.** Left ventricles of phospholemman-null (KO) mice are injected with recombinant adeno-associated virus, serotype 9, expressing either green fluorescent protein (GFP)(□; n=5) or the phosphomimetic phospholemman S68E mutant (●; n=7). S68E mutant inhibits Na<sup>+</sup>/Ca<sup>2+</sup> exchanger but has no effect on Na<sup>+</sup>-K<sup>+</sup>-ATPase (69, 70, 87). Five weeks after virus injection, in vivo hemodynamics (+dP/dt) are measured in anesthetized mice (80). Note with increasing doses of isoproterenol, KO-S68E hearts contract significantly better than KO-GFP hearts. Since isoproterenol has no effect on Na<sup>+</sup>-K<sup>+</sup>-ATPase in phospholemman-null myocytes (17, 80), [Na<sup>+</sup>]<sub>i</sub> is similar between KO-S68E and KO-GFP hearts (data not shown). Enhanced contractility in KO-S68E hearts is therefore due to inhibition of Na<sup>+</sup>/Ca<sup>2+</sup> exchanger by the S68E mutant.