

7-22-2018

## Another "D" in MUDPILES? A Review of Diet-Associated Nondiabetic Ketoacidosis.

Waqas Ullah  
*Abington Jefferson Health*

Mohsin Hamid  
*Abington Jefferson Health*

Hafez Mohammad Ammar Abdullah  
*University of South Dakota*

Mamoon Ur Rashid  
*Florida Hospital*

Faisal Inayat  
*Allama Iqbal Medical College*  
Follow this and additional works at: <https://jdc.jefferson.edu/abingtonfp>

 Part of the [Internal Medicine Commons](#)

[Let us know how access to this document benefits you](#)

---


### Recommended Citation

Ullah, Waqas; Hamid, Mohsin; Ammar Abdullah, Hafez Mohammad; Ur Rashid, Mamoon; and Inayat, Faisal, "Another "D" in MUDPILES? A Review of Diet-Associated Nondiabetic Ketoacidosis." (2018). *Abington Jefferson Health Papers*. Paper 5.  
<https://jdc.jefferson.edu/abingtonfp/5>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Abington Jefferson Health Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

# Another “D” in MUDPILES? A Review of Diet-Associated Nondiabetic Ketoacidosis

Journal of Investigative Medicine High  
Impact Case Reports  
Volume 6: 1–8  
© 2018 American Federation for  
Medical Research  
DOI: 10.1177/2324709618796261  
journals.sagepub.com/home/hic  


Waqas Ullah, MD<sup>1</sup>, Mohsin Hamid, MD<sup>1</sup>,  
Hafez Mohammad Ammar Abdullah, MD<sup>2</sup>,  
Mamoon Ur Rashid, MD<sup>3</sup>, and Faisal Inayat, MBBS<sup>4</sup>

## Abstract

Ketogenic diet or very-low-carbohydrate diet gained widespread popularity in the 1990s due to their favorable effects on weight loss and diabetes among others with good short-term safety data. People on ketogenic diets exist in a state of “dietary ketosis” in which the body production of ketone is equal to consumption and no harmful effects of ketonemia occur. However, in face of stress, the harmless “dietary ketosis” can lead to profound acid-base disturbances due to massive overproduction of ketone bodies that overwhelms the acid buffer system of the body. A handful of case reports have been published on this topic calling the safety of ketogenic diet into question. In this article, we chronicle a unique case of ketogenic (Atkins) diet-associated ketoacidosis, and we present a comprehensive literature review on the etiology of ketoacidosis.

## Keywords

Diet, Ketoacidosis, NDKA

Received June 18, 2018. Revised July 17, 2018. Accepted July 22, 2018.

## Introduction

Ketone bodies are formed by the deamination of amino acids and degradation of fatty acids in times of severe stress, diabetes, alcoholism, starvation, and pregnancy.<sup>1</sup> Rarely, a low-carbohydrate ketogenic diet can also give rise to ketoacidosis.<sup>2</sup> We present a case of a young nonpregnant female with no history of alcoholism, diabetes, or starvation presenting with dialysis-refractory metabolic acidosis due to Atkins diet-associated ketoacidosis. Urgent intubation, intensive care, and aggressive fluid resuscitation with insulin therapy led to the successful recovery of the patient. We also performed an extensive literature search and highlighted all reported literature of nondiabetic ketoacidosis (NDKA).

## Case Presentation

A 33-year-old Caucasian female was brought to the emergency department with possible syncope following lethargy and extreme exhaustion. Her mother found her on the floor of the restroom after hearing her falling down. According to her parents, she had mild flu-like symptoms, low-grade fever, and multiple episodes of nonbilious vomiting for 3 days before presentation. Her parents denied her ingestion of any medications or toxic substances intentionally or accidentally, and she did not have a past history of suicide attempts

or ideation. Past medical history was only significant for high-functioning autism; she worked as a cashier at a fast food restaurant and was living with her parents. Her medications included methylphenidate and sertraline for years without any recent changes.

On presentation, her vitals included temperature 98.2°F, blood pressure 140/71 mm Hg, heart rate 136 beats per minute, respiratory rate 38/min, and oxygen saturation of 96% on ambient air. Examination revealed a Glasgow Coma Scale score of 10/15; mucous membranes were dry, and skin was cold to touch with decreased turgor. Breathing was deep and labored, chest was otherwise clear to auscultation; gastrointestinal and cardiovascular examinations were unremarkable.

## Investigations

The laboratory values are shown in Table 1.

<sup>1</sup>Abington Memorial Hospital, Abington, PA, USA

<sup>2</sup>University of South Dakota, Vermillion, SD, USA

<sup>3</sup>Florida Hospital, Orlando, FL, USA

<sup>4</sup>Jinnah Hospital, Allama Iqbal Medical College, Lahore, Pakistan

## Corresponding Author:

Waqas Ullah, Abington Memorial Hospital, 1200 Old York Road, Abington, PA 19001-3720, USA.

Email: waqasullah.dr@gmail.com



**Table 1.** Laboratory Studies of Our Patient While She Was in the Intensive Care Unit.

|  |                           |
|--|---------------------------|
| Complete blood count                   |                           |
| Hemoglobin (g/dL)                      | 12.2 (12-16)              |
| White blood cells (/cmm)               | 34 800 (4000-12 000)      |
| Platelet count (/cmm)                  | 578 000 (140 000-400 000) |
| Differential type                      |                           |
| Neutrophils (%)                        | 91                        |
| Lymphocytes (%)                        | 2                         |
| Monocytes (%)                          | 7                         |
| Chemistry                              |                           |
| Glucose (mg/dL)                        | 126 (70-100)              |
| Serum osmolarity (mOsm/kg)             | 323 (275-295)             |
| Sodium (mmol/L)                        | 141 (135-145)             |
| Potassium (mmol/L)                     | 5.7 (3.5-5.1)             |
| Chloride (mmol/L)                      | 111 (98-110)              |
| Blood urea nitrogen (mg/dL)            | 17 (<23)                  |
| Creatinine (mg/dL)                     | 1.32 (<1.11)              |
| Bicarbonate (mmol/L)                   | <5 (20-31)                |
| AST (U/L)                              | 21 (4-34)                 |
| ALT (U/L)                              | 20 (<55)                  |
| ALP (U/L)                              | 130 (40-150)              |
| Total bilirubin (mg/dL)                | 0.2 (0.2-1.2)             |
| HbA1c (mg/dL)                          | 5.6                       |
| Serum lactate (mEq/L)                  | 1.1 (normal <2.0)         |
| Anion gap                              | 27 (8-12)                 |
| Arterial blood gas                     |                           |
| pH                                     | 6.8 (7.35-7.45)           |
| pCO <sub>2</sub> (mm Hg)               | 16.9 (35-45)              |
| pO <sub>2</sub> (mm Hg)                | 120 (80-100)              |
| HCO <sub>3</sub> <sup>-</sup> (mmol/L) | 2.6 (21-29)               |
| SaO <sub>2</sub> (%)                   | 97.4 (95-100)             |

Abbreviations: AST, aspartate aminotransferase; ALT, alanine aminotransferase; ALP, alkaline phosphatase; HbA1c, hemoglobin A1c; pCO<sub>2</sub>, partial pressure of carbon dioxide; pO<sub>2</sub>, partial pressure of oxygen; HCO<sub>3</sub><sup>-</sup>, bicarbonate; SaO<sub>2</sub>, oxygen saturation.

Serum and urine toxicology was negative for alcohol, salicylate, acetaminophen, and other toxic alcohols. Urine pregnancy test was negative; however, urinalysis was positive for ketones (+4). Computed tomography scan of head and chest X-ray were unremarkable.

## Differential Diagnosis

Initial suspicion of the cerebrovascular accident was ruled out by the normal computed tomography scan of the head. Sepsis, diabetic ketoacidosis, poisoning with toxic alcohols, and pregnancy-induced ketosis were considered, but they were ruled out with the appropriate studies.

## Treatment

She was initially stabilized with aggressive volume resuscitation, sodium bicarbonate therapy, and ventilator support.

Broad-spectrum antibiotics (vancomycin and piperacillin/ticarcillin) were started due to the possibility of sepsis from unknown origin. Persistent metabolic acidosis in spite of bicarbonate therapy led to one session of hemodialysis that improved serum bicarbonate levels to 16 mEq/L from 2.5 mmol/L. After a few hours of hemodialysis, repeat laboratory tests showed worsening acidosis with serum bicarbonate dipping down to 12 mmol/L. At this point, endocrinology was consulted and serum was tested for possible ketonemia, which turned out to have β-hydroxybutyrate (β-hB) levels of 120.1 mg/dL (normally less than 3 mg/dL). On further questioning, her parents mentioned that she was on a “weight-loss diet” for about 2 months. Suspecting the diagnosis of ketogenic diet-induced ketoacidosis, she was started on intravenous (IV) insulin along with dextrose 10%.


## Outcome and Follow-up

She improved rapidly after the start of IV insulin and dextrose infusions. Metabolic acidosis and anion gap (AG) resolved within a day and β-hB levels normalized. She was then extubated and shifted to a general medicine floor on the second day of her presentation. On further questioning, she admitted to following a strict commercially available ketogenic diet (Atkins diet) for the past 2 months. She was discharged home in a stable condition on day 5 of hospitalization with the diagnosis of the ketogenic diet (Atkins diet)-induced euglycemic metabolic ketoacidosis. A follow-up after 3 months revealed that she was doing well and was not following the Atkins diet anymore.

## Discussion

Over the past century, numerous different diets have been studied and tried to help weight loss. The 2 basic diets popularized are the low-fat and the low-carbohydrate diet (LCD). The LCD has increased popularity in the latter part of the past century, especially after the introduction of Atkins diet by R. C. Atkins in 1972.<sup>3</sup> Atkins diet is a special type of LCD, which is protein rich and has high-fat contents. The beneficial effect of LCD in an Atkins regime is believed to be due to the use of higher energy ketones rather than glucose for the production of energy. Even though studies have shown that LCD is effective for weight loss and causes improvement in cardiovascular risk factors, there are still controversies surrounding the use of this diet, one of the most serious of which is NDKA.<sup>4</sup>

NDKA is the presence of ketoacidosis in the absence of diabetes and is typically associated with starvation, alcoholism, hyperthyroidism, and lactation. However, this can also occur rarely with LCD, especially in patients with comorbid conditions and intercurrent illness. Our case is unique in the sense that it occurred in an otherwise healthy young female with no significant comorbidities.




**Figure 1.** Flow sheet of the pathophysiology of nondiabetic ketoacidosis.

There are a few mechanisms by which an LCD induces ketosis. Normally in high-carbohydrate states, aerobic glycolysis produces citrate as one of the intermediate metabolites. This citrate is a signal for a high-glucose state and inhibits the activity of carnitine palmitoyltransferase complex-1 (CPT-1).<sup>2</sup> CPT-1 complex is responsible for the  $\beta$ -oxidation of fatty acids, and its inhibition prevents the metabolism of fatty acids and the production of ketones in carbohydrate-rich states. The opposite occurs in low-carbohydrate states. With no citrate, the CPT-1 is activated and extensive  $\beta$ -oxidation of fatty acids occurs. The liver then uses this energy to make ketone bodies, which provide energy to important organs, including the brain, heart, and kidneys.<sup>2</sup> Moreover, fat-rich diet and LCD can enhance glucagon secretion and lower insulin secretion, and a high glucagon-to-insulin ratio also promotes ketosis.<sup>5,6</sup> Excessive accumulation of ketone bodies in the body occurs when its production is greater than consumption, leading to ketoacidosis. We believe that the low-carbohydrate and high-plasma fatty acid concentrations, compounded by the absence of carbohydrate-induced inhibition of  $\beta$ -oxidation of fatty acids might be the cause of NDKA in patients on an Atkins diet.

The pathophysiology of LCD or Atkins diet-induced ketoacidosis is shown in Figure 1.

A comprehensive literature search using the MEDLINE databases (PubMed and Ovid) showed only a handful of cases on ketoacidosis in nondiabetic patients. Different Medical Subject Headings (MeSH) like “non-diabetic,” “euglycemic,” “normoglycemic,” “normal sugar levels,” “ketoacidosis,” “ketonuria,” “ketonemia,” “ketosis,” “metabolic acidosis,” “fatty acids,” “metabolites,” “acetoacetate,” and “beta-hydroxybutyrate” were combined using Boolean operators. A total of 127 cases were initially obtained; after removing 36 duplicate studies, 93 studies were read in full-text form. Twenty-seven out of 93 cases were found relevant to our study, but only 22 cases had sufficient data for extraction.<sup>2,5-25</sup>

Further stratification of the selected cases showed that NDKA was most commonly reported in chronic alcoholics (5 case reports including 9 patients) and LCD users (5 cases).<sup>2,6,7,9,11,17,21</sup> Only 3 cases were reported in pregnancy and lactation each.<sup>12,13,15</sup> It is interesting to note that the pregnant and lactating patients had overriding starvation or low-carbohydrate intake as precipitating factors for ketoacidosis.



**Figure 2.** Flow sheet of studies on nondiabetic ketoacidosis.

There was only a single case of euglycemic ketoacidosis reported with diets like Dukan diet, diet coke, and fruitarian diet each and a single case each with hyperthyroidism, bariatric surgery, and salicylate use<sup>2224</sup> (Figure 2). Most of these patients had an underlying condition like pregnancy, asthma, pneumonia, or hyperthyroidism, which might have made them vulnerable to develop ketoacidosis. Our patient is unique as she was a previously healthy individual who presented as syncope mimicking cerebrovascular accident. The underlying cause was the Atkins diet, which has never been reported before as the cause of severe NDKA.

NDKA was most commonly reported in female patients ( $n = 16$ ) when compared with male patients ( $n = 11$ ). The age distribution ranged from 5 to 79 years. More than 80% of the patients were at the extreme of ages and with other comorbid conditions signifying that stressful or lower metabolic reserve makes patients susceptible to NDKA. Our patient,

however, was unique that flu was the only association, and she was relatively younger than the majority of reported patients. Most patients presented with gastrointestinal symptoms like nausea, vomiting, and diarrhea ( $n = 13$ ) or mild central symptoms like a headache, dizziness, and confusion ( $n = 5$ ). Cardiovascular symptoms like palpitations ( $n = 3$ ) were also noted in a few cases; our patient surprisingly presented with syncope with a low Glasgow Coma Scale score of 10/15 and respiratory failure. Almost all patients were normotensive and were maintaining oxygen saturation at room air. All of them were found to have positive urinary ketones or serum ketonemia in association with high AG metabolic acidosis in the absence of hyperglycemia and a history of diabetes. This was confirmed by arterial blood gases, serum and urinary ketones, and an HbA1c. The  $\beta$ -hB was the most commonly used tested ketone body with a value ranging from 80 till 9000  $\mu\text{mol/L}$

**Table 2. Previously Reported Cases of Nondiabetic Euglycemic Ketoacidosis<sup>3</sup>.**

| No. | Author/Reference                    | Age/Gender | Presentation   | BMI, T, BP, HR, RR, SaO <sub>2</sub>                          | Laboratory Studies  | Diagnosis                                      | Associations/Comorbidities   | Management   | Outcome                            | Follow-up        |
|-----|-------------------------------------|------------|--|---|---|--|--|--|------------------------------------|------------------|
| 1   | Shah and Isley <sup>2</sup>         | 51/Female  | Vomiting   | BMI 21.7  | Anion gap acidosis, ketonuria, normal HbA1c   | HAGMA and urinary ketones                      | Low-carb for 4 years (20 g/day)  | IV fluids and insulin  | Improved and discharged            | NA               |
| 2   | Hudak et al <sup>5</sup>            | 32/Female  | Tachypnea, tachycardia                                     | NA  | Urine acetone +3, Hb 14.5, WBC 10 200, RBS 108, pH 6.8, HCO <sub>3</sub> 3, PCO <sub>2</sub> 8, AG 28, lactate 1      | HAGMA and urinary acetone                      | Fasting and lactation  | ICU care: IV normal saline, sodium bicarbonate (8.4%), potassium, phosphate, and glucose | Improved and discharged on day 10  | NA               |
| 3   | von Geijer and Ekelund <sup>6</sup> | 32/Female  | Nausea/vomiting, palpitations, trembling, extremity spasms | T36; BP 110/80; HR 102; RR 12; SaO <sub>2</sub> 96%           | Ketones 7.1, pH 7.20, pCO <sub>2</sub> 25, RBS 103, lactate 1.0   | HAGMA and ketosis                              | LCHF diet (carb <20 g/day)   | IV glucose infusion, IV insulin  | Improved and discharged on day 3   | 1 month, stable  |
| 4   | Iwata et al <sup>7</sup>            | 76/Female  | Dizziness, headache  | BMI 22.8; T36; BP 106/94; HR 100; RR 15; SaO <sub>2</sub> 99% | β-hb 6663, Hb 15.6, BUN 26.3, WBC 12 500, lactate 27.2, pH 7.28, AG 27.3, RBS 93, HbA1c 6.8%                          | HAGMA and ketosis                              | Hamburger steak only(carb = 12.7 g/day)  | IV normal saline and regular diet  | Improved and discharged on day 3   | 2 months, stable |
| 5   | Lewis <sup>8</sup>                  | 5/Male     | Vomiting/diarrhea, fever                                   | T36.8; BP 110/70; HR 100; RR 20                               | Serum ketone 80, Hb 13.5, HCO <sub>3</sub> 6, RBS 82, AG 27, pH 7.14, pCO <sub>2</sub> 14, lactate 0.7                | HAGMA and serum ketones                        | Diet of aspartame-sweetened 7-LUP  | IV dextrose in water with 0.45% NaCl and potassium acetate                               | Improved and discharged on day 2   | NA               |
| 6   | Freeman et al <sup>9</sup>          | 42/Female  | Nausea/vomiting  | BMI 25.6; T36; BP 114/65; HR 76; RR 18; SaO <sub>2</sub> 100% | WBC 16 900, RBS 99, pH 7.21, HCO <sub>3</sub> 10, lactate 0.9, urine ketone >150                                      | HAGMA and urinary ketones                      | Dukan diet (low-carb, high-protein), fasting                                     | IV 2 L of normal saline, ondansetron, and promethazine                                   | Improved and discharged on day 2   | NA               |
| 7   | Dahl et al <sup>10</sup>            | 24/Male    | Malaise, vomiting/dyspnea                                  | HR 140; RR 60   | pH 7.08, AG 36, Hb 14.3, WBC 19 600, RBS 145, lactate 0.9, urine ketones +4, serum acetoacetate 4.5                   | HAGMA, serum acetoacetate and urinary ketones  | Water and Diet   | IV normal saline and sodium bicarbonate  | Improved and discharged            | None             |
| 8   | Causso et al <sup>11</sup>          | 35/Male    | Aggressiveness, incoherent speech, fasting for over a week | BMI 16  | RBS 57, pH 7.08, HCO <sub>3</sub> 7.4, pCO <sub>2</sub> 25, Hb 13.7, lactate 2.6, acetoacetate 5.1, urine ketone >150 | HAGMA, serum acetoacetate, and urinary ketones | Fruitarian (ate only fruits for 10 years) due to undetermined psychotic disorder | IV normal saline, IV dextrose, bicarbonate   | Improved and discharged            | None             |
| 9   | Monnier et al <sup>12</sup>         | 29/Female  | Dyspnea/anorexia, fatigue, weight loss                     | NA  | pH 7.11, Hb 11, WBC 10 000, PaCO <sub>2</sub> 8, PaO <sub>2</sub> 136 mm Hg, HCO <sub>3</sub> 2.9                     | HAGMA and ketosis                              | Restricted diet, lactation, and gastric bariatric surgery                        | IV normal saline   | Improved and discharged in 7 hours | NA               |
| 10  | Wuopio et al <sup>13</sup>          | 21/Female  | Nausea/dyspnea, headache                                   | NA  | pH 6.9, RBS 120, B-acetones 10, AG 21   | HAGMA and serum acetone                        | Low-carbohydrate diet and lactation  | IV Ringer lactate and IV glucose   | Improved and discharged on day 1   | NA               |
| 11  | Milroy and Parai <sup>14</sup>      | 3/Female   | Death  | NA  | β-hb 3966, RBS 89, HbA1c 4.7%   | HAGMA and ketosis                              | Starvation, pneumonia, and Armani-Ebstein lesion                                 | NA   | NA                                 | NA               |
| 12  | Wei et al <sup>15</sup>             | 36/Female  | Dyspnea/vomiting/anorexia                                  | BMI 26.5; T37; BP 126/78; HR 138; RR 25                       | β-hb 7216, WBC 15 850, RBS 152, HbA1c 4.8%, pH 7.21, HCO <sub>3</sub> 11, PCO <sub>2</sub> 28, AG 15, lactate normal  | HAGMA and ketosis                              | Starvation, pregnancy, asthma  | ICU care: intubation, albuterol, dexamethasone, dextrose in normal saline and C-section  | Improved and discharged on day 14  | NA               |
| 13  | Scholte and Boer <sup>16</sup>      | 26/Female  | Dyspnea/vomiting and abdominal discomfort                  | BMI 44.3; T37; BP 150/90                                      | Urine ketone 7.8, HbA1c 5.4%, pH 7.30, HCO <sub>3</sub> 9, AG 21, RBS 120   | HAGMA and urinary ketones                      | Starvation, pregnancy, hemochromatosis   | ICU care: IV antiemetics and normal saline, 8.4% sodium bicarbonate                      | Improved and discharged on day 5   | NA               |

(continued)



Table 2. (continued)

| No. | Author/Reference                    | Age/Gender             | Presentation                                    | BMI, T, BP, HR, RR, SaO <sub>2</sub>                           | Laboratory Studies   | Diagnosis   | Associations/Comorbidities                   | Management  | Outcome                           | Follow-up               |
|-----|-------------------------------------|------------------------|---|--|--|---|--|---|-----------------------------------|-------------------------|
| 14  | Jain et al <sup>17</sup>            | 50/Female              | Comatose state, vomiting/anorexia               | T37; BP 110/80; HR 120; RR 34; SaO <sub>2</sub> 93%            | β-Hb 14.860, RBS <20 mg/dL, HCO <sub>3</sub> <sup>-</sup> 9, AG 9, pH 7.2, pCO <sub>2</sub> 24, lactate 2.3, ketonuria           | HAGMA urine and serum ketones                           | Chronic alcoholism and chronic liver disease | ICU incubate, IV normal saline and 5% dextrose with thiamine and folate | Vegetative state                  | 10 months               |
| 15  | Gill and Yong <sup>18</sup>         | 39/Male                | Abdominal pain and vomiting                     | HR 120   | Ketonuria, WBC 11 500, pH 7.18, pCO <sub>2</sub> 20, pO <sub>2</sub> 94, HCO <sub>3</sub> <sup>-</sup> 10                        | HAGMA and urine ketones                                 | Chronic alcoholism and pancreatitis          | IV 0.9% NaCl infusion and glucose, potassium, insulin infusion          | Improved and discharged on day 3  | 6 months, third episode |
| 16  | Devenyi <sup>19</sup>               | 71/Female              | Vomiting/fasting                                | NA   | Raised serum β-Hb and ketonuria, RBS 31, pH 6.9  | HAGMA urine and serum ketones                           | Chronic alcoholism                           | IV hypertonic dextrose  | Improved twice                    | Died, third episode     |
| 17  | Bernau et al <sup>20</sup>          | N/A                    | Vomiting/starvation                             | NA   | Raised serum β-Hb and ketonuria, AG 25 to 41, lactate 0.9  | HAGMA urine and serum ketones                           | Chronic alcoholism                           | IV fluids   | NA                                | NA                      |
| 18  | Platia and Hsu <sup>21</sup>        | 5 patients, 28-39/Male | Vomiting/stuporous, comatose                    | NA   | Raised β-Hb up to 9800, low glucose level  | HAGMA urine and serum ketones                           | Chronic alcoholism                           | IV thiamine, glucose, and dextrose                                      | Improved and discharged on day 2  | 1 and 3 months, stable  |
| 19  | Arena et al <sup>22</sup>           | 78/Female              | Confusion and lethargy                          | T 37.4; BP 140/72; HR 84; RR 28                                | Raised serum ketones, salicylate 60.8 mg/dL, RBS 38, pH 7.28   | HAGMA and serum ketones                                 | Salicylates for degenerative joint disease   | IV dextrose and sodium bicarbonate                                      | Improved and discharged           | NA                      |
| 20  | Wood and Kinlaw <sup>23</sup>       | 41/Female              | Vomiting/diarrhea, palpitation, and weight loss | T 27.3; BP 146/75; HR 112                                      | β-Hb 52.10, pH 7.36, pCO <sub>2</sub> 26, HCO <sub>3</sub> <sup>-</sup> 14.5, RBS normal   | HAGMA and serum ketones                                 | Hyperthyroidism                              | IV normal saline, propranolol, propylthiouracil, thiamine               | Improved and discharged           | NA                      |
| 21  | Vestergaard et al <sup>24</sup>     | 79/Female              | Stroke and weight loss ~13 kg                   | NA   | NA   | Metabolic acidosis with elevated poly-3-hydroxybutyrate | Pneumonia and dehydration                    | IV normal saline, potassium, bicarbonate, and insulin                   | Improved and discharged on day 14 | NA                      |
| 22  | Valkenburgh and Braal <sup>25</sup> | 19/Female              | Metabolic acidosis after bariatric surgery      | BMI 40.5; T 36; BP 106/94; HR 100; RR 16; SaO <sub>2</sub> 92% | pH 7.28, pCO <sub>2</sub> 34, HCO <sub>3</sub> <sup>-</sup> 16   | Starvation-induced ketoacidosis                         | Bariatric surgery                            | Oral glucose therapy  | Improved and discharged on day 3  | NA                      |
| 23  | The present report                  | 33/Female              | Syncopal, respiratory failure                   | BMI 27; T 37; BP 140/71; HR 136; RR 22; SaO <sub>2</sub> 96%   | β-Hb 10.600, pH 6.6, pCO <sub>2</sub> 16.9, pO <sub>2</sub> 120.5, HCO <sub>3</sub> <sup>-</sup> 2.6, AG 30, RBS 126, HbA1c 5.1% | HAGMA and serum ketones                                 | Atkins diet and flu                          | ICU: intubation, hemodialysis and IV dextrose saline and insulin        | Improved and discharged on day 7  | 3 months, stable        |

Abbreviations. BMI, body mass index; HbA1c, hemoglobin A1c; HAGMA, high anion gap metabolic acidosis; IV, intravenous; NA, not available; WBC, white blood cells; RBS, random blood sugar; HCO<sub>3</sub><sup>-</sup>, bicarbonate; pCO<sub>2</sub>, partial pressure of carbon dioxide; AG, anion gap; ICU, intensive care unit; BP, blood pressure; HR, heart rate; RR, respiratory rate; SaO<sub>2</sub>, oxygen saturation; LCHF, low-carb high fat; β-Hb, β-hydroxybutyrate; Hb, hemoglobin; BUN, blood urea nitrogen; NaCl, sodium chloride; pO<sub>2</sub>, partial pressure of oxygen.

<sup>a</sup>Units of measurement: BMI in kg/m<sup>2</sup>; Temperature (T) in Celsius; BP in mm Hg; HR is in beats per minute; RR is in breaths per minute; SaO<sub>2</sub> is in percentage; β-Hb is in μmol/L; acetoacetate is in mmol/L; Hb, RBS, lactate, and BUN are in mg/dL; WBC is in counts/μL; AG is in mEq/L; HCO<sub>3</sub><sup>-</sup> is in mmol/L; pCO<sub>2</sub> is in mm Hg; and β-acetones in 10 mmol/L.

in the previously reported cases; in our case, the value was above 10 000  $\mu\text{mol/L}$ . High AG metabolic acidosis in reported cases was spread between a pH of 6.8 to 7.2,  $\text{HCO}_3^-$  2.9 to 10 mEq/L, and AG 21 to 28. In our patient, however, the pH was very low 6.6,  $\text{HCO}_3^-$  of 2.6, and AG 30. These findings were alarming for us and are important for readers suspecting that Atkins diet is associated with serious NDKA in terms of its presentation and laboratory findings. It also signifies that a ketogenic diet can cause severe symptoms and complications like syncope and respiratory failure even in a previously healthy young patient when they encounter a stressful condition like seasonal flu. The characteristics of previously reported cases are summarized in Table 2.

NKDA is usually managed with IV dextrose along with insulin, IV bicarbonate, correction of electrolyte imbalances, and supportive treatment until the acidosis is resolved. Though NDKA is a serious and potentially life-threatening condition, timely diagnosis and proper management as in our case leads to good outcomes. Outcomes were not reported in all the cases; however, in cases where details were available, all the patients recovered with no significant long-term sequelae.

## Conclusions

- Starvation, pregnancy, lactation, and ketogenic (Atkins) diet-associated ketoacidosis can be severe and possibly fatal if not treated promptly.
- Physicians should consider diet-associated ketoacidosis in previously nondiabetic patients presenting with severe metabolic acidosis.
- Hemodialysis is only helpful as a temporizing measure but has no role as definitive management.
- Intravenous insulin is the mainstay of treatment even in patients with normal glucose levels; however, intravenous dextrose should be given to avoid hypoglycemia.
- To summarize, Atkins diet and other previously reported diets like Dukan, fruitarian, aspartame, and other low-carb high-fat diets should be considered as an extra “D” in the mnemonic “MUDPILES” for causes of high anion gap metabolic acidosis.

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.



## Ethics Approval

Our institution does not require ethical approval for reporting individual cases or case series.

## Informed Consent

Verbal informed consent was obtained from the patient(s) for their anonymized information to be published in this article.

## ORCID iDs

Waqas Ullah  <https://orcid.org/0000-0002-4850-0309>  
 Mohsin Hamid  <https://orcid.org/0000-0002-7105-4802>  
 Faisal Inayat  <https://orcid.org/0000-0001-7576-7319>

## References

1. O'Meara M. Pediatric emergency medicine secrets. *Emerg Med.* 2002;14:113-114.
2. Shah P, Isley WL. Ketoacidosis during a low-carbohydrate diet. *N Engl J Med.* 2006;354:97-98.
3. Gilman SL. *Diets and Dieting: A Cultural Encyclopedia.* New York, NY: Routledge; 2008.
4. Jenkins DJ, Wong JM, Kendall CW, et al. Effect of a 6-month vegan low-carbohydrate (“Eco-Atkins”) diet on cardiovascular risk factors and body weight in hyperlipidaemic adults: a randomised controlled trial. *BMJ Open.* 2014;4:e003505.
5. Hudak SK, Overkamp D, Wagner R, Häring HU, Heni M. Ketoacidosis in a non-diabetic woman who was fasting during lactation. *Nutr J.* 2015;14:117.
6. von Geijer L, Ekelund M. Ketoacidosis associated with low-carbohydrate diet in a non-diabetic lactating woman: a case report. *J Med Case Rep.* 2015;9:224.
7. Iwata H, Tsuzuki S, Iwata M, Terasawa T. Ketoacidosis due to a low-carbohydrate diet in an elderly woman with dementia and abnormal eating behavior. *Intern Med.* 2017;56:2671-2675.
8. Lewis P. Starvation ketosis after rehydration with diet soda. *Pediatrics.* 1991;88:806-807.
9. Freeman TF, Willis B, Krywko DM. Acute intractable vomiting and severe ketoacidosis secondary to the Dukan Diet®. *J Emerg Med.* 2014;47:e109-e112.
10. Dahl GT, Woldseth B, Lindemann R. Metabolic acidosis mimicking diabetic ketoacidosis after use of calorie-free mineral water. *Eur J Pediatr.* 2012;171:1405-1407.
11. Causso C, Arrieta F, Hernández J, et al. Severe ketoacidosis secondary to starvation in a frutarian patient. *Nutr Hosp.* 2010;25:1049-1052.
12. Monnier D, Goulenok T, Allary J, Zarrouk V, Fantin B. Starvation ketosis in a breastfeeding woman [in French]. *Rev Med Interne.* 2015;36:854-858.
13. Wuopio J, Schiborr R, Charalampakis G. Severe ketoacidosis in breastfeeding woman with low energy and carbohydrate intake [in Swedish]. *Lakartidningen.* 2015;112. <http://lakartidningen.se/Klinik-och-vetenskap/Fallbeskrivning/2015/09/Svar-ketoacidosis-hos-ammande-kvinna-med-lagt-energi-och-kolhydratintag/>. Accessed August 10, 2018.
14. Milroy CM, Parai JL. Armani-Ebstein lesion, ketoacidosis and starvation in a child. *Forensic Sci Med Pathol.* 2011;7:213-216.
15. Wei KY, Chang SY, Wang SH, Su HY, Tsai CL. Short-term starvation with a near-fatal asthma attack induced ketoacidosis in a nondiabetic pregnant woman: a case report. *Medicine (Baltimore).* 2016;95:e4042.
16. Scholte JB, Boer WE. A case of nondiabetic ketoacidosis in third term twin pregnancy. *J Clin Endocrinol Metab.* 2012;97:3021-3024.



17. Jain H, Beriwal S, Singh S. Alcohol induced ketoacidosis, severe hypoglycemia and irreversible encephalopathy. *Med Sci Monit.* 2002;8:CS77-CS79.
18. Gill G, Yong A. Recurrent alcoholic ketoacidosis with hyperglycaemia in a non-diabetic patient. *Hosp Med.* 2000;61:506-507.
19. Devenyi P. Alcoholic hypoglycemia and alcoholic ketoacidosis: sequential events of the same process? *Can Med Assoc J.* 1982;127:513.
20. Bernuau J, Nouel O, Degos F, Rueff B, Benhamou JP. Alcoholic ketoacidosis (author's translation) [in French]. *Nouv Presse Med.* 1979;8:3625-3627.
21. Platia EV, Hsu TH. Hypoglycemic coma with ketoacidosis in nondiabetic alcoholics. *West J Med.* 1979;131:270-276.
22. Arena FP, Dugowson C, Saudek CD. Salicylate-induced hypoglycemia and ketoacidosis in a nondiabetic adult. *Arch Intern Med.* 1978;138:1153-1154.
23. Wood ET, Kinlaw WB. Nondiabetic ketoacidosis caused by severe hyperthyroidism. *Thyroid.* 2004;14:628-630.
24. Vestergaard T, Aaen JF, Bruun JM. Starvation is a rare cause of metabolic ketoacidosis [in Danish]. *Ugeskr Laeger.* 2014;176:V05140275.
25. Valkenborgh T, Bral P. Starvation-induced ketoacidosis in bariatric surgery: a case report. *Acta Anaesthesiol Belg.* 2013;64:115-117.