6-2012

Revision Total Knee Arthroplasty: Infection should be Ruled Out in All Cases

Mohammad R. Rasouli

Rothman Institute of Orthopaedics at Thomas Jefferson University Hospital

Armin Aalami Harandi

Rothman Institute of Orthopaedics at Thomas Jefferson University Hospital

Bahar Adeli

Rothman Institute of Orthopaedics at Thomas Jefferson University Hospital

James J. Purtill

Rothman Institute of Orthopaedics at Thomas Jefferson University Hospital

Javad Parvizi

*Rothman Institute of Orthopaedics at Thomas Jefferson University Hospital*Follow this and additional works at: https://jdc.jefferson.edu/rothman_institute Part of the [Orthopedics Commons](#)[Let us know how access to this document benefits you](#)

Recommended Citation

Rasouli, Mohammad R.; Harandi, Armin Aalami; Adeli, Bahar; Purtill, James J.; and Parvizi, Javad, "Revision Total Knee Arthroplasty: Infection should be Ruled Out in All Cases" (2012).

Rothman Institute Faculty Papers. Paper 7.https://jdc.jefferson.edu/rothman_institute/7

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Rothman Institute Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Revision Total Knee Arthroplasty: Infection should be Ruled Out in All Cases

Abstract:

We hypothesized that some aseptic revision knee arthroplasty (TKA) failures are indeed due to occult infection. This prospective study recruited 65 patients undergoing revision TKA. Mean follow-up period was 19 months. Collected synovial fluid was analyzed by Ibis T5000 biosensor (a multiplex PCR technology). Cases were considered as infected or aseptic based on the surgeon's judgment and Ibis findings. Based on Ibis biosensor, 17 aseptic cases were indeed infected that had been missed. Of these 17 cases, 2 developed infection following the index revision. A considerable number of so called aseptic failures seem to be occult infections who were not adequately investigated and/or miss-categorized as aseptic failure. We recommend that all patients undergoing revision arthroplasty need to be investigated for PJI.

Key words: Revision knee arthroplasty, Aseptic failure, Periprosthetic joint infection, Ibis T5000 biosensor, Diagnosis

18 **Introduction**

19 Periprosthetic joint infection (PJI), that occurs following 1 to 3% of TKAs [1, 2] is the
20 most common cause of failure after total knee arthroplasty (TKA) [3-5]. Diagnosis of PJI
21 continues to pose a challenge to the medical community because of lack of a “gold standard” [6].
22 It is, however, critical that aseptic cases be distinguished from PJI, as treatment for these
23 conditions is vastly different [7, 8].

24 History taking, physical examination, and radiographic findings can be similar in PJI and
25 aseptic loosening and may not allow distinction in most cases [7]. Joint aspiration and serologic
26 tests such as erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP) are usually
27 performed during the work-up of a patient with suspected PJI. Although a very high sensitivity
28 has been reported for ESR and CRP, these laboratory tests have their own limitations in the
29 diagnosis of PJI [9]. Even intraoperative culture and pathology have limited sensitivity in
30 diagnosis of PJI particularly in those receiving antibiotics [10, 11]. It has been demonstrated that
31 intraoperative culture yields false negative results in 6.6% [12] to 17.7% [13] and false positive
32 in 13% of cases [14]. Imaging modalities such as nuclear scans (tc-99, indium 111, gallium) have
33 also been employed in diagnosis of PJI but these tests are also unable to accurately differentiate
34 between aseptic loosening and septic failure after joint arthroplasty [15-17].

35 One of the main reasons for inability to isolate the pathogen relates to the presence of a
36 biofilm [8, 18] and/or internalization of organism by osteoblasts [19]. Biofilm is a complex
37 structure comprising microorganisms enveloped in macromolecules of glycocalyx and other
38 protective films [18, 20]. As a result, it is probable that some of PJIs to be missed and treated as
39 aseptic failure which subsequently cause recurrent failure [21].

Using molecular techniques may improve diagnosis of PJI as these methods have high sensitivity and are culture independent [11]. Polymerase chain reaction (PCR) has been used in several studies to diagnose PJI [22-25]. Using a specific PCR or a broad-range (16S ribosomal DNA) PCR which are respectively able to detect only a single microorganism or previously unknown organisms were limitations of these studies. Compared to the specific PCR, the sensitivity and specificity of the broad-range PCR is lower, needs subsequent sequencing for bacterial identification, and fails to detect mixed infections [11]. Recently, the Ibis T5000 universal biosensor has been introduced as a sensitive and specific method for identification of bacteria, viruses, fungi, and protozoa. The system operates based on broad-range PCR and high-performance mass spectrometry and seems to be more accurate than conventional PCR. [26] However, it has not yet been approved by the Food and Drug Administration (FDA) for routine use in clinical practice.

The hypothesis of this study was that some cases of aseptic failure were indeed due to infection that had either escaped diagnosis using conventional modalities or had not been adequately investigated. This prospective study was set up to examine the postoperative course of a consecutive cohort of patients undergoing revision TKA in whom an intra-articular tissue and/or fluid sample was also sent for analysis by Ibis T5000 biosensor.

Materials and Methods

After approval of the Institutional Review Board of the Thomas Jefferson University, all patients who underwent TKA revision from February 2009 to May 2010 were recruited for this study. The study consists of 65 patients of whom 33 were men. The mean age of the patients was

65 $\pm$ 11 years. All patients underwent appropriate preoperative work-up based on the recommendation of the treating surgeon and then categorized as infected or uninfected based on these investigations and surgeon's judgment. In our center, patients who are suspicious for PJI are evaluated by measurement of serum erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP), intraoperative culture and synovial fluid analysis. However, frozen section isn't use as a part of PJI work-up in our institute. Intraoperatively tissue and/or fluid samples were collected and analyzed using the Ibis T5000 biosensor. After discharging from the hospital, patients are followed-up based on the protocol which is used routinely in our institution at 6 weeks, 6 months and 2 years after the revision surgery. In this study, all patients were followed-up for a minimum of six months with a mean follow-up of 19 months (range; 12 to 26). In particular, "aseptic" patients in whom the Ibis T5000 biosensor had detected an infecting pathogen were followed-up closely for development of subsequent failure after the index revision.

Preoperative work-up

Detailed history taking, examination, and routine radiographs were performed in all patients in this cohort. It is institutional policy that all patients undergoing revision arthroplasty at our institution have preoperative Erythrocyte Sedimentation Rate (ESR) and C-Reactive Protein (CRP) measured. In addition, and based on the findings of serology, majority of patients (n=57) underwent joint aspiration with the fluid sent for analysis for neutrophil count, neutrophil differential, and culture. As this was an observational and not interventional study, we did not make any changes to the preoperative work-up of any patients. Collection of synovial fluid and/or tissue sample for analysis by Ibis T5000 biosensor was a requirement of the study.

84 *Sample collection and the Ibis T5000 method*

85 Joint fluid and/or tissue were collected intraoperatively. Joint fluid was aspirated prior to
86 the arthrotomy and was also sent for WBC count (if indicated) and culture. Tissue-sampling was
87 performed from areas that were considered by the surgeon that to be most suggestive of
88 infection. In addition to the Ibis analysis, tissues were sent for histopathologic assessment and
89 culture. The samples were not sent for measurement of inflammatory markers for this study.
90 Samples then processed appropriately for later analysis by Ibis T5000 in batches. Appropriate
91 cryogenic vials were used to store fluid samples in a Styrofoam container. The vials were
92 transferred in ice bags from the operating room to liquid nitrogen. About 0.5 to 1 mL of the
93 liquid was stored for further analysis in liquid nitrogen. The syringe was changed in order to
94 minimize possibility of accidental microbial contaminations before synovial fluid was transferred
95 to the labeled vials. These vials were transferred to -140o F freezer where they were stored until
96 they were shipped in batch to the Center for Research and Genomic Studies in Allegheny, PA.

97 For DNA extraction, 1 ml of the aspirate was centrifuged at 10,000 rpms for 3 min and
98 900uL of supernatant was removed. Then, ATL lysis buffer and proteinase K were added and the
99 samples were incubated at 56°C until lysis occurred. The Qiagen DNeasy Tissue kit (Qiagen. Inc.
100 cat # 69506) was used to extract nucleic acid from the lysed sample. After DNA was extracted,
101 10 uL of sample was loaded per well into each of 16 wells on the BAC detection PCR plate that
102 each contained a different primer pair (Abbott Molecular. cat # PN 05N13-01). The BAC
103 detection plate is a 96 well, 6 sample plate which contains 16 primers that identify all bacterial
104 organisms, Candida species, and determines the presence of several key antibiotic resistance
105 markers such as van-A and van-B (vancomycin resistance) in Enterococcus species, KPC
106 (carbapenem resistance) in Gram-negative bacteria, and mec-A (methicillin resistance) in

Staphylococcus species. Once PCR was completed, the plate was loaded onto the Ibis T5000 machine. The products from the PCRs were desalted in a 96-well plate format and sequentially electrosprayed into the time-of-flight mass spectrometer. The resultant spectral signals were then processed to determine the masses of each of the PCR products present with sufficient accuracy that the base composition of each amplicon could be unambiguously deduced. Using combined base compositions from multiple PCRs, the identities of the pathogens and their relative concentrations in the starting sample were determined.

The isolated microorganism from the Ibis biosensor was considered as an “orthopedic pathogen” based on extensive search of the available literature. In other words, if there was any evidence even a case report that shows the isolated microorganism is able to cause bone and/or joint infection, that microorganism was defined as an “orthopedic pathogen”. However, Ibis biosensor results did not change the treatment strategy and all patients were treated based on results obtained from conventional diagnostic tests and surgeon’s judgment.

Results

Based on preoperative investigations and surgeon’s judgment, of the 65 patients recruited for this study, 21 patients were undergoing revision arthroplasty for PJI and the remaining 44 patients had aseptic failure. Among the 21 patients with PJI, synovial culture was negative in 11 cases. In the remaining 10 patients the isolated organisms were coagulase negative *Staphylococcus* (5 cases), *Staphylococcus aureus* (3 patients), *Streptococcus mitis* plus *Streptococcus sanguis* (1 case) and *Peptostreptococcus* species (1 case). Ibis identified a pathogen with confidence ≥ 0.7 in total of 36 cases. Ibis T5000 isolated an organism in 19 PJI cases and failed to isolate any organism in 2 cases that were categorized as infected. The isolated organism by Ibis was coagulase negative *Staphylococcus* in 10 patients and *Staphylococcus*

aureus in 4 patients. In the PJI group, comparison of the isolated organisms from the culture and the detected organism by the Ibis T5000 biosensor showed the same pathogens in 9 cases samples whereas in 11 cases, the Ibis biosensor found additional pathogen. Table 1 demonstrates comparison between isolated organism by culture and Ibis T5000 results in patients with PJI. On the other hand, the Ibis T5000 found additional non-pathogen organisms in 3 cases in which the Ibis T5000 had also detected an orthopedic pathogen.

All cultures in the aseptic group were negative whereas in 17 cases the Ibis T5000 found orthopedic pathogens. In 27 patients, the biosensor failed to find any orthopedic pathogens. Table 2 demonstrates more details on aseptic cases. During the follow-up period, 2 patients failed and needed re-revision who were both initially revised for aseptic failure. The cause of failure in these 2 patients was subsequent PJI with the same organism (Coagulase negative *Staphylococcus*) as one isolated by the Ibis T5000. At the latest follow-up which ranged from 12 to 26 months after the index revision, all the remaining patients appear to be doing well with no evidence of infection.

Discussion

Given the completely different management of aseptic loosening and PJI as well as the importance of early diagnosis of PJI for establishment of a more effective treatment, distinguishing between these two conditions needs special attention. Absence of a “gold standard” for diagnosis of PJI [27] in addition to various defensive mechanisms of pathogens such as biofilm production [8] make this differentiation more difficult. The infecting organism that segregate in biofilm evade detection by conventional culture as the latter relies on isolation of planktonic organisms. As a result it is suggested that some cases of PJI escape detection and

erroneously are categorized as aseptic failures [8]. Although many factors contribute to development of PJI after revision surgery, the latter point may be considered as one of the contributing factor for the much higher incidence of PJI after revision arthroplasty than that after primary replacement.

The Ibis T5000 universal biosensor is a promising technology that has been used to identify a wide spectrum of pathogens in sepsis [28] and it may cover limitations of PCR method for diagnosis of PJI. Because of reliance on mass spectrometry and further “purification” of DNA it is assumed that Ibis does not suffer the same extreme sensitivity as conventional PCR. Further, because of pan-genomic amplification, Ibis may be able to detect infecting organisms that could be missed by conventional PCR.

This prospective study was designed to examine the possibility of escaping some cases of TKA failures which are assumed to be “aseptic” from conventional diagnostic tests. These cases may be indeed infections, that have escaped diagnosis and have been miss-categorized as “aseptic” failures. The study relied on Ibis T5000 for isolation of organism. Although we did not accept Ibis as the “gold standard” for diagnosis of PJI, we were interested to know in what percentage of patients with aseptic failure Ibis T5000 biosensor was able to isolate a pathogen. Further, we sought to examine the correlation between conventional culture and Ibis in terms of their ability for isolation of a pathogen and its resistance profile.

At our institution since 2006, we have utilized an algorithmic approach for work-up of patients with failed arthroplasty which includes determination of ESR and CRP prior to revision arthroplasty and selective aspiration of the failed joint in those with abnormal serology or high index of suspicion for PJI [7]. In addition, intraoperative culture is performed for all cases

undergoing revision arthroplasty. In spite of employing such a comprehensive and strict approach, the present study revealed that a few PJIs cannot be detectable by using routine diagnostic tools. It appears that reliance on conventional investigations is likely to miss occult PJI at least in 30% of patients (13 out of 44 if *Enterococcus faecalis* cases were considered as contamination). The 2 patients who were originally assumed to have aseptic failure, developed infection shortly after the index revision by organisms that had been isolated by the Ibis T5000 but failed to be detected by conventional culture. The infecting organism in one case was low-virulence, but a recognized pathogen [29, 30] namely *Staphylococcus Caprae*. Although these patients did not receive any treatment for isolated additional pathogens from the Ibis biosensor, our findings may indirectly indicate the clinical importance of isolated pathogens from the Ibis biosensor. However, we are not able to make a statement about effect of treatment on outcome of these patients with negative culture in whom the Ibis biosensor isolates additional pathogens. It is possible that the conventional culture may have identified these organisms if supplemented culture was utilized or the culture was kept for an extended period of time.

PCR has been used previously for the purpose of isolating organism in cases of suspected PJI [22, 23, 31]. However, PCR methods suffer several limitations, most important of which is the high incidence of false positive results [11]. The technique is so sensitive that it may amplify contaminating and non-infecting organism such as those residing on the skin that may have been picked up by aspirating needle [25]. Ibis T5000 is a multiplex PCR technology that was designed to overcome some of the limitations of conventional PCR.

Compared to conventional PCR, the Ibis T5000 utilizes a pan-genomic amplifier that is capable of isolating atypical bacteria and even non-bacterial pathogens such as fungi. Unlike conventional PCR, Ibis does not rely on universal primers for amplification of DNA which may

detect contaminating organism. Instead Ibis T5000 biosensor uses multiple pairs of species-specific primers to amplify regions of an organism's genome. This process is followed by the identification of that region's base composition using mass spectrometry, the results of which are compared to a database which matches it to the closest microorganism [26]. The Ibis T5000 universal biosensor technology combines nucleic acid amplification to high-performance electrospray ionization mass spectrometry and base-composition analysis. The system can identify and quantify all known bacteria, all major groups of pathogenic fungi, and the major families of pathogen viruses in humans and animals. Moreover, the system is capable of detecting virulence factors and antibiotic resistance markers [26].

Despite its appeal, Ibis T5000 may still be a victim of high sensitivity. Ibis isolated a "pathogen" in 17 cases (38%) of "aseptic" cases. Of the latter 2 patients have failed so far due to infection which we believe was missed during the index revision arthroplasty. It is possible that occult PJI may have been present in a few more cases that were either eradicated during index revision arthroplasty, effectively with the patient undergoing a one stage exchange arthroplasty, or are likely to manifest a failure with further follow-up. It is unlikely that the isolated organism by Ibis in all 17 aseptic cases represent a true pathogen. Thus, this technology, despite its appeal, should be reserved for patients in whom high index of suspicion or PJI exists but no organism can be isolated. In other words the indication for use of Ibis, in our opinion, is for cases of culture negative PJI.

The study suffers a few limitations. Perhaps the most important limitation of this study is the relatively short follow-up. As mentioned above, it is possible that with longer surveillance we are likely to encounter more patients who may fail. Although plausible, the latter is unlikely to alter the message of this study. The study highlights the importance of routine preoperative

work-up using conventional serology for all and selected aspiration for some, in line with the recent American Academy of Orthopedic Surgeons guidelines for diagnosis of PJI [6, 32]. It also highlights the fact that a sophisticated technology is available for use by orthopedic surgeons to isolate the infecting organism in cases of culture negative PJI. Another limitation of the study relates to lack of a “standard” definition for PJI. It is possible that using a different diagnostic criteria, some of the PJI cases in our cohort may have been considered as uninfected and vice versa. The latter is unfortunately a limitation inherent to any studies related to topic of PJI as various definitions for PJI exist and depending on the definition used the percent of infected versus uninfected cases in a given cohort may change. It is hoped that orthopedic societies in collaboration with other organizations may be able to address this shortfall in the future.

Despite the aforementioned limitations, the present study demonstrated that some aseptic loosening are not “truly” aseptic and are low grade PJIs that remain undiagnosed using conventional modalities. Some of these cases may fail early for a subsequent infection. This may explain the relatively high incidence of infection following revision arthroplasty, compared to primary, and also the high rate of early failure of revision cases. It is thus recommended that all patients undergoing revision for failed arthroplasty should be subjected to routine conventional work-up which includes routine serology (ESR and CRP) and joint aspiration in patients with abnormal serology tests and high index of suspicion for PJI. With further refinements of molecular techniques such as multiplex PCR, the true nature of some of these so called “aseptic” failures is likely to be revealed.

243 **References**

- 244 1. Della Valle CJ, Zuckerman JD, Di Cesare PE. Periprosthetic sepsis. Clin Orthop Relat Res
245 2004; 420: 26.
- 246 2. Phillips JE, Crane TP, Noy M, et al. The incidence of deep prosthetic infections in a specialist
247 orthopaedic hospital: a 15-year prospective survey. J Bone Joint Surg Br 2006; 88: 943.
- 248 3. Sharkey PF, Hozack WJ, Rothman RH, et al. Insall Award paper. Why are total knee
249 arthroplasties failing today? Clin Orthop Relat Res 2002; 7.
- 250 4. Bozic KJ, Kurtz SM, Lau E, et al. The epidemiology of revision total knee arthroplasty in the
251 United States. Clin Orthop Relat Res 2010; 468: 45.
- 252 5. Mortazavi SM, Molligan J, Austin MS, et al. Failure following revision total knee
253 arthroplasty: infection is the major cause. Int Orthop 2010 Oct 21. [Epub ahead of print];
- 254 6. Parvizi J, Della Valle CJ. AAOS Clinical Practice Guideline: diagnosis and treatment of
255 periprosthetic joint infections of the hip and knee. J Am Acad Orthop Surg 2010; 18: 771.
- 256 7. Parvizi J, Ghanem E, Menashe S, et al. Periprosthetic infection: what are the diagnostic
257 challenges? J Bone Joint Surg Am 2006; 88 Suppl 4: 138.
- 258 8. Nelson CL, McLaren AC, McLaren SG, et al. Is aseptic loosening truly aseptic? Clin Orthop
259 Relat Res 2005; 25.
- 260 9. Greidanus NV, Masri BA, Garbuz DS, et al. Use of erythrocyte sedimentation rate and C-
261 reactive protein level to diagnose infection before revision total knee arthroplasty. A prospective
262 evaluation. J Bone Joint Surg Am 2007; 89: 1409.
- 263 10. Bare J, MacDonald SJ, Bourne RB. Preoperative evaluations in revision total knee
264 arthroplasty. Clin Orthop Relat Res 2006; 446: 40.

265 11. Achermann Y, Vogt M, Leunig M, et al. Improved diagnosis of periprosthetic joint infection
266 by multiplex PCR of sonication fluid from removed implants. *J Clin Microbiol* 2010; 48: 1208.

267 12. Berbari EF, Marculescu C, Sia I, et al. Culture-negative prosthetic joint infection. *Clin Infect*
268 *Dis* 2007; 45: 1113.

269 13. Trampuz A, Piper KE, Jacobson MJ, et al. Sonication of removed hip and knee prostheses for
270 diagnosis of infection. *N Engl J Med* 2007; 357: 654.

271 14. Barrack RL, Harris WH. The value of aspiration of the hip joint before revision total hip
272 arthroplasty. *J Bone Joint Surg Am* 1993; 75: 66.

273 15. Reing CM, Richin PF, Kenmore PI. Differential bone-scanning in the evaluation of a painful
274 total joint replacement. *J Bone Joint Surg Am* 1979; 61: 933.

275 16. Weiss PE, Mall JC, Hoffer PB, et al. ^{99m}Tc-methylene diphosphonate bone imaging in the
276 evaluation of total hip prostheses. *Radiology* 1979; 133: 727.

277 17. Gelman MI, Coleman RE, Stevens PM, et al. Radiography, radionuclide imaging, and
278 arthrography in the evaluation of total hip and knee replacement. *Radiology* 1978; 128: 677.

279 18. Costerton JW, Montanaro L, Arciola CR. Biofilm in implant infections: its production and
280 regulation. *Int J Artif Organs* 2005; 28: 1062.

281 19. Jevon M, Guo C, Ma B, et al. Mechanisms of internalization of *Staphylococcus aureus* by
282 cultured human osteoblasts. *Infect Immun* 1999; 67: 2677.

283 20. Gristina AG, Costerton JW. Bacterial adherence and the glycocalyx and their role in
284 musculoskeletal infection. *Orthop Clin North Am* 1984; 15: 517.

285 21. Parvizi J, Suh DH, Jafari SM, et al. Aseptic Loosening of Total Hip Arthroplasty: Infection
286 Always Should be Ruled Out. *Clin Orthop Relat Res* 2011; 469: 1401.

287 22. Tunney MM, Patrick S, Curran MD, et al. Detection of prosthetic hip infection at revision
288 arthroplasty by immunofluorescence microscopy and PCR amplification of the bacterial 16S
289 rRNA gene. J Clin Microbiol 1999; 37: 3281.

290 23. Mariani BD, Martin DS, Levine MJ, et al. The Coventry Award. Polymerase chain reaction
291 detection of bacterial infection in total knee arthroplasty. Clin Orthop Relat Res 1996; 11.

292 24. Levine MJ, Mariani BA, Tuan RS, et al. Molecular genetic diagnosis of infected total joint
293 arthroplasty. J Arthroplasty 1995; 10: 93.

294 25. Panousis K, Grigoris P, Butcher I, et al. Poor predictive value of broad-range PCR for the
295 detection of arthroplasty infection in 92 cases. Acta Orthop 2005; 76: 341.

296 26. Ecker DJ, Sampath R, Massire C, et al. Ibis T5000: a universal biosensor approach for
297 microbiology. Nat Rev Microbiol 2008; 6: 553.

298 27. Della Valle CJ, Bogner E, Desai P, et al. Analysis of frozen sections of intraoperative
299 specimens obtained at the time of reoperation after hip or knee resection arthroplasty for the
300 treatment of infection. J Bone Joint Surg Am 1999; 81: 684.

301 28. Andrade SS, Bispo PJ, Gales AC. Advances in the microbiological diagnosis of sepsis.
302 Shock 2008; 30 Suppl 1: 41.

303 29. Arciola CR, Campoccia D, An YH, et al. Prevalence and antibiotic resistance of 15 minor
304 staphylococcal species colonizing orthopedic implants. Int J Artif Organs 2006; 29: 395.

305 30. Shuttleworth R, Behme RJ, McNabb A, et al. Human isolates of *Staphylococcus caprae*:
306 association with bone and joint infections. J Clin Microbiol 1997; 35: 2537.

307 31. Tarkin IS, Henry TJ, Fey PI, et al. PCR rapidly detects methicillin-resistant staphylococci
308 periprosthetic infection. Clin Orthop Relat Res 2003; 89.

309 32. Della Valle C, Parvizi J, Bauer TW, et al. Diagnosis of periprosthetic joint infections of the
310 hip and knee. J Am Acad Orthop Surg 2010; 18: 760.

311

312

313