

2-25-2016

Microcystic, Elongated, and Fragmented (MELF) Pattern Invasion in Ovarian Endometrioid Carcinoma: Immunohistochemical Profile and Prognostic Implications

Allison F. Goldberg, MD
Thomas Jefferson University

Lauren Cooper, MD
Thomas Jefferson University

D. De Cotiis
University of Pennsylvania

Follow this and additional works at: <https://jdc.jefferson.edu/pacbresidentposters>

Norman G. Rosenblum, MD, PhD

Part of the [Medical Anatomy Commons](#), [Medical Cell Biology Commons](#), and the [Medical Pathology Commons](#)

Joanna Chan, MD

Department of Pathology, Jefferson Medical College of Thomas Jefferson University, Philadelphia, PA

Recommended Citation

Goldberg, MD, Allison F.; Cooper, MD, Lauren; Cotiis, D. De; Rosenblum, MD, PhD, Norman G.; and Chan, MD, Joanna, "Microcystic, Elongated, and Fragmented (MELF) Pattern Invasion in Ovarian Endometrioid Carcinoma: Immunohistochemical Profile and Prognostic Implications" (2016). *Department of Pathology, Anatomy, and Cell Biology Resident's Posters*. Paper 8.
<https://jdc.jefferson.edu/pacbresidentposters/8>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Pathology, Anatomy, and Cell Biology Resident's Posters by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Microcystic, Elongated, and Fragmented (MELF) Pattern Invasion in Ovarian Endometrioid Carcinoma: Immunohistochemical Profile and Prognostic Implications

AF Goldberg¹, L Cooper², D De Cotiis³, N Rosenblum², JSY Chan¹

¹Department of Pathology, Anatomy, and Cell Biology, Thomas Jefferson University Hospital, Philadelphia, PA

²Department of Obstetrics and Gynecology, Thomas Jefferson University Hospital, Philadelphia, PA


³Department of Radiology, University of Pennsylvania, Philadelphia PA

BACKGROUND

- Microcystic, Elongated and Fragmented (MELF) is a well-recognized pattern of uterine endometrioid carcinoma (UEC) associated with lymphovascular space invasion and occult lymph node metastasis
- MELF in UEC may be seen with Lynch Syndrome
- MELF in UEC is hypothesized to be histologic evidence of an epithelial mesenchymal transition
- MELF pattern invasion in ovarian endometrioid carcinoma (OEC) was first described at USCAP 2015
- Current study evaluates MELF in OEC for
 - Prognostic implications
 - Immunohistochemical (IHC) profile related to
 - Lynch Syndrome
 - Epithelial mesenchymal transition

DESIGN

- 42 consecutive cases of OEC without concurrent UEC (1996-2014) evaluated by 2 pathologists
- MELF defined as at least three glands fulfilling histologic criteria
- 32 cases had blocks available for staining
 - MLH1, PMS2, MSH2 and MSH6 for mismatch repair (MMR) protein expression
 - Graded as “retained” or “lost”
 - β -catenin, e-cadherin, CK19 and cyclin D1 for evidence of epithelial mesenchymal transition
 - Graded as “rare” (<25% cells stain), “moderate” (25-75% cells stain), or “strong” (>75% cells stain)
- Retrospective chart review of clinical and demographic features and overall survival
- Data analyzed using Fisher exact test analysis
- Survival analyzed using Kaplan-Meier method



RESULTS

- MELF pattern invasion was identified in 45% of the cases reviewed
- Clear cell features were only seen in cases with MELF pattern invasion (p-value=0.044)
- Overall, 13% of cases demonstrate MMR protein loss
 - MELF: MSH2/MSH6 deficiency (n=2)
 - Non-MELF: PMS2 deficiency (n=2)
- No difference was identified in:
 - Overall survival
 - Cancer recurrence
 - IHC staining for β -catenin, e-cadherin, CK19 and cyclin D1
 - Serous features
 - Concurrent endometriosis
 - Lymphovascular space invasion
 - Lymph node metastasis
 - Bilaterality of disease
 - Extranodal metastasis

CONCLUSIONS

- MELF occurs in ovarian endometrioid carcinoma at a similar or higher frequency than in uterine endometrioid carcinoma.
- Clear cell features were identified exclusively in MELF pattern invasion cases.
- Different MMR proteins are lost in MELF and non-MELF pattern invasion carcinomas.
- As there is no current consensus on Lynch screening in patients with ovarian endometrioid carcinoma, perhaps the presence of MELF pattern invasion should prompt screening.
- MELF should be considered when assessing ovarian endometrioid carcinoma, as the pattern may be confused with endometriosis or endosalpingiosis.