

3-22-2017

Total Parenteral Nutrition in Patients Following Pancreaticoduodenectomy: Lessons from 1184 Patients

Cullen Worth

Thomas Jefferson University

Talar Tatarian, MD, PGY-4

Department of Surgery, Thomas Jefferson University

Awinder Singh

Thomas Jefferson University

Michael J. Pucci, MD

Thomas Jefferson University

Jordan M Winter, MD

Thomas Jefferson University

Follow this and additional works at: <https://jdc.jefferson.edu/surgeryposters>



Next page for additional authors

[Let us know how access to this document benefits you](#)

Recommended Citation

Worth, Cullen; Tatarian, MD, PGY-4, Talar; Singh, Awinder; Pucci, MD, Michael J.; Winter, MD, Jordan M; Yeo, MD, Charles J.; and Lavu, MD, Harish, "Total Parenteral Nutrition in Patients Following Pancreaticoduodenectomy: Lessons from 1184 Patients" (2017). *Department of Surgery Posters*. 7. <https://jdc.jefferson.edu/surgeryposters/7>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Surgery Posters by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Cullen Worsh; Talar Tatarian, MD, PGY-4; Awinder Singh; Michael J. Pucci, MD; Jordan M Winter, MD; Charles J. Yeo, MD; and Harish Lavu, MD

Total Parenteral Nutrition in Patients Following Pancreaticoduodenectomy: Lessons from 1184 Patients

Cullen E Worsh, Talar Tatarian, Awinder Singh, Michael J Pucci, Jordan M Winter, Charles J Yeo, Harish Lavu
Jefferson Pancreas, Biliary, and Related Cancer Center and the Department of Surgery, Thomas Jefferson University


BACKGROUND

- Total parenteral nutrition (TPN) has historically been used conservatively in the management of patients after pancreaticoduodenectomy (PD).
- In this study, we evaluate the indications for and outcomes associated with TPN use in a high-volume pancreatic surgery center.

METHODS

- Following IRB approval, all cases of PD from December 2006 through December 2015 were reviewed utilizing our pancreatic surgery database.
- Following identification of peri-operative TPN use, patients were separated into two cohorts with the hypothesis that was those patients who were started on TPN but received it for less than 3 days due to an improvement in their clinical condition, most likely were overprescribed.

Figure 1. Study Design



RESULTS

Table 1. Patient Demographics and Operative Data

	No TPN n=952 (80%)	TPN n=232 (20%)	p-value
Male	453 (48)	156 (67)	0.0001
Age	66.8 (18-92)	69.3 (37-90)	0.007
BMI	25.6 (10.8-52.6)	26 (15-43.9)	0.30
Diabetic	245 (26)	60 (26)	1.00
Smoker (Current or Past)	448 (47)	131 (56)	0.01
Pylorus Preservation	801 (84)	183 (79)	0.06
EBL (mL)	430 (50-1600)	630 (40-8000)	0.08
Soft Gland (n=872;n=204)	421 (48)	114 (56)	0.05
Pathology			
PDA	601 (63)	164 (71)	0.03
IPMN	138 (15)	27 (12)	0.29
Pancreatitis	52 (5)	6 (3)	0.09

Table 2. Indications for and Patterns of TPN Use

Complication	TPN ≤3 Days n=44 (19%)	TPN >3 Days n=188 (81%)	p-value
Any Complication	39 (89)	179 (95)	0.15
DGE	32 (73)	139 (74)	0.85
Grade A	25 (78)	59 (42)	0.0003
Grade B	6 (19)	53 (38)	0.04
Grade C	1 (3)	27 (19)	0.03
Pancreatic Fistula	8 (18)	94 (50)	0.0001
Grade A	5 (63)	24 (26)	0.04
Grade B	2 (25)	57 (61)	0.07
Grade C	1 (13)	13 (14)	1.00
Deep Surgical Site Infection	7 (16)	91 (48)	0.0001
Pneumoniae	0 (0)	31 (16)	0.001

Table 3. Postoperative Imaging

Patient Imaging	TPN ≤3 Days n=44 (%)	TPN >3 Days n=188 (%)	p-value
Upper GI Imaging	20 (46)	109 (58)	0.18
Normal function	5 (25)	35 (32)	0.61
Aperistalsis	6 (30)	40 (37)	0.62
Anastomotic edema	8 (40)	18 (17)	0.03
CT Imaging	22 (50)	157 (84)	0.0001
Fluid collection	17 (77)	126 (80)	0.78
Drainable collections	5 (23)	94 (60)	0.001

Table 4. Post TPN Outcomes

Postoperative Measure	TPN ≤3 Days n=44 (%)	TPN >3 Days n=188 (%)	p-value
CLABSI	0 (0)	6 (3)	0.60
Total Days on TPN	2 (1-3)	12 (4-115)	0.0001
LOS (days)	8.5 (6-16)	14 (3-92)	0.0005
Readmission	16 (36)	66 (35)	0.86
30-Day Mortality	3 (7)	5 (3)	0.18

RESULTS

Table 5. Multivariate Logistic Regression Analysis of Factors Affecting Length of TPN Use

Risk Factor	Odds Ratio	95% CI	p-value
Diabetes	0.49	0.24-1.03	0.060
DGE	1.22	0.57-2.60	0.105
Pancreatic Fistula	2.57	1.03-6.41	0.043
Deep Surgical Site Infection	3.09	1.16-5.06	0.018

KEY FINDINGS

- The most common indications for TPN were delayed gastric emptying (DGE, n=171, 73.7%), pancreatic fistula (PF, n=102, 44%), and generalized malnutrition (n=25, 10.8%).
- The median day of TPN initiation was POD 4 (range: minus 31 to 22), with a median usage of nine days (range: 1 to 115).
- Forty-four (19%) patients received short-course TPN (≤3 days), primarily those diagnosed with isolated Grade A DGE without associated complications (p=0.0003).
- Multivariate analysis suggests the presence of deep surgical site infection (OR 3.09, [1.16 - 5.06], p=0.018) or PF (OR 2.57, [1.03 - 6.41], p=0.043) at the time of DGE presentation as predictive of long-term TPN requirement.

CONCLUSIONS

- TPN is safe and effective in patients following PD.
- Avoid TPN in patients with isolated DGE secondary to anastomotic edema.
- Use TPN in patients with DGE and another serious complication such as deep surgical site infection or PF.