

5-24-2016

Factors Influencing Resection in Locoregional Pancreatic Cancer Patients

Amy Cunningham, MPH, PhD(c)

Jefferson College of Population Health, Thomas Jefferson University

Brian P.H. Chen, ScM, PharmD

Jefferson College of Population Health, Thomas Jefferson University

David Delgado, PhD

Jefferson College of Population Health, Thomas Jefferson University

Follow this and additional works at: <https://jdc.jefferson.edu/jcphposters>

 Part of the [Public Health Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Cunningham, MPH, PhD(c), Amy; Chen, ScM, PharmD, Brian P.H.; and Delgado, PhD, David, "Factors Influencing Resection in Locoregional Pancreatic Cancer Patients" (2016). *College of Population Health Posters*. 9.

<https://jdc.jefferson.edu/jcphposters/9>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in College of Population Health Posters by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.



HOME OF SIDNEY KIMMEL MEDICAL COLLEGE

Factors Influencing Resection in Locoregional Pancreatic Cancer Patients

Amy T. Cunningham, MPH, PhD(c), Brian Po-Han Chen, ScM, David Delgado, PhD
Thomas Jefferson University College of Population Health, Philadelphia, PA

Background

- Pancreatic cancer has surpassed breast cancer as the **3rd leading cause of U.S. cancer deaths**, with 41,780 deaths in 2016¹
- Projected to be **2nd leading cause of cancer death by 2030**²
- Mean direct medical costs of **\$65,700 per patient** (2000-2007)³
- **Five-year survival rate is 7.7%**¹, due to often advanced stage at diagnosis, lack of effective treatment options
- **Five-year survival rate for pancreatic cancer patients undergoing resection, the only potentially curative treatment, is 18-24%**⁴
- Pancreatic cancer resection is underused in eligible patients⁵⁻⁷
- **Factors associated with underuse of pancreatic cancer resection are poorly understood**

Objective

Identify factors associated with resection in a national sample of locoregional pancreatic cancer patients

Methods

Data Source

- Surveillance, Epidemiology and End Results (SEER) dataset⁸
- National Cancer Institute cancer statistics program
- Cancer demographics, incidence, treatment and survival data
- Combines data from 20 regional and state registries; covers 30% of U.S. population
- 8,689,771 total cancer cases from 1973-2014

Study Population

Inclusion criteria

- 2004-2012 SEER patients, age 15-89, with primary diagnosis of locoregional (Stage I or II) pancreatic cancer in the pancreas head, body, or tail
- Staging based on American Joint Committee on Cancer (AJCC6)⁹ criteria instituted in 2004

Exclusion criteria:

- Pancreatic cancer diagnosis via death certificate or autopsy
- Incomplete survival data
- Incomplete surgical data

Statistical Analyses

- Primary outcome: receipt of pancreatic cancer resection
- Descriptive statistics to characterize sample
- Chi-square tests to identify associations between demographic variables and pancreatic cancer resection
- Multivariate logistic regression to build final model of associations between covariates and pancreatic cancer resection
- Model fit assessed using Hosmer-Lemeshow test
- All analyses performed using Statistical Analysis System (SAS) Studio software¹⁰


Results

Final sample: 16,676 locoregional pancreatic cancer patients (Figure 1, Table 1), of whom **8152 (48.9%)** did not receive surgery (Figure 2)

Variables associated with not receiving surgery (controlling for registry, stage, and tumor location)(Table 2):

1. **Age:** 65-74 **30%** less likely; 75 or older **72%** less likely to undergo resection
2. **Sex:** Men **9%** less likely to undergo resection
3. **Race/ethnicity:** Non-Hispanic black individuals **31%** less likely, Hispanic **20%** less likely, other non-white race/ethnicity **14%** less likely to undergo resection
4. **Marital status:** Unmarried individuals **29%** less likely to undergo resection

Figure 1. Stepwise Cohort Ascertainment



Results(contd.)

Table 1. Demographics of final sample

Characteristic	Resection N(%)	No resection N(%)	p-value
Total (N=16,676)	8524(51.1)	8152(48.9)	
Age Category			<.001
<65	4088(48.0)	2325(28.5)	
65-74	2647(31.1)	2120(26.0)	
≥75	1780(21.0)	3707(45.5)	
Sex			<.001
Male	4290(52.7)	3857(47.3)	
Female	4234(49.6)	4295(50.4)	
Race/Ethnicity			<.001
White/Non Hispanic	6187(52.2)	5665(47.8)	
Non-Hispanic Black	879(45.9)	1034(54.1)	
Hispanic	804(48.9)	840(51.1)	
Other Race/Ethnicity	654(51.6)	613(48.4)	
Marital Status			<.001
Married	5328(56.2)	44159(43.8)	
Not married	3196(44.5)	3993(55.5)	
Registry			<.001
Diagnosis Year			.313
2004-2006	2390(49.9)	2398(50.1)	
2007-2009	2934(52.1)	2699(47.9)	
2010-2012	3200(51.2)	3055(48.8)	
AJCC Stage			<.001
Stage I	1537(39.7)	2332(60.3)	
Stage II	6987(54.6)	5820(45.4)	
Tumor Location			<.001
Head	6930(49.8)	6994(50.2)	
Body	555(41.0)	799(59.0)	
Tail	1039(74.3)	359(25.7)	

Figure 2. Reason Given for No Surgery (N=8152)




Table 2. Odds of Undergoing Resection

Variable	Univariate Analysis Odds Ratio (CI)	Multivariate Analysis* Odds Ratio (CI)
Age Category		
>65	Reference	Reference
65-74	.71 [.66-.77]	.70 [.65-.76]
≥ 75	.28 [.26-.30]	.28 [.25-.30]
Sex		
Female	Reference	Reference
Male	.89 [.83-.94]	.91 [.85-.98]
Race/Ethnicity		
White/Non-Hispanic	Reference	Reference
Non-Hispanic Black	.78 [.71-.86]	.69 [.62-.77]
Hispanic	.88 [.79-.97]	.80 [.71-.89]
Other race/ethnicity	.98 [.87-1.10]	.86 [.75-.98]
Marital Status		
Married	Reference	Reference
Not married	.63 [.59-.67]	.71 [.66-.76]

* Controlling for registry, stage, and tumor location

Discussion

- Nearly half of patients with locoregional pancreatic cancer do not receive surgery; the reason why is often unclear
- Factors associated with lower resection rates are non-white race/ethnicity, older age, male sex and being unmarried
- Understanding and addressing these disparities could increase pancreatic cancer resection rates and improve survival

Limitations

- Observational data limits ability to make causal inferences
- Some variables associated with resection (comorbidities, insurance, socioeconomic status)⁴ are not available in SEER
- No details on resection decision-making process

Recommendations for Future Research

- Perform sensitivity analysis and instrumental variable analysis
- Explore SEER-Medicare linked data
- Analyze regional care patterns and the impact of high-volume pancreatic surgery hospitals on resection rates
- Conduct interviews with physicians, patients and their caregivers on resection decision-making process

Acknowledgements

The authors would like to thank Harish Lavu, MD, FACS, Associate Professor, Thomas Jefferson University Department of Surgery, for his review of the poster abstract.

References

1. National Cancer Institute. SEER Stat Fact Sheets: Pancreas Cancer. 2016; retrieved from: <http://seer.cancer.gov/statfacts/html/pancreas.html>
2. Rahib L, Smith BD, Aizenberg R, Rosenzweig AB, Fleshman JM, Matrisian LM. Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and pancreas cancers in the United States. Cancer research. 2014 Jun 1;74(11):2913-21.
3. O'Neill CB, Atona CL, O'Reilly EM, LaFemina J, Henman MC, Elkin EB. Costs and trends in pancreatic cancer treatment. Cancer. 2012 Oct 15;118(20):5132-9.
4. National Cancer Institute. Pancreatic Cancer Treatment –Health Professional Version (PDQ®). General information about pancreatic cancer. 2016; retrieved from <http://www.cancer.gov/types/pancreatic/hp/pancreatic-treatment-pdq>
5. Blimoria KY, Bentrem DJ, Ko CY, Stewart AK, Winchester DP, Talamonti MS. National failure to operate on early stage pancreatic cancer. Annals of surgery. 2007 Aug 1;246(2):173-80.
6. Riall TS, Townsend CM, Kuo YF, Freeman JL, Goodwin JS. Dissecting racial disparities in the treatment of patients with locoregional pancreatic cancer. Cancer. 2010 Feb 15;116(4):930-9.
7. Enevoild L, Harlan LC, Tucker T, McKenzie S. Pancreatic cancer in the USA: persistence of undertreatment and poor outcome. Journal of gastrointestinal cancer. 2015 Mar 1;46(1):9-20.
8. Surveillance, Epidemiology, and End Results (SEER) Program (www.seer.cancer.gov) Research Data (1973-2012), National Cancer Institute, DCCPS, Surveillance Research Program, Surveillance Systems Branch, released April 2015, based on the November 2014 submission.
9. American Joint Committee on Cancer (AJCC). Pancreas cancer staging, 6th edition. 2002.
10. Statistical Analysis System (SAS) Studio. Copyright © 2016 SAS Institute Inc. Cary, NC, USA.