

2-2019

BOVA is Superior to sPESI in Identification of High Risk Pulmonary Embolism Patients

J. DiBello

Thomas Jefferson University, joseph.dibello@jefferson.edu

P. Khaing

Thomas Jefferson University, phue.khaing@jefferson.edu

E. Gupta

Thomas Jefferson University, ena.gupta@jefferson.edu

C. Au

Thomas Jefferson University

T. Chengsupanimit

*Thomas Jefferson University, tayoot.chengsupanimit@jefferson.edu**See next page for additional authors*

[Let us know how access to this document benefits you](#)

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2021_phase1 Part of the [Critical Care Commons](#), and the [Pulmonology Commons](#)

Recommended Citation

DiBello, J.; Gupta, E.; Au, C.; Chengsupanimit, T.; Awsare, B.K.; Baram, M.; West, F.M.; and Kane, G.C., "BOVA is Superior to sPESI in Identification of High Risk Pulmonary Embolism Patients" (2019). SKMC JeffMD Scholarly Inquiry, Phase 1, Project 1.

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

J. DiBello, P. Khaing, E. Gupta, C. Au, T. Chengsupanimit, B. K. Awsare, M. Baram, F. M. West, and G. C. Kane

Joe DiBello
SKMC Class of 2021
SI CTR Abstract
12/15/18

Title: BOVA is Superior to sPESI in Identification of High Risk Pulmonary Embolism Patients

J. DiBello¹, P. Khaing², E. Gupta², C. Au¹, B.K. Awsare², M. Baram², F.M. West², G.C. Kane²

1. Sidney Kimmel Medical College at Thomas Jefferson University Hospital, Philadelphia, PA
2. Division of Pulmonary and Critical Care Medicine, Jane and Leonard Korman Respiratory Institute, Sidney Kimmel Medical College at Thomas Jefferson University Hospital, Philadelphia, PA

Introduction: Prognostic models exist for the purpose of stratifying patients with acute pulmonary embolism. Of these, the Pulmonary Embolism Severity Index (PESI) and the simplified PESI (sPESI) are the most well-known, although more recent composite models, like the BOVA score, are now being studied and implemented. Comparative efficacy of these scores to predict long term mortality is not well established.

Methods: We performed a retrospective analysis of all consecutive patients diagnosed with PE using computed tomography scan from 2014-2016 at an urban tertiary-referral medical center. Cox proportional hazard analyses were performed to compare the performance of two prognostic models – sPESI and BOVA – to predict all-cause in-hospital and cumulative one-year mortality.

Results: The all-cause in-hospital mortality rate was 6.0%, and cumulative one-year mortality rate was 21.3%.

In adjusted analyses, a BOVA score >4 was significantly associated with an increased in-hospital mortality (HR 3.5, 95% CI: 1.4-9.0, p = 0.009) and one-year mortality (HR 2.0, 95% CI: 1.0-3.9, p = 0.04), as compared to a BOVA score <4. However, the sPESI (p = 0.14) did not show a significant association with one-year mortality.

In identifying in-hospital mortality, the sPESI had high sensitivity (100%) and low specificity (10.1%), whereas the BOVA score had low sensitivity (20.0%) and high specificity (92.7%).

Similar trends were seen for one-year mortality.

Conclusion: In this study, a high BOVA score was found to be the best predictor of both short and long-term mortality in PE patients. A low sPESI score identified with high sensitivity patients with low-risk PEs.