

1-2020

## Anxiety and Depressive Symptoms Pre-Trauma as a Predictor of Dissociative Symptoms Post-Trauma – An AURORA Study

Vishal Swaminathan

Thomas Jefferson University, [vishal.swaminathan@jefferson.edu](mailto:vishal.swaminathan@jefferson.edu)

Follow this and additional works at: [https://jdc.jefferson.edu/si\\_dh\\_2022\\_phase1](https://jdc.jefferson.edu/si_dh_2022_phase1)

 Part of the [Mental and Social Health Commons](#), and the [Psychiatry Commons](#)

[Let us know how access to this document benefits you](#)

---

### Recommended Citation

Swaminathan, Vishal, "Anxiety and Depressive Symptoms Pre-Trauma as a Predictor of Dissociative Symptoms Post-Trauma – An AURORA Study" (2020). *Phase 1*. Paper 4.  
[https://jdc.jefferson.edu/si\\_dh\\_2022\\_phase1/4](https://jdc.jefferson.edu/si_dh_2022_phase1/4)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

## **SKMC Class of 2022: SI/DH Abstract**

**Word count: 300**

### **Anxiety and Depressive Symptoms Pre-Trauma as a Predictor of Dissociative Symptoms Post-Trauma – An AURORA Study**

**Vishal Swaminathan, Dr. AnnaMarie Chang\***

(\*) indicates primary project advisor

#### **Introduction:**

Increased states of anxiety or depression may lead to increased occurrences of dissociative behavior, justifying the need to validate pre-traumatic cognitive states as a predictor of dissociation post-trauma. Early prediction can lead to medical intervention for high-risk patients and prevent potential adverse sequelae. The AURORA Study evaluates biomarkers and neuropsychological sequelae in patients immediately following traumatic events. We predict patients who suffered from depressive symptoms and anxiety prior to the traumatic event will experience increased dissociative symptoms post-trauma.

#### **Methods:**

This study includes survey data collected at ED's from survivors (n = 666) of traumatic motorized vehicle collisions (MVC) at time points 30 days pre-collision, and 2- and 8-weeks post-collision. Inclusion criteria included those enrolled until the closure of the week 8 survey window and completed at least 90% of follow up surveys at weeks 2 and 8. Our primary outcomes were dissociative symptoms at 2- and 8- weeks post-

MVC. We adjusted for sex at birth, age, race/ethnicity, and overall mental health prior to the MVA. Simple and multiple linear regression models were utilized for analysis, along with ANOVA.

**Results:**

Anxiety was not a significant predictor for dissociative symptoms at 2- (0.042, 95% CI -0.11-0.096, P=0.12) and 8- weeks (0.030, 95% CI -0.11-0.096, P=0.275). Depressive symptoms were a significant predictor at 2- (0.032, 95% CI 0.006-0.058, P=0.016) and 8- weeks (0.042, 95% CI 0.016-0.069, P=0.002). Additionally, overall mental health was a significant predictor at 2 weeks (-0.027, 95% CI -0.047- -0.008, P=0.006), and white ethnicity was significant at 8 weeks (-0.418, 95% CI -0.757- -0.079, P=0.016).

**Discussion:**

Symptoms of anxiety prior to traumatic events are reliable predictors of post-traumatic stress, while depressive symptoms are not. Additional long-term data points are needed, but preliminary analysis justifies the potential use of pre-traumatic anxiety for early medical intervention before deterioration of neurocognitive functioning.