

8-29-2024

## Lower Extremity Injury Rates on Artificial Turf Versus Natural Grass Surfaces in the National Football League During the 2021 and 2022 Seasons

Nikit Venishetty

Angel X Xiao

Ramesh Ghanta

Rohit Reddy

Nirav Kiritkumar Pandya

*See next page for additional authors*

Follow this and additional works at: <https://jdc.jefferson.edu/skmcstudentworks>



Part of the [Orthopedics Commons](#)

**[Let us know how access to this document benefits you](#)**

---

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in SKMC Student Presentations and Publications by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

---

**Authors**

Nikit Venishetty, Angel X Xiao, Ramesh Ghanta, Rohit Reddy, Nirav Kiritkumar Pandya, and Brian T Feeley

# Lower Extremity Injury Rates on Artificial Turf Versus Natural Grass Surfaces in the National Football League During the 2021 and 2022 Seasons

Nikit Venishetty,\* BA, Angel X. Xiao,<sup>†</sup> MD, Ramesh Ghanta,<sup>†</sup> MD, Rohit Reddy,<sup>‡</sup> BS, Nirav Kiritkumar Pandya,<sup>†</sup> MD, and Brian T. Feeley,<sup>†§</sup> MD  
*Investigation performed at the Department of Orthopaedic Surgery, University of California, San Francisco, California, USA*

**Background:** It has been argued that the use of artificial turf football fields in the National Football League (NFL) increases player injury risk compared with natural grass surfaces.

**Purpose/Hypothesis:** The purpose of this study was to quantify the rate of lower extremity injuries occurring in NFL players on artificial turf compared with natural grass surfaces and characterize the time missed due to injury and proportion of injuries requiring surgery. It was hypothesized that lower extremity injuries requiring surgical intervention would occur at a higher rate on artificial turf than on natural grass.

**Study Design:** Descriptive epidemiology study.

**Methods:** Lower extremity injury data for the 2021 and 2022 NFL seasons were obtained using publicly available records. Data collected included injury type, player position, player age, playing surface, weeks missed due to injury, and whether the patient underwent season-ending or minor surgery. Multivariable logistic regression was performed to determine the risk of season-ending surgery according to playing surface.

**Results:** When combining injuries for the 2021 and 2022 seasons (N = 718 injuries), the incidence rate of lower extremity injury was 1.22 injuries/game for natural grass and 1.42 injuries/game for artificial turf. The odds of a season-ending surgery were found to be significantly higher on artificial turf compared with natural grass (odds ratio = 1.60; 95% CI, 1.28-1.99;  $P < .05$ ), while additional variables, including weather, age, position, week of injury occurrence, and history of prior injury, did not influence the odds of season-ending surgery.

**Conclusion:** The 2021 and 2022 NFL seasons of our analysis demonstrated a higher incidence rate of injuries on artificial turf surfaces compared with natural grass surfaces. In addition, the odds of injury requiring season-ending surgery were found to be significantly higher on artificial turf compared with natural grass.

**Keywords:** NFL injuries; playing surfaces; season-ending surgeries; natural grass; artificial turf

As the emphasis on player safety in the National Football League (NFL) has increased in recent years, numerous risk factors for injury have been closely scrutinized for their roles in placing players at increased risk. One such factor, the use of artificial turf as a playing surface, has become a controversial topic, with many current and former players stating that football fields that use

artificial turf place players at higher risk for injury, while alternative sources have suggested there is no significant injury rate between artificial and natural grass surfaces.<sup>12-14,21,23,24</sup>

Artificial playing fields have been utilized in the NFL for the past 5 decades, with the first installed in the Houston Astrodome in 1966.<sup>19</sup> Overtime, these artificial fields have transformed from carpet-like asphalt bases to the current use of permeable elastomeric layers of recycled rubber, polyurethane, and mineral aggregate.<sup>23</sup> Recently, designers of these fields have transitioned into using infill

The Orthopaedic Journal of Sports Medicine, 12(8), 23259671241265378  
 DOI: 10.1177/23259671241265378  
 © The Author(s) 2024

This open-access article is published and distributed under the Creative Commons Attribution - NonCommercial - No Derivatives License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits the noncommercial use, distribution, and reproduction of the article in any medium, provided the original author and source are credited. You may not alter, transform, or build upon this article without the permission of the Author(s). For article reuse guidelines, please visit SAGE's website at <http://www.sagepub.com/journals-permissions>.

surfaces, a mat consisting of long polyethylene fibers with spaces between the fibers that are filled with rubber particles or sand.<sup>4</sup> By the start of the 2005 NFL season, all NFL stadiums with artificial playing fields had installed infill surfaces in place of the older generation artificial surfaces.<sup>4</sup>

With the popularity of artificial turf football fields, there has been discordance regarding player safety, especially at the professional level. Previous studies that have assessed injuries on collegiate and professional playing fields have found statistically significantly higher rates of injuries on artificial surfaces in comparison with those on natural grass.<sup>4,18,21,23</sup> Among these studies, there was variability in the consistency of study design and statistical significance, prompting future studies to assess the rate of injuries on artificial turf playing surfaces.<sup>7,15</sup>

The purpose of this study was to quantify the rate of lower extremity injuries occurring in NFL players on artificial turf compared with natural grass surfaces and characterize the time missed due to injury and proportion of injuries requiring surgery. We hypothesized that lower extremity injuries requiring surgical intervention would occur at a higher rate on artificial turf than on natural grass.

## METHODS

Lower extremity injury data for the 2021 and 2022 NFL seasons were obtained using publicly available records from the internet ([www.ESPN.com](http://www.ESPN.com), [www.NFL.com](http://www.NFL.com)).<sup>16,17</sup> The databases on ESPN provided in-game injury information regarding NFL players from the season. As we were unable to obtain information regarding injuries in practice, we focused on in-game injuries that were reported on the ESPN game summaries. All reported lower extremity injuries during the regular and playoff seasons were analyzed. Upper extremity, head, and back injuries were excluded. Preseason data were also excluded due to the unavailability of injuries reported on the ESPN game summaries. Data collected included injury type, player position and age, playing surface, weeks missed due to injury, and whether the patient underwent minor surgery or major (season-ending) surgery.

Statistical analysis was performed using R Software (Rlabs). Descriptive statistics, including percentages and counts for categorical and ordinal data and means with interquartile ranges for continuous data, were used to summarize data. The injury incidence was calculated for the 2021 and 2022 seasons by calculating the ratio of the number of injuries that occurred on grass and artificial

TABLE 1  
Summary of Lower Extremity Injury  
Data Collected (N = 718 Injuries)<sup>a</sup>

Variable	Value
Injuries by year	
2021	327
2022	391
Playing position	
Control (quarterback, kicker, punter)	22
Defensive line	82
Offensive line	127
Running back, fullback, tight end	134
Secondary (cornerback, safety)	151
Wide receiver	96
Unspecified <sup>b</sup>	106
Age, y, mean (range)	26.29 (20-38)
Injuries by playing surface	
Artificial turf	386
Natural grass	332
History of injury at injured site	100

<sup>a</sup>Data are reported as No. of injuries unless otherwise indicated.

<sup>b</sup>Players without designated positions on team (special teams, practice squad etc.).

turf and the number of games played on the respective surfaces. Additionally, injuries were stratified into 4 categories according to the number of games missed: 0, 1-2, 3-6, or >6. Multivariable logistic regression was performed to determine the influence of playing surface on the risk of major surgery.  $P < .05$  was considered statistically significant.

## RESULTS

A total of 718 lower extremity injuries were included in the analysis, 327 injuries (45.54%) in 2021 and 391 injuries (54.46%) in 2022. A demographic summary of these injuries by position, age, field, and history of previous injury is displayed in Table 1. Within the NFL as of the end of the 2022 season, there were 16 stadiums surfaced with natural grass and 13 stadiums with artificial turf (Table 2).

To account for injury severity, we documented the subsequent weeks of gameplay missed after an injury. The incidence of these injuries by severity is shown in Table 3 and Figure 1. The overall incidence rate of lower extremity injuries for the combined 2021 and 2022 seasons was 1.22 injuries/game for natural grass and 1.42 injuries/game for artificial turf.

<sup>§</sup>Address correspondence to Brian T. Feeley, MD, Department of Orthopaedic Surgery, University of California, San Francisco, 1500 Owens Street, San Francisco, CA 94158, USA (email: [brian.feeley@ucsf.edu](mailto:brian.feeley@ucsf.edu)) (Twitter/X: DrBrianFeeley).

\*Paul L. Foster School of Medicine, Texas Tech Health Sciences Center, El Paso, Texas, USA.

<sup>†</sup>Department of Orthopaedic Surgery, University of California, San Francisco, San Francisco, California, USA.

<sup>‡</sup>Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, Pennsylvania, USA.

Final revision submitted November 18, 2023; accepted February 12, 2024.

One or more of the authors has declared the following potential conflict of interest or source of funding: N.K.P. has received education payments from Evolution Surgical. B.T.F. has received education payments from Evolution Surgical. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.

Ethical approval was not sought for the present study.

TABLE 2  
Summary of Stadiums in the NFL According to Field Surface<sup>a</sup>

Stadium Name (NFL Team)	Natural Grass	Artificial Turf
Empower Field at Mile High (Denver Broncos)		AT&T Stadium (Dallas Cowboys)
Allegiant Stadium, Las Vegas (Las Vegas Raiders)		Bank of America Stadium (Carolina Panthers)
Arrowhead (Kansas City Chiefs)		Caesars Superdome (New Orleans Saints)
FedEx (Washington Commanders)		Ford Field (Detroit Lions)
First Energy Stadium (Cleveland Browns)		Gillette Stadium (New England Patriots)
Hard Rock Stadium (Miami Dolphins)		Highmark Stadium (Buffalo Bills)
Heinz Field (Pittsburgh Steelers)		Lucas Oil Field (Indianapolis Colts)
Lambeau Stadium (Green Bay Packers)		Lumen Filed (Seattle Seahawks)
Levi stadium (San Francisco 49ers)		Mercedez-Benz Stadium (Atlanta Falcons)
Lincoln Financial Field (Philadelphia Eagles)		MetLife Stadium (New York Giants/Jets)
M&T Bank Stadium (Baltimore Ravens)		NRG Stadium (Houston Texans)
Nissan Stadium (Tennessee Titans)		Paycor Stadium (Cincinnati Bengals)
Raymond James Stadium (Tampa Bay Buccaneers)		SoFi Stadium (Los Angeles Rams/Chargers)
Soldier Field (Chicago Bears)		
State Farm Stadium (Arizona Cardinals)		
TIAA Bank Field (Jacksonville Jaguars)		

<sup>a</sup>NFL, National Football League.

TABLE 3  
Injuries Within the 2021 and 2022 NFL Seasons Stratified by Amount of Time Missed<sup>a</sup>

Season	Total	Games Missed				Surgeries <sup>b</sup>
		0	1-2	3-6	>6	
2021						
Injuries on natural grass	171	54	35	46	37	24
Incidence (injuries/game)	1.25	0.39	0.26	0.34	0.26	0.18
Injuries on artificial turf	156	41	36	39	40	36
Incidence (injuries/game)	1.16	0.30	0.27	0.29	0.29	0.27
2022						
Injuries on natural grass	161	44	44	31	45	26
Incidence (injuries/game)	1.18	0.33	0.33	0.21	0.33	0.19
Injuries on artificial turf	230	59	64	44	63	47
Incidence (injuries/game)	1.68	0.43	0.47	0.34	0.44	0.34
Combined						
Injuries on natural grass	332	98	79	77	78	50
Incidence (injuries/game)	1.22	0.36	0.29	0.28	0.29	0.18
Injuries on artificial turf	386	100	100	83	102	83
Incidence (injuries/game)	1.42	0.37	0.37	0.31	0.38	0.31


<sup>a</sup>Data are reported as No. NFL, National Football League.

<sup>b</sup>Data reported as number of surgeries on natural grass vs artificial turf, and their corresponding incidence rates.

In total, there were 50 players injured on natural grass who required lower extremity surgery and 83 players injured on artificial turf who required lower extremity surgeries. The most common surgeries after a lower extremity on grass were anterior cruciate ligament (ACL) reconstruction (n = 17), Achilles repair (n = 9), and ankle or syndesmosis open reduction and internal fixation (n = 7). The most common surgeries after a lower extremity on artificial turf were ACL reconstruction (n = 30), Achilles repair

(n = 12), and patellar open reduction and internal fixation or patellar tendon repair (n = 8) (Table 4).

Bivariate modeling showed that the history of previous injury as well as field type influenced the risk of surgery. In the multivariate model, the odds of season-ending surgery were significantly higher on artificial turf compared with grass (odds ratio = 1.60; 95% CI, 1.28-1.99; *P* < .05). Weather, age, position, week, and history of previous injury were not significant influences on the odds of season-ending surgery.



**Figure 1.** Injuries on artificial turf versus natural grass during the 2021 and 2022 National Football League seasons.

## DISCUSSION

The study findings indicated that NFL players were at a higher incidence of developing lower extremity injuries on artificial turf compared with natural grass. On top of this, the odds of a more serious injury requiring season-ending surgery were found to be significantly higher on artificial turf compared with natural grass. The findings from our study contribute to the ongoing discussion of injury differences while playing on artificial turf surfaces versus natural grass, with the hope of prioritizing player safety.

Despite significant advancements in equipment, a growing emphasis on player safety, and the progress made in treatments, the persisting rate of injuries in the NFL has sparked ongoing debates regarding the safety of playing on artificial turf fields. Previous studies have demonstrated an increase in injury risk in artificial turf versus natural grass surfaces.<sup>12-14,21,23,24</sup> However, there has been no recent assessment of the differences. The purpose of this study was to compare the injury rates in the NFL for artificial turf and natural grass over the 2 seasons of 2021 and 2022.

Players in the NFL and the NFL Players Association have voiced concerns regarding the safety of artificial turf fields.<sup>2-20</sup> However, there has been substantial push-back from several NFL owners and the league itself. Concerns from players throughout the professional league have warranted studies on the differences in injury rates between artificial turf and natural grass playing fields. Our results are similar to a previous study, as we found that artificial turf is a serious risk factor for increasing lower extremity injury.<sup>9</sup> Moreover, a study that looked at injury rates from 2000 to 2009 in the NFL similarly found a 22% increase in both knee and ankle injuries on FieldTurf in comparison with natural grass.<sup>4</sup> Our findings align closely with the results of both studies,

**TABLE 4**  
Summary of Lower Extremities Surgeries  
Performed in the 2021 and 2022 NFL Seasons<sup>a</sup>

	Artificial Turf	Natural Grass
Achilles tendon repair	12	9
ACL reconstruction	30	17
Meniscus	2	2
Patellar ORIF or tendon repair	8	3
Hamstring repair	1	1
Ankle or syndesmotic ORIF	7	7
Foot ORIF	6	3
Lisfranc repair	2	1
MCL repair	4	—
Quadriceps repair	1	2
Multiligamentous knee	2	2
Adductor repair	1	—
Knee surgery, unspecified	4	3
Hip surgery, unspecified	1	—
Tibial ORIF	2	—
<b>Total</b>	<b>83</b>	<b>50</b>

<sup>a</sup>Data are reported as No. Dashes indicate data not reported. ACL, anterior cruciate ligament; MCL, medial collateral ligament; ORIF, open reduction and internal fixation.

revealing a notable trend: over the past 3 years, there has been a substantial surge (20%) in lower extremity injuries on FieldTurf in comparison with those sustained on natural grass.<sup>3</sup> It is thought that the increasingly integrated use of analytics and higher reporting of in-game injuries has led to this increase, despite advancements in player equipment.<sup>9</sup>

Additionally, a recent systematic review analyzed 53 articles across football, soccer, and other sports. Of the included articles, 32 compared injury rates on artificial turf and natural grass, and >37.5% (since over 12) reported a higher overall injury rate on artificial turf. Of the 32 articles, 3 (9.4%) reported a higher overall injury rate on natural grass.<sup>3</sup> However, it was interesting to note that all 3 of those studies utilized ad hoc cohort selection and were supported by funding from the artificial turf industry.<sup>10-12</sup> Our findings corroborate previous studies and underscore the importance of addressing injury risks on artificial turf fields.

While these injuries are severely affecting players in the NFL, the injuries associated with playing on artificial turf fields extend to the collegiate and high school levels, further emphasizing the increased injury risk. In a separate study, 2460 knee injuries sustained over the course of 10 National Collegiate Athletic Association (NCAA) seasons were examined by Loughran et al.<sup>8</sup> They came to the conclusion that using artificial turf increases the chance of sustaining certain knee ligament ailments. According to their research, posterior cruciate ligament tears were substantially more common during competitions played on artificial turf than on natural grass. During competitions on artificial turf as opposed to natural grass, lower NCAA divisions (II and III) were found to have greater rates of ACL injuries.<sup>8</sup>

Interestingly, in our data analysis, we found a significantly higher number of extensor mechanism injuries and a very high number of ACL injuries in artificial turf fields versus natural grass fields.

There are several technical and clinical indications that can be drawn from our findings. First, it is important to continually evolve synthetic turf to maintain its desirable traits but also emphasize player safety. Its increased use in recent years warrants further evaluation to protect players from season-ending injuries that could significantly alter their future. The altering of the future can start at a young age, as studies have found that high school athletes are 58% more likely to sustain injuries on artificial turf than on natural grass.<sup>22</sup> Not only does this negatively affect their future health but it also affects their ability to go into professional sports in the future.

The capacity of natural grass and artificial turf to develop divots and release the cleat at loading magnitudes and rates generated during elite sports competition has been demonstrated by biomechanical testing of various football cleats on a range of athletic surfaces.<sup>5,6</sup> Artificial turf is known to have less absorption of force for the player's knee and ankle, which increases the risk of injury.<sup>5,6,9</sup> The lesser ability of artificial turf to release a cleat puts more strain on a player's extremities through greater force and increased torque.<sup>5,6,9</sup> On the other hand, natural grass has been shown to often release the cleat before reaching an injurious load.<sup>1,5,6,9</sup> As a result, it is crucial that its design takes into account the type of surface that will be utilized with it as well as the precise mechanics of how its cleat pattern will interface with synthetic surfaces.

### Limitations

Our study has several limitations. Due to the use of public records to collect data in this study, it is possible that the method of data collection did not capture all sustained injuries, providing an underestimate of the incidence of lower extremity injuries. We understand this may limit the validity of our results; however, we still believe this method captures the general trend in injuries between the 2 fields. Moreover, while we tried to control for confounding factors such as weather, age, and history of a prior injury, we are unable to account for personal factors such as player preference or league-specific factors that may influence the decision to proceed with surgery. To the best of our knowledge, this is the first study that has evaluated NFL injury differences in the past 3 years. We hope this information provides players, coaches, and teams with pertinent information to promote player safety.

### CONCLUSION

Our study showed that NFL athletes were more likely to get injured on artificial surfaces and that artificial turf dramatically increased the likelihood of sustaining a serious injury requiring surgery. These findings support prior research highlighting the necessity of putting player safety

first and reducing danger on pitches with artificial turf. For players, coaches, and teams to make decisions about playing surfaces, the study offers useful information regarding playing safety.

### REFERENCES

- Andersson H, Ekblom B, Krstrup P. Elite football on artificial turf versus natural grass: movement patterns, technical standards, and player impressions. *J Sports Sci.* 2008;26(2):113-122. doi:10.1080/02640410701422076
- Chavkin D. Aaron Rodgers jabs NFL regarding artificial turf, player safety. *Sports Illustrated.* Published November 15, 2022. Accessed July 16, 2023. <https://www.si.com/nfl/2022/11/15/green-bay-packers-aaron-rodgers-jabs-nfl-regarding-artificial-turf-player-safety>
- Gould HP, Lostetter SJ, Samuelson ER, Guyton GP. Lower extremity injury rates on artificial turf versus natural grass playing surfaces: a systematic review. *Am J Sports Med.* 2023;51(6):1615-1621. doi:10.1177/03635465211069562
- Hershman EB, Anderson R, Bergfeld JA, et al. An analysis of specific lower extremity injury rates on grass and FieldTurf playing surfaces in National Football League games: 2000-2009 seasons. *Am J Sports Med.* 2012;40(10):2200-2205. doi:10.1177/0363546512458888
- Kent R, Forman JL, Crandall J, Lessley D. The mechanical interactions between an American football cleat and playing surfaces in situ at loads and rates generated by elite athletes: a comparison of playing surfaces. *Sports Biomech.* 2015;14(1):1-17. doi:10.1080/14763141.2015.1024277
- Kent R, Forman JL, Lessley D, Crandall J. The mechanics of American football cleats on natural grass and infill-type artificial playing surfaces with loads relevant to elite athletes. *Sports Biomech.* 2015;14(2):246-257. doi:10.1080/14763141.2015.1052749
- Latest NFL injury data fuels grass vs. turf debate. *ESPN.com.* Published April 20, 2023. Accessed July 28, 2023. [https://www.espn.com/nfl/story/\\_/id/36243906/nflpa-new-injury-data-shows-grass-significantly-safer-turf](https://www.espn.com/nfl/story/_/id/36243906/nflpa-new-injury-data-shows-grass-significantly-safer-turf)
- Loughran GJ, Vulpis CT, Murphy JP, et al. Incidence of knee injuries on artificial turf versus natural grass in National Collegiate Athletic Association American Football: 2004-2005 through 2013-2014 seasons. *Am J Sports Med.* 2019;47(6):1294-1301. doi:10.1177/0363546519833925
- Mack CD, Hershman EB, Anderson RB, et al. Higher rates of lower extremity injury on synthetic turf compared with natural turf among National Football League athletes: epidemiologic confirmation of a biomechanical hypothesis. *Am J Sports Med.* 2019;47(1):189-196. doi:10.1177/0363546518808499
- Meyers MC. Incidence, mechanisms, and severity of game-related college football injuries on FieldTurf versus natural grass: a 3-year prospective study. *Am J Sports Med.* 2010;38(4):687-697. doi:10.1177/0363546509352464
- Meyers MC. Incidence, mechanisms, and severity of match-related collegiate men's soccer injuries on FieldTurf and natural grass surfaces: a 6-year prospective study. *Am J Sports Med.* 2017;45(3):708-718. doi:10.1177/0363546516671715
- Meyers MC. Incidence, mechanisms, and severity of match-related collegiate women's soccer injuries on FieldTurf and natural grass surfaces: a 5-year prospective study. *Am J Sports Med.* 2013;41(10):2409-2420. doi:10.1177/0363546513498994
- Meyers MC, Barnhill BS. Incidence, causes, and severity of high school football injuries on FieldTurf versus natural grass: a 5-year prospective study. *Am J Sports Med.* 2004;32(7):1626-1638. doi:10.1177/0363546504266978
- Motamedi AR, Gowd AK, Nazemi AK, Gardner ST, Behrend CJ. Incidence, positional distribution, severity, and time missed in medial collateral ligament injuries of the knee in NCAA Division I football athletes. *J Am Acad Orthop Surg Glob Res Rev.* 2017;1(5):e019. doi:10.5435/JAAOSGlobal-D-17-00019

15. NFL data shows injury rates same on grass, turf. *ESPN.com*. Published November 9, 2022. Accessed July 26, 2023. [https://www.espn.com/nfl/story/\\_/id/34982032/nfl-data-shows-recent-injury-rates-same-grass-artificial-turf](https://www.espn.com/nfl/story/_/id/34982032/nfl-data-shows-recent-injury-rates-same-grass-artificial-turf)
16. NFL on ESPN—scores, stats and highlights. *ESPN.com*. Accessed July 28, 2023. <https://www.espn.com/nfl/>
17. NFL.com. Accessed July 28, 2023. <https://www.nfl.com/>
18. Nicholas JA. A historical perspective of injuries in professional football: twenty-six years of game-related events. *JAMA*. 1988;260(7):939. doi:10.1001/jama.1988.03410070067030
19. NRG Astrodome website. Published December 8, 2016. Accessed June 30, 2023. <https://www.nrgpark.com/nrg-park-facilities-2/nrg-astrodome/>
20. Only natural grass can level the NFL's playing field. NFL Players Association. Accessed July 16, 2023. <https://nflpa.com/posts/only-natural-grass-can-level-the-nfls-playing-field>
21. Orchard JW, Powell JW. Risk of knee and ankle sprains under various weather conditions in American football. *Med Sci Sports Exerc*. 2003;35(7):1118-1123. doi:10.1249/01.MSS.0000074563.61975.9B
22. Paliobeis A, Sivasundaram L, Knapik DM, et al. Injury incidence is higher on artificial turf compared with natural grass in high school athletes: a retrospective cohort study. *Curr Orthop Pract*. 2021;32(4):355. doi:10.1097/BCO.0000000000001012
23. Powell JW, Schootman M. A multivariate risk analysis of selected playing surfaces in the National Football League: 1980 to 1989: an epidemiologic study of knee injuries. *Am J Sports Med*. 1992;20(6):686-694. doi:10.1177/036354659202000609
24. Williams S, Hume PA, Kara S. A review of football injuries on third and fourth generation artificial turfs compared with natural turf. *Sports Med*. 2011;41(11):903-923. doi:10.2165/11593190-000000000-00000