

9-24-2013

## Scavenger receptor class B type I regulates cellular cholesterol metabolism and cell signaling associated with breast cancer development.

Christiane Danilo  
*Thomas Jefferson University*

Jorge L Gutierrez-Pajares  
*Thomas Jefferson University*

Maria Antonietta Mainieri  
*Thomas Jefferson University*

Follow this and additional works at: [https://jdc.jefferson.edu/stem\\_regenerativefp](https://jdc.jefferson.edu/stem_regenerativefp)

Isabelle Mercier

Part of the [Medical Biochemistry Commons](#), and the [Medical Molecular Biology Commons](#)

**Let us know how access to this document benefits you**

Michael P. Lisanti

*Thomas Jefferson University*

### Recommended Citation

Danilo, Christiane; Gutierrez-Pajares, Jorge L; Mainieri, Maria Antonietta; Mercier, Isabelle; Lisanti, Michael P.; and Frank, Philippe G., "Scavenger receptor class B type I regulates cellular cholesterol metabolism and cell signaling associated with breast cancer development." (2013). *Department of Stem Cell Biology and Regenerative Medicine Faculty Papers & Presentations*. Paper 16.

[https://jdc.jefferson.edu/stem\\_regenerativefp/16](https://jdc.jefferson.edu/stem_regenerativefp/16)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Stem Cell Biology and Regenerative Medicine Faculty Papers & Presentations by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

---

## Authors

Christiane Danilo, Jorge L. Gutierrez-Pajares, Maria Antonietta Mainieri, Isabelle Mercier, Michael P. Lisanti, and Philippe G. Frank

# Scavenger receptor class B type I regulates cellular cholesterol metabolism and cell signaling associated with breast cancer development

Christiane Danilo<sup>1</sup>  
Email: christianed@gmail.com

Jorge L Gutierrez-Pajares<sup>1</sup>  
Email: jlgutierrez2005@gmail.com

Maria Antonietta Mainieri<sup>1</sup>  
Email: ma.mainieri@gmail.com

Isabelle Mercier<sup>1</sup>  
Email: isabelle.mercier@jefferson.edu

Michael P Lisanti<sup>2</sup>  
Email: michaelp.lisanti@gmail.com

Philippe G Frank<sup>1,3,\*</sup>  
Email: Philippe.Frank@jefferson.edu

<sup>1</sup> Department of Stem Cell Biology and Regenerative Medicine, Kimmel Cancer Center, Thomas Jefferson University, Philadelphia, PA 19107, USA

<sup>2</sup> Manchester Breast Centre & Breakthrough Breast Cancer Research Unit; Paterson Institute for Cancer Research; Institute of Cancer Sciences; Manchester Academic Health Science Centre, University of Manchester, Manchester, UK

<sup>3</sup> Department of Biochemistry and Molecular Biology, Kimmel Cancer Center, Thomas Jefferson University, Philadelphia, PA 19107, USA

\* Corresponding author. Department of Biochemistry and Molecular Biology, Kimmel Cancer Center, Thomas Jefferson University, Philadelphia, PA 19107, USA

## Abstract

### Introduction

Previous studies have identified cholesterol as an important regulator of breast cancer development. High-density lipoprotein (HDL) and its cellular receptor, scavenger receptor class B type I (SR-BI) have both been implicated in the regulation of cellular cholesterol homeostasis, but their functions in cancer remain to be established.

## Methods

In the present study, we have examined the role of HDL and SR-BI in the regulation of cellular signaling pathways in breast cancer cell lines and in the development of tumor in a mouse xenograft model.

## Results

Our data show that HDL is capable of stimulating migration and can activate signal transduction pathways in the two human breast cancer cell lines, MDA-MB-231 and MCF7. Furthermore, we also show that knockdown of the HDL receptor, SR-BI, attenuates HDL-induced activation of the Phosphatidylinositol 3-kinase (PI3K)/Protein Kinase B (Akt) pathway in both cell lines. Additional investigations show that inhibition of the PI3K pathway, but not that of the mitogen-activated protein kinase (MAPK) pathway, could lead to a reduction in cellular proliferation in the absence of SR-BI. Importantly, while the knockdown of SR-BI led to decreased proliferation and migration *in vitro*, it also led to a significant reduction in tumor growth *in vivo*. Most importantly, we also show that pharmacological inhibition of SR-BI can attenuate signaling and lead to decreased cellular proliferation *in vitro*. Taken together, our data indicate that both cholesteryl ester entry via HDL-SR-BI and Akt signaling play an essential role in the regulation of cellular proliferation and migration, and, eventually, tumor growth.

## Conclusions

These results identify SR-BI as a potential target for the treatment of breast cancer.

## Introduction

Breast cancer is estimated to have affected over 200,000 women in the United States in 2012 and will be responsible for approximately 40,000 deaths, making it the second leading cause of cancer deaths [1]. Recently, attention has focused on examining the role of components of the diet, such as cholesterol, in breast cancer development. Plasma cholesterol levels are often lower in patients with advanced lung [2] and breast cancers [3-5], a phenomenon known as the “preclinical effect of cancer” [6]. This effect is believed to be consequential and not causal [6] but does suggest a role for cholesterol in cancer. Several epidemiologic studies have shown a positive correlation between elevated high density lipoprotein cholesterol (HDL-C) levels and breast cancer risk [5,7-9], and recent data have corroborated this claim [10-16]. *In vivo* works have suggested that hypercholesterolemia induced by diet and/or genetic background leads to increased tumor burden and metastasis in murine breast cancer models [10,12]. *In vitro* analyses have shown that human breast cancer cell lines exhibit increased proliferation and migration in the presence of HDL [11,13,15-17]. The effect of cholesterol on breast cancer may be attributed to several of its properties and functions. Cholesterol is the precursor of bioactive steroid hormones such as estrogen. It is also necessary for the formation of plasma membrane microdomains known as lipid rafts [18]. Lipid rafts are believed to organize signaling molecules in the plasma membrane and, as a result, have been implicated in the development of human cancers [19]. Therefore, cholesterol may play an essential role in the regulation of tumor growth [20,21].

The HDL lipoprotein is an important carrier of plasma cholesterol and can function as a signaling molecule by initiating MAPK and AKT signaling pathways and stimulate migration in endothelial cells [22-24]. The activation of these signaling pathways is dependent on HDL binding to the HDL receptor, the scavenger receptor class B, type I (SR-BI) and subsequent lipid transfer to the cell [25-27]. SR-BI functions as the HDL receptor and has been shown to mediate the selective transfer of cholesteryl ester from HDL molecules to cells in a process known as the selective HDL-cholesteryl ester uptake [28]. Its role in the development of atherosclerosis has been well documented [28], but its role in cancer has not been extensively investigated. Nevertheless, SR-BI has been implicated in prostate [29] and breast cancer [15,30]. In the case of breast cancer, SR-BI protein levels were found to be increased in malignant tissue samples compared to the normal surrounding tissue [30].

In the present study, we have examined the role of HDL and SR-BI in the regulation of cellular signaling pathways in breast cancer cell lines and in the development of tumors in a mouse xenograft model. Our data show that HDL can stimulate migration and can activate signal transduction pathways in the two human breast cancer cell lines, MDA-MB-231 and MCF7. Furthermore, we also show that knockdown of the HDL receptor, SR-BI, attenuates HDL-induced activation of the MAPK and PI3K/Akt pathways in both cells lines. A more detailed analysis revealed that SR-BI regulates signaling pathways via Akt activation, and the regulation of SR-BI expression or activity could limit tumor development in a mouse model.

## **Methods**

### **Materials**

The following antibodies were used: SR-BI was from Novus Biologicals, Inc. (Littleton, CO). CD31 antibody was from Abcam, Inc. (Cambridge, MA). Phospho-Erk1/2 (T202/Y204), Erk1/2, Phospho-Akt (S473), and Akt were from Cell Signaling Technology, Inc. (Beverly, MA). GAPDH was from Fitzgerald Industries International (Acton, MA), and  $\beta$ -Actin was from Sigma-Aldrich Corp. (St. Louis, MO). Anti-mouse secondary antibody was from Thermo Fisher Scientific, Inc. (Rockford, IL) and anti-rabbit secondary antibody was from BD Biosciences (San Jose, CA). The signaling inhibitors U0126 and LY294002 were from Cell Signaling Technology and Sigma-Aldrich, respectively. BLT-1 was from EMD Millipore (Billerica, MA).

### **Cell culture**

MCF7 cells were obtained from the American Tissue Culture Collection (ATCC) (Manassas, VA) and MDA-MB-231 cells were as previously described [31]. MDA-MB-231 and MCF7 cells were grown in Dulbecco's modified Eagle's media (DMEM) containing 10% Fetal Bovine Serum (FBS) in an incubator kept at 37°C with 5% CO<sub>2</sub>.

### **Purification of lipoproteins**

Human plasma was obtained from adult female volunteers. Approval for the use of human plasma was obtained from the Office of Human Research at Thomas Jefferson University, and consent was obtained from the volunteers for the use of their plasma samples. Lipoproteins [LDL (1.019-1.063 g/mL), HDL<sub>2</sub> (1.063-1.125 g/mL), HDL<sub>3</sub> (1.125-1.21 g/mL)] were separated by density gradient ultracentrifugation as previously described [32]. HDL<sub>3</sub>

was dialyzed against PBS, 0.5 mM EDTA, 1% NaN<sub>3</sub>. In experiments with MCF7 cells, lipoproteins were stripped of estrogens with activated charcoal to remove endogenous steroid hormones, as previously described [16].

### **Migration and invasion assays**

Migration was assayed using modified, non-coated Boyden chambers from BD Biosciences (San Jose, CA). Both MDA-MB-231 and MCF7 cells were cultured in 10-cm dishes in DMEM containing 10% FBS. Cells were harvested, counted, and washed three times with PBS. Cells were resuspended in DMEM containing 1% BSA, and  $5.0 \times 10^4$  cells were added to the upper chambers. The bottom chambers contained 1% FBS or 100 µg/mL lipoprotein in 1% BSA. Cells were incubated at 37°C overnight. At the end of the experiment, upper chambers were swabbed with a cotton swab to remove non-migrating cells, stained with crystal violet and quantified using phase contrast microscopy. Migration was quantified by counting the number of cells in 5 separate fields at 10× magnification. Invasion assays were performed as described above for transwell migration assays but used matrigel-coated Boyden chambers (BD Biosciences). The upper chambers contained cells in 1% BSA and the bottom chambers contained 1% FBS or 100 µg/mL lipoprotein in 1% BSA.

### **Immunoblot analysis**

Cells were lysed in radio-immunoprecipitation assay (RIPA) buffer containing protease and phosphatase inhibitors from Roche Applied Science (Indianapolis, IN) and Sigma-Aldrich Corp., respectively, prepared and analyzed as previously described [33]. Briefly, proteins were separated by SDS-PAGE and transferred to a nitrocellulose membrane. Membranes were blocked in 5% BSA in Tris-buffered saline containing 0.1% Tween-20 (TBS-T). Primary antibodies were diluted in 5% BSA in TBS-T and incubated for either 1 hour at room temperature or overnight at 4°C. Membranes were washed three times in TBS-T, and incubated with horseradish peroxidase-conjugated secondary antibody for 1 hour at room temperature. Membranes were washed three times in TBS-T, and visualized using Pierce Chemiluminescent Substrate from Thermo Fisher Scientific, Inc.

### **Knockdown of SR-BI**

MDA-MB-231 and MCF7 stably transfected cell lines were produced by lentiviral transduction. Lentiviral particles containing a pool of 3 short hairpin RNA (shRNA) constructs targeted against SR-BI were purchased from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA). Control lentiviral particles containing scrambled shRNA sequences were also purchased from Santa Cruz. Cells were transduced with the lentiviral particles according to manufacturer's protocol. Two days post-transduction, target cells containing either control shRNA (shCTL) or shRNA against SR-BI (shSRBI) were selected using 2.5 µg/mL puromycin. Successful knockdown was verified by immunoblot analysis.

### **Cholesterol content determination**

Cells were grown to confluence in 10-cm dishes in the presence of complete media (DMEM, 10% FBS). In half of the plates cholesterol was extracted with isopropanol. The extract was dried down in glass tubes under nitrogen and re-solubilized in a smaller volume of isopropanol. Cholesterol content was determined using the Cholesterol E kit from Wako

Chemicals USA, Inc. (Richmond, VA) as per the manufacturer's instructions. In the other plates, cells were lysed with 0.5 NaOH, and lysates were collected. Protein concentration was determined by the bicinchoninic acid assay (BCA) from Thermo Fisher Scientific as per the manufacturer's instruction. Total cellular cholesterol levels were calculated by dividing the total cholesterol by the total protein per dish.

### **<sup>3</sup>H-Thymidine incorporation proliferation assays**

Cell proliferation was measured by [<sup>3</sup>H]-thymidine incorporation to assess DNA synthesis and proliferation. MDA-MB-231 cells were seeded ( $5.0 \times 10^4$  cells/well) in 12-well plates in 1 mL of DMEM containing 10% fetal bovine serum (FBS) and grown overnight. The following day, media was aspirated, cells were washed twice with PBS and were serum-starved with 1% fatty acid free bovine serum albumin (BSA) for 1 hour. Media containing FBS, 1% BSA, or 1% BSA with lipoproteins (100 µg/mL) and 1 µCi/mL of <sup>3</sup>H-thymidine from Perkin Elmer (Waltham, MA) was added to cells. Cells were incubated for 6 hours at which time media was removed, cells were washed twice with PBS and incubated in 10% trichloroacetic acid to precipitate DNA. Cells were solubilized in 0.1 M NaOH and 1% SDS. Radioactivity was measured by liquid scintillation counting. Protein concentration was determined using the BCA assay.

### **Tumor studies**

All mice were housed and maintained in a barrier facility at the Kimmel Cancer Center at Thomas Jefferson University. Mice used in this study were athymic nude mice obtained from Taconic (Hudson, NY). Animal protocols used for these studies were approved by the Institutional Animal Care and Use Committee of Thomas Jefferson University. MDA-MB-231 ( $10^6$ ) cells containing either shRNA targeted against SR-BI (shSRBI) or control shRNA containing scrambled shRNA (shCTL) were subcutaneously injected in the flank of 7–9 week old nude mice. MCF7 cells ( $5 \times 10^6$ ) were orthotopically injected into the mammary fat pad of 9 week-old athymic nude mice implanted with slow release 17β-estradiol pellets (0.36mg/pellet, 60 days) from Innovative Research of America (Sarasota, FL). Four weeks post-injection, tumors were excised, weighed, and the volume was determined using the formula ( $\text{width}^2 \times \text{length}$ )/2. Half of each tumor was flash frozen and stored at –80°C and subsequently homogenized and lysed in RIPA buffer for immunoblot analysis as previously described [33]. The other half was fixed in formalin for 24 hours and then used to prepare paraffin-embedded sections.

### **Immunohistochemical analysis**

Paraffin-embedded tumor sections were deparaffinized in xylene and rehydrated. Antigen retrieval was performed in 10 mM citrate buffer pH6 for 10 min using a pressure cooker. Endogenous peroxidase activity was blocked with 3% H<sub>2</sub>O<sub>2</sub>, and sections were blocked in 10% goat serum obtained from Vector Laboratories, Inc. (Servion, Switzerland) and incubated with primary antibody overnight at 4°C. Sections were washed three times with PBS, incubated with biotinylated secondary antibody for 30 minutes followed by HRP-conjugated streptavidin for 30 minutes using a Streptavidin-HRP kit from Dako North America, Inc. (Carpinteria, CA). After 3 washes in PBS, the presence of bound antibody was visualized using 3,3' diaminobenzidine (DAB). Slides were counterstained with hematoxylin, dehydrated, and mounted with coverslips.

## **TUNEL assay**

Apoptosis was measured by TUNEL assay using the TUNEL-based ApopTag Peroxidase In Situ Apoptosis Detection Kit from Millipore (Temecula, CA) as per manufacturer's instructions. Briefly, paraffin-embedded tumor sections were de-paraffinized and rehydrated. Sections were treated with 20 µg/ml proteinase K from Roche Applied Science (Indianapolis, IN) for 15 min at room temperature and washed, and peroxidase activity was blocked by incubation in 3% hydrogen peroxide for 5 min. Sections were then incubated with equilibration buffer, followed by incubation in TdT enzyme for 1 h at 37°C. After washing, sections were incubated with HRP-conjugated antibody directed against digoxigenin for 30 min at RT, washed, and apoptotic positive cells were visualized using DAB. The percentage of apoptotic cells was quantified by dividing the number of TUNEL positive cells by the total number of cells observed in four distinct fields per section.

## **Statistical analyses**

All values are expressed as the mean  $\pm$  standard deviation (SD). The Prism 4.0 program (GraphPad Software, Inc., San Diego, CA) was used for statistical analysis. Statistical significance was examined using the Student's t test or ANOVA when appropriate.

## **Results**

### **HDL<sub>3</sub> stimulates migration and activates Akt and Erk1/2 in MCF7 and MDA-MB-231 cells**

Previous studies have shown that HDL can induce migration of endothelial cells [24]. In cancer, tumor cell migration represents the initial step associated with the development of metastasis [34]. To examine the effect of HDL on breast cancer cell migration, we studied the effect of lipoproteins on the migration of two breast cancer cell lines, MCF7 and MDA-MB-231. Interestingly, we found that when HDL<sub>3</sub> was used as the chemoattractant, it induced migration of both MCF7 and MDA-MB-231 cells by 3.5 and 61 fold, respectively, compared to the controls (CTL) as a chemoattractant (Figure 1A and B). Interestingly, LDL had no effect on the migration of neither MCF7 nor MDA-MB-231 cells (Figure 1A and B). Since lipoproteins, specifically HDL, can act as signaling molecules in endothelial cells and prostate cancer cells and activate Akt and MAPK pathways [23,35], we examined their effect on signaling in MCF7 and MDA-MB-231 cells. However, HDL<sub>3</sub> (100µg/mL) stimulated the activation of Erk1/2 and Akt in both MCF7 and MDA-MB-231 cells (Figure 1C). A modest increase in the phosphorylation of Erk1/2 was observed in MDA-MB-231 cells after 30 minutes of incubation with HDL<sub>3</sub>. However, a more robust and quicker response was observed in MCF7 cells (Figure 1C). Additionally, HDL<sub>3</sub> rapidly activated Akt in both cell lines, an effect that was prolonged in MCF7 cells (Figure 1C). These results indicate that HDL<sub>3</sub> can function as a signaling molecule in these two breast cancer cell lines. LDL had a modest effect on Akt activation, and no effect on Erk1/2 activation in neither MDA-MB-231 nor MCF-7 cells was observed (Figure 1D).



**Figure 1 HDL<sub>3</sub> induces migration and acts as a signaling molecule in MDA-MB-231 and MCF7 cells.** **A.** and **B.** Migration of MCF7 (A.) and MDA-MB-231 (B.) cells was induced by HDL<sub>3</sub>. Migration of MCF7 and MDA-MB-231 cells was increased when HDL<sub>3</sub>, but not LDL, was used as a chemoattractant compared to CTL (1% BSA alone) in Boyden Chamber assays. Stained cells were quantified using phase contrast microscopy. Columns represent the mean number of migrated cells; bars represent  $\pm$  standard deviation (SD) (n=3). The results obtained with CTL and HDL<sub>3</sub> groups are significantly different. There was no statistical difference between CTL and LDL groups. (A. \* $P$  < 0.05 Student's t test, B. \*\* $P$  < 0.01, versus 1% BSA by Student's t test). Results shown are representative of three independent experiments. **C.** MDA-MB-231 and MCF7 cells were incubated for the indicated amounts of time with 100  $\mu$ g/mL HDL<sub>3</sub>. Whole cell lysates were analyzed by Western blotting for the indicated proteins. The arrow indicates the protein corresponding to phospho-Akt. Results are representative of 3 independent experiments. **D.** MDA-MB-231 and MCF7 cells were incubated for the indicated amounts of time with 100  $\mu$ g/mL LDL. Whole cell lysates were analyzed by Western blotting for the indicated proteins. The arrow indicates the protein corresponding to phospho-Akt. Results are representative of 3 independent experiments.

### **Knockdown of the HDL receptor, SR-BI, attenuates the effects of HDL<sub>3</sub> on signaling in MDA-MB-231 and MCF7 cells**

In the following experiments, we examined the effect of downregulating the HDL receptor, SR-BI, on signaling in MDA-MB-231 and MCF7 cells. As demonstrated in Figure 2, we were able to successfully downregulate SR-BI in both MDA-MB-231 cells (Figure 2A) and MCF7 cells (Figure 2B). Knockdown of SR-BI was achieved by stable transduction of a pool of lentiviral particles containing shRNA sequences specific for SR-BI (shSRBI). shCTL cells were created by stable transduction of lentiviral particles containing a scrambled version of the shRNA. Knockdown of SR-BI was assessed by western blot analysis. In MDA-MB-231 cells, SR-BI expression was reduced by 5.3 fold and in MCF7 cells, SR-BI expression was reduced by 4.0 fold (Figure 2A). To determine the role of SR-BI on the regulation of signaling pathways, both shCTL and shSRBI MDA-MB-231 and MCF7 cells were serum starved overnight and then incubated in media containing 10% FBS for 30 minutes or 100  $\mu$ g/mL of HDL<sub>3</sub> for 0, 5, 15 and 30 minutes, as indicated. We found that the activation of Akt was greatly reduced in the shSRBI cells compared to the shRNA control cells. Similar results were obtained with both MDA-MB-231 and MCF7 cell lines in the presence of FBS (lanes 1 and 6 in both Figures 2A and 2B). Consistent with the results presented in Figure 1C, HDL<sub>3</sub> was able to stimulate the activation of Akt in both cell lines in a time-dependent manner. However, activation of Akt in shSRBI MDA-MB-231 cells was greatly reduced when stimulated by HDL<sub>3</sub> for 15 and 30 minutes (Figure 2A, lanes 9 and 10), compared to the Akt activation observed in shCTL MDA-MB-231 cells when stimulated by HDL<sub>3</sub> for the same periods of time (Figure 2A, lanes 4 and 5). Similar results were obtained in MCF7 cells. In that case, Akt activation was reduced in the shSRBI MCF7 cells when stimulated by HDL<sub>3</sub> for 15 and 30 minutes (Figure 2B, lanes 9 and 10), compared to shCTL MCF7 cells stimulated by HDL<sub>3</sub> for the same periods of time (Figure 2B, lanes 4 and 5). Finally, Erk1/2 appeared to be constitutively active in MDA-MB-231 cells (Figure 2A, lanes 1–10). However, almost no change in Erk1/2 activation was detected in shSRBI MDA-MB-231 cells treated with HDL<sub>3</sub> for 30 minutes (Figure 2A, lane 10) compared to shCTL MDA-MB-231 treated with HDL<sub>3</sub> for 30 minutes (Figure 2A, lane 5). This effect was in contrast with observations made with MCF7 cells (Figure 2B). In shCTL MCF7 cells, HDL<sub>3</sub> rapidly stimulated Erk1/2 activation, reaching a peak at 5 minutes (Figure 2B, lane 3) but maintaining a sustained effect at 30 minutes (Figure 2B, lanes 4 and 5). Activation of Erk1/2

in shSRBI MCF7 cells followed a similar pattern, but the intensity of activation was greatly reduced (Figure 2B, lanes 7–10). These results suggest that downregulation of SR-BI in MDA-MB-231 and MCF7 cells attenuates signaling via the AKT and MAPK pathways. Additionally, our results also show that the interaction between HDL and SR-BI regulates activation of these signaling pathways. Finally, the effect of LDL was also tested in these cells lines. Results presented in Figure 2C and D demonstrate that the downregulation of SR-BI in MDA-MB231 and MCF7 cells had no effect on the regulation of Akt and Erk1/2 activation by LDL.

---

**Figure 2 Knockdown of SR-BI attenuates HDL<sub>3</sub>-induced signal transduction in MDA-MB-231 and MCF7 cells.** A, B. MDA-MB-231 and MCF7 cells were stably transduced by lentiviruses carrying either shRNA against SR-BI (shSRBI) or control shRNA (shCTL). Cells were incubated with either 10% FBS for 30 minutes or with 100 µg/mL HDL<sub>3</sub> for the indicated times. Whole cell lysates were analyzed by Western blot for the indicated proteins. The arrow indicates the protein corresponding to phospho-Akt. GAPDH was used as loading control. C, D. MDA-MB-231 and MCF7 cells were stably transduced by lentiviruses carrying either shRNA against SR-BI (shSRBI) or control shRNA (shCTL). Cells were incubated with either 10% FBS for 30 minutes or with 100 µg/mL LDL for the indicated times. Whole cell lysates were analyzed by Western blot for the indicated proteins. The arrow indicates the protein corresponding to phospho-Akt. GAPDH was used as loading control. Note that, in panel C, image was spliced (between line 5 and 6) to eliminate an irrelevant line.

---

### **Knockdown of the HDL receptor, SR-BI, inhibits proliferation and migration of MDA-MB-231 cells**

We observed decreased signaling in shSRBI MDA-MB-231 cells compared to shCTL MDA-MB-231 cells in the presence of FBS. Therefore, we used media containing 10% FBS for the remainder of the experiments described below. FBS contains large amounts of lipoproteins and provides suitable ligands for SR-BI. Previous studies have shown that expression of SR-BI in COS-7 cells significantly increased cholesterol mass in these cells compared to control vector-transfected cells [36]. To determine if cellular cholesterol homeostasis was affected by the downregulation of SR-BI, we quantified the cholesterol content of shCTL and shSRBI MDA-MB-231 cells. Under basal conditions, shCTL MDA-MB-231 cells contained a significantly greater (1.3-fold increase) ( $P<0.01$ ) total cholesterol content compared to shSRBI MDA-MB-231 cells (Figure 3A).

---

**Figure 3 Knockdown of the HDL receptor, SR-BI, inhibits proliferation, migration, and invasion of MDA-MB-231 cells.** **A.** Cholesterol content of shCTL and shSRBI MDA-MB-231 cells. Cholesterol was extracted from cells with isopropanol, and total cholesterol was measured enzymatically using a colorimetric assay. Protein concentration was also determined, and total cellular cholesterol content was expressed as the amount of cholesterol per milligram of protein. Columns represent the mean cholesterol content and bars represent  $\pm$  SD. shCTL MDA-MB-231 cells displays significantly more cholesterol than shSRBI MDA-MB-231 cells (\*\* $P < 0.01$  by student's t test). **B.** Knockdown of SR-BI inhibits proliferation of MDA-MB-231 cells. MDA-MB-231 cells containing control shRNA (shCTL) or shSRBI were incubated in the presence of  $^3\text{H}$ -thymidine for 6 hours in DMEM containing 10% FBS. Proliferation was measured by  $^3\text{H}$ -thymidine incorporation. Columns represent the mean  $^3\text{H}$ -Thymidine incorporation (cpm/ $\mu\text{g}$  protein); bars represent  $\pm$  (SD). The results obtained for shCTL and shSRBI cells are significantly different (\* $P < 0.01$  by student's t test). Results are representative of three independent experiments. **C.** Knockdown of SR-BI reduces cellular migration. Cellular migration of shCTL and shSRBI MDA-MB-231 cells was assessed for 24h in a Boyden Chamber using 1% FBS as a chemoattractant. Columns represent the mean number of migrated cells from three independent experiments, bars represent  $\pm$  standard deviation (SD), (n=12). The results obtained from the shCTL and shSRBI groups are significantly different (\* $P < 0.05$  versus shCTL, as determined by student's t test). **D.** Knockdown of SR-BI leads to a marginally significant reduction in cellular invasion. Invasion of shCTL and shSRBI MDA-MB-231 cells was assessed by matrigel-coated Boyden Chamber Assays using 1% FBS as a chemoattractant. Columns represent the mean number of invaded cells from three separate experiments (n=12); bars represent  $\pm$  SD ( $P = 0.0517$ , Student's t test).

---

Previous studies have shown that a mutant of SR-BI inhibits proliferation of the luminal B subtype of human breast cancer cells, MCF7, in the presence of HDL [30]. To further investigate the role of SR-BI in a triple negative (lacking the estrogen receptor (ER), progesterone receptor (PR), and Her2) basal B subtype breast cancer cell line, we determined the effect of knocking down SR-BI on the proliferation of MDA-MB-231 cells. The proliferation of shSRBI MDA-MB-231 cell was reduced by 2.0 fold compared to the proliferation observed with shCTL MDA-MB-231 cells ( $P < 0.01$ ) (Figure 3B). Knockdown of SR-BI also significantly reduced cellular migration ( $P < 0.05$ ) by 1.65 fold (Figure 3C). Finally, a reduction of SR-BI protein levels was associated with a marginally significant reduction of the ability of MDA-MB-231 cells to invade (Figure 3D) ( $P = 0.0517$ ).

### **Pharmacological inhibition of SR-BI reduces proliferation and signal transduction in MDA-MB-231 cells**

To further elucidate the inhibitory effect of SR-BI ablation on proliferation and signal transduction obtained by molecular biological means, we treated cells with the pharmacological inhibitor of SR-BI, BLT-1 [37]. Previous work has identified BLT-1 as a specific inhibitor of SR-BI function. BLT-1 has been shown to act by blocking the transfer of lipids from HDL particles to cells [37]. The  $\text{IC}_{50}$  for this compound was determined to be 50 nM [37]. The ability of BLT-1 to regulate proliferation was evaluated in the presence of varying concentrations of this inhibitor. Accordingly, BLT-1 could inhibit growth of shCTL MDA-MB-231 cells in a dose-dependent manner (Figure 4A). At doses over 50 nM, BLT-1 treatment could significantly decrease proliferation of shCTL MDA-MB-231 cells compared to the control untreated cells ( $P < 0.001$  for concentrations of 50 nM, 75 nM, and 100 nM compared to shCTL MDA-MB-231 incubated with 0 nM BLT-1). Importantly, there was no

significant effect of BLT-1 treatment on the proliferation of shSRBI MDA-MB-231 cells: Proliferation of shSRBI cells treated with concentrations of BLT-1 between 0.1nM and 100nM were not statistically different from that of vehicle-treated shSRBI cells. Interestingly, there was also no significant difference between shCTL MDA-MB-231 cells treated with BLT-1 at doses above 20nM and vehicle-treated (0nM) shSRBI MDA-MB-231 cells. Taken together, these results suggest that down regulation or pharmacological inhibition of SR-BI have similar effect on MDA-MB-231 proliferation.

---

**Figure 4 Inhibition of SR-BI by the small molecule BLT-1 inhibits proliferation of MDA-MB-231 cells.** **A.** Pharmacological inhibition of SR-BI in MDA-MB-231 cells inhibits cellular proliferation. MDA-MB-231 shCTL cells were incubated in the presence of the indicated doses of BLT-1 and <sup>3</sup>H-thymidine added to the culture media for six hours. At this time, the assay was stopped, and lysates were collected. Columns represent the mean <sup>3</sup>H-thymidine incorporation (cpm/μg protein); bars represent ±SD. Results obtained with shSRBI MDA-MB-231 CTL (0 nM BLT-1) are significantly different from those obtained with shCTL 0 nM BLT-1 (CTL) (*a*, *P*<0.001), shCTL 0.1 nM BLT-1 (*b*, *P*<0.001), and shCTL 1.0 nM (*c*, *P*<0.05). Results obtained with shCTL MDA-MB-231 CTL (0nM BLT-1) were significantly different from those obtained with shCTL 50 nM BLT-1 (*d*, *P*<0.001), shCTL 75 nM BLT-1 (*e*, *P*<0.001), and shCTL 100 nM (*f*, *P*<0.001). Significance was determined by one-way ANOVA with Tukey's post-test analysis. **B.** Pharmacological inhibition of SR-BI attenuates signaling in MDA-MB-231 cells. Western blot analysis was performed with extracts obtained from shSRBI MDA-MB-231 cells treated with vehicle, shCTL MDA-MB-231 cells treated with vehicle, and shCTL MDA-MB-231 cells treated with BLT-1 (50 nM). Cells were serum starved overnight, incubated with vehicle alone or BLT-1 for 3 hours, and then treated with 10% FBS for 30 minutes.

---

We also examined the effect of BLT-1 on signal transduction in these cells. In agreement with the finding described in Figure 2A, Akt activation in shSRBI MDA-MB-231 cells treated with FBS for 30 minutes was reduced compared to shCTL MDA-MB-231 cells (Figure 4B). Similar results were obtained with shCTL MDA-MB-231 cells with treated BLT-1. Akt activation was reduced in the treated shCTL MDA-MB-231 cells compared to untreated control cells. Finally, SR-BI knockdown or pharmacological inhibition had no effect on Erk1/2 activation (Figure 4B) compared to the control cells. A statistical difference between untreated shSRBI MDA-MB-231 cells and shSRBI MDA-MB-231 cells treated with 100 nM BLT-1 was also detected. This observation may be due to the presence of some residual SR-BI protein. Collectively, these data suggest that Akt activation may be mediated, in part, by SR-BI, and the downregulation of SR-BI is responsible for the observed reduction in the cellular proliferation.

### **Inhibition of PI3K, not MEK1/2, inhibits growth of shCTL MDA-MB-231 cells**

In order to elucidate the mechanism by which SR-BI knockdown inhibits proliferation, we used pharmacological agents to inhibit PI3K and MAPK signaling pathways. We show that the PI3K inhibitor, LY294002, abolished FBS-induced activation of Akt (Figure 5A) in shCTL and shSRBI MDA-MB-231 cells. Importantly, PI3K inhibition significantly reduced proliferation of shCTL MDA-MB-231 cells to levels similar to those observed with untreated shSRBI MDA-MB-231 cells (Figure 5B). In addition, PI3K inhibition had no effect on the proliferation of shSRBI MDA-MB-231 cells suggesting that downregulation of SR-BI in these cells was sufficient to inhibit proliferation. Conversely, U0126-induced inhibition of

MEK1/2, which activates Erk1/2, did not affect proliferation of shCTL MDA-MB-231 or shSRBI MDA-MB-231 cells (Figure 5B). Erk1/2 activation, however, was significantly reduced by inhibition of MEK1/2 in both cell types (Figure 5A). These results suggest that the MAPK pathway does not play a significant role in SR-BI-mediated signaling and proliferation unlike the PI3K pathway.

---

**Figure 5 Inhibition of PI3K, not MEK1/2, prevents proliferation of MDA-MB-231 cells.**

**A.** LY294002 and U0126 effectively inhibit Akt and Erk1/2 activation in MDA-MB-231 cells. Serum-starved shCTL and shSRBI MDA-MB-231 cells were incubated with or without the inhibitors LY294002 (15  $\mu$ M) or U0126 (10  $\mu$ M) for 2 hours. Medium containing 10% FBS was added for 30 minutes, cells were lysed, and whole cell lysates were analyzed by Western blot for the indicated proteins. GAPDH was used as a loading control. **B.** PI3K inhibition reduces cellular proliferation of shCTL MDA-MB-231 cells. shCTL and shSRBI MDA-MB-231 cells were incubated with culture media containing either LY294002 (15 $\mu$ M) or U0126 (10  $\mu$ M) and  $^3$ H-thymidine for 6 hours, at which time the assay was stopped, and lysates were collected. Columns represent the mean  $^3$ H-thymidine incorporation (cpm/ $\mu$ g protein); bars represent  $\pm$  SD. Results obtained from vehicle-treated (CTL) shCTL MDA-MB-231 cells are significantly different from those obtained with shSRBI MDA-MB-231 cells (\*\* $P$ <0.001, one-way ANOVA with Tukey's post-test analysis). Results obtained from vehicle-treated shCTL MDA-MB-231 cells (CTL) are significantly different from those obtained with shCTL MDA-MB-231 cells treated with LY294002. Note that proliferation of shCTL MDA-MB-231 cells treated with U0126 was not significantly different from that observed with vehicle-treated shCTL MDA-MB-231 cells. Proliferation of shSRBI MDA-MB-231 cells treated with LY294002 or U0126 was also not significantly different from that observed with vehicle-treated shCTL MDA-MB-231 cells. Results are representative of three independent experiments.

---

**Knockdown of SR-BI results in decreases in *in vivo* tumor growth of MDA-MB-231 and MCF7 cells**

To assess the effect of SR-BI knockdown *in vivo*, we subcutaneously injected shSRBI and shCTL MDA-MB-231 cells in the flanks of nude mice. Four weeks after injection, tumors were excised from sacrificed mice, and mass and volume were measured. Tumors obtained with shCTL MDA-MB-231 were significantly larger than those obtained from shSRBI MDA-MB-231: tumor volume and mass were increased by 3.8 fold and 3.7 fold, respectively (Figure 6A). To determine the role of SR-BI in tumor growth in MCF7 cells, shCTL and shSRBI MCF7 cells ( $5 \times 10^6$ ) were orthotopically injected into the mammary fat pad of athymic nude mice after implantation with slow release 17 $\beta$ -estradiol pellets. Four weeks post-injection, mice were sacrificed, tumors were excised, and mass and volume of tumors were determined (Figure 6B). Tumors obtained with shCTL MCF7 were 1.5-fold larger than those obtained with shSRBI MCF7 ( $P$ <0.05), and 1.3-fold larger by mass (marginally significant).

---

**Figure 6 Knockdown of SR-BI inhibits xenograft tumor growth *in vivo*.** **A.** Knockdown of SR-BI in MDA-MB-231 cells decreases tumor growth *in vivo*. Athymic nude mice were injected with  $10^6$  MDA-MB-231 cells carrying control shRNA (shCTL) or shSRBI. Four weeks post-injection, mice were sacrificed, and tumors were excised. Tumors were weighed and volume determined using the formula  $(\text{length} \times \text{width}^2)/2$ . Columns represent the mean volume and mass respectively, bars represent  $\pm$  SD. Tumors obtained with shCTL cells were significantly larger by weight and volume than the tumors obtained with shSRBI. (\* $P < 0.05$ ,  $n = 14$  shCTL,  $n = 13$  shSRBI) **B.** Knockdown of SR-BI in MCF7 cells decreases tumor growth *in vivo*. MCF7 cells ( $5 \times 10^6$ ) were orthotopically injected into the mammary fat pad of 9 week-old athymic nude mice implanted with slow release  $17\beta$ -estradiol pellets (0.36mg/pellet, 60 days). Four weeks post injection, mice were sacrificed, and the tumors were excised. Tumors were weighed and volume determined using the formula  $(\text{length} \times \text{width}^2)/2$ . Columns represent the mean volume and mass respectively; bars represent  $\pm$  SD. The shCTL tumors were significantly larger by volume than the shSRBI tumors (\* $P < 0.05$ ,  $n = 8$  shCTL,  $n = 7$  shSRBI). **C.** shCTL tumors display increased expression of proliferative markers compared to shSRBI tumors. Immunohistochemistry was performed to determine expression patterns of SR-BI and pErk1/2 in MDA-MB231 tumors. **D.** shCTL tumors display increased expression of proliferative markers compared to shSRBI tumors. Three tumors per cell type were analyzed by western blot analysis for the indicated proteins. GAPDH was used as a loading control.

---

To elucidate the mechanism by which SR-BI regulates tumor formation, tissue immunohistochemical analyses (Figure 6C) and immunoblot analyses of homogenized tumors (Figure 6D) were performed. Immunohistochemistry analyses demonstrated the reduction in SR-BI protein expression in shSRBI MDA-MB-231-derived tumors compared to shCTL MDA-MB231. Results also revealed that levels of the proliferative marker, pErk1/2, were decreased in shSRBI MDA-MB-231 tumors, compared to those observed in control tumors. Consistent with *in vitro* findings, pAkt levels were decreased in shSRBI MDA-MB-231 tumors compared to those observed in control tumors.

Since cholesterol has been shown to play a role in the regulation of angiogenesis [38], microvessel density in the tumors was assessed by staining tumor sections for CD31, a specific marker of endothelial cells (Figure 7A). A significant increase in microvessel density was observed with tumors obtained from shCTL MDA-MB-231 cells compared to those obtained from shSRBI MDA-MB-231 cells. These data suggest that SR-BI can regulate angiogenesis in these tumors. Finally, SR-BI has been shown to activate Akt [23], which may inhibit apoptosis thereby promoting cell survival [39]. Therefore, we assessed apoptosis by TUNEL staining in tissue sections obtained from shCTL and shSRBI MDA-MB-231 xenograft tumors (Figure 7B). As anticipated, we observed a significant increase in apoptosis in shSRBI MDA-MB-231 tumors compared to shCTL MDA-MB-231 tumors.

---

**Figure 7 Knockdown of SR-BI inhibits xenograft tumor growth *in vivo*.** **A.** Knockdown of SR-BI results in decreased angiogenesis in MDA-MB-231 xenograft tumors. Immunohistochemistry was performed to determine expression of CD31. CD31 staining was quantified by averaging the number of vessels per field in 5 separate fields per sample. Columns represent the mean microvessel density; bars represent  $\pm$  SD. Microvessel density was statistically greater in shCTL MDA-MB-231 tumors compared to shSRBI MDA-MB-231 cells ( $*P<0.05$  student's t test,  $n=4$  per group). **B.** Knockdown of SR-BI results in increased apoptosis in MDA-MB-231 xenograft tumors. Images shown are representative of TUNEL assay staining. Percentage of TUNEL positive cells was calculated by dividing the number of TUNEL-positive cells by the total number of cells in 4 separate fields. Columns represent the percent of TUNEL-positive cells; bars represent  $\pm$  SD. TUNEL staining is statistically increased in shSRBI MDA-MB-231 tumors compared to shCTL MDA-MB-231 tumors ( $***P<0.001$ , student's t test, shCTL  $n=3$  and shSRBI  $n=4$ ).

---

## Discussion

In the present study, we have examined the role of HDL and its receptor, SR-BI, in breast cancer development and progression. We found that HDL<sub>3</sub> stimulates migration and activates signaling pathways such as MAPK and PI3K in two breast cancer cell lines. Inhibiting selective HDL-cholesteryl ester uptake by knocking down or pharmacologically inhibiting SR-BI resulted in an attenuation of cell signaling events induced by HDL. Additionally, loss of SR-BI resulted in decreased proliferation, migration, and tumor growth of MDA-MB-231 cells. These findings suggest that regulating cholesterol metabolism and cellular signaling pathways via SR-BI may be linked and may additionally identify new targets associated with tumor progression.

### HDL, signal transduction and cellular migration

HDL has a well-established role in the etiology of atherosclerosis, particularly in reverse cholesterol transport whereby HDL removes excess cholesterol molecules from peripheral tissues and returns them to the liver for excretion or recycling [40]. In addition, HDL functions in a number of other cellular processes including inhibition of apoptosis in macrophages [41], induction of migration in endothelial cells [24], and the initiation of cell signaling events in multiple cells types [22,23,42]. Although clinical studies have suggested that plasma HDL levels may be correlated with increased breast cancer [5,7-9,12], the mechanisms by which HDL exerts its effect have yet to be elucidated. HDL has been shown to activate Erk1/2 in fibroblasts [23,42], Chinese hamster ovary cells [22], endothelial cells [23], and prostate cancer cells [35,43]. Studies have also shown that it can activate Akt in endothelial [23] and prostate cancer cells [35,43]. Interestingly, the activation of Erk1/2 [44] and Akt [39,45] has been implicated in several human cancers, including breast cancer. In the present study, we have established a role for HDL as a mediator of signal transduction in two breast cancer cell lines. Consistent with the results obtained in other cell types, we found that, in both MCF7 and MDA-MB-231 cells, incubation with HDL<sub>3</sub> induces a rapid activation of both Erk1/2 and Akt signaling pathways. These novel findings in breast cancer suggest that HDL may regulate various signaling pathways and may therefore alter tumor progression.

In the present study, we found that HDL can induce migration of two breast cancer cell lines, MCF7 and MDA-MB-231, suggesting that HDL may play a role in the early stages of metastasis. This finding is consistent with previous studies showing that HDL can stimulate

migration of endothelial cells [24,46]. Interestingly, the observed migration was shown to be mediated by SR-BI [24]. By contrast, a recent study reported that HDL inhibits migration of MDA-MB-231 in Boyden chamber assays [14]. However, in this study, the investigators used serum as the chemoattractant, and HDL was added to the upper chamber, thereby measuring the ability of HDL to prevent cellular migration induced by serum. By contrast, our method allows the analysis of the role of HDL in the regulation of cellular migration and therefore allows a direct measurement of the capability of HDL to induce migration of MDA-MB-231 cells. As a result, our results indicate that HDL may play a role in the pathogenesis of breast cancer, especially in the later stages.

## **SR-BI, signal transduction regulation, and tumor formation**

SR-BI has been implicated as a mediator of several cell signaling events in the context of atherosclerosis [22,23,25-27]. Previous studies have shown that HDL binding to SR-BI and subsequent lipid transfer are sufficient to activate Src, which subsequently activates the PI3K/Akt and MAPK pathways [25,27]. In endothelial cells, one of the downstream effectors of Akt is eNOS, which catalyzes the production of NO. The results obtained in the present study are consistent with the hypothesis that SR-BI may also play a role in signal transduction in the context of cancer. In agreement with this hypothesis, upon knockdown or pharmacological inhibition of SR-BI in MDA-MB-231 cells, Akt activation was significantly reduced, suggesting that SR-BI may be mediating this response. In addition, down-regulation of SR-BI was accompanied by a reduction of total cholesterol levels in MDA-MB-231 cells. These results are consistent with reports that indicate that the cholesterol flux mediated by SR-BI plays a role in the regulation of signal transduction initiation [26]. In our model, decreased total cholesterol levels may represent a reduction in SR-BI-mediated cholesterol flux and therefore significantly reduce signal transduction activation. SR-BI also binds LDL, which can, like HDL, promote the cellular entry of cholesteryl ester. While LDL may promote the entry of cholesteryl ester via SR-BI, it is not sufficient to induce migration of breast cancer cells, and it does not appear to alter Akt activation (Figure 2C and D). Taken together, our data suggest that both cholesteryl ester entry via HDL-SR-BI and Akt activation are required for cellular proliferation and migration, and, eventually, tumor growth.

Activation of the PI3K/Akt pathway promotes growth, survival, and proliferation [45] and has been implicated in a variety of human cancers [39]. Importantly, Akt is aberrantly hyper-activated in approximately 40% of breast cancers [39]. Interestingly, we observed a reduction in proliferation and migration in the SR-BI-knockdown cells compared to control cells in association with reduced Akt activation. These results suggest that SR-BI may mediate the activation of Akt and its downstream effects in the presence of HDL. Mechanistically, we showed that the inhibition of the PI3K/Akt pathway results in significantly reduced proliferation of shCTL MDA-MB-231 cells similar to the reduction in proliferation observed in shSRBI MDA-MB-231 cells. Importantly, no further reduction in proliferation of shSRBI MDA-MB-231 cells was detected upon inhibition of the PI3K/Akt pathway. Taken together, these data suggest that reduced Akt activation observed in the shSRBI MDA-MB-231 cells may be responsible for reduced proliferation of these cells compared to shCTL MDA-MB-231 cells.

Previous studies have suggested a role for SR-BI in the etiology of breast cancer. Cao et al. [30] have shown that expression of SR-BI is increased in human breast tumors compared to the normal surrounding tissue. They also demonstrated that recombinant expression of a mutant form of SR-BI, which lacked the carboxyl-terminal tail of the protein, could inhibit



proliferation of breast cancer cells. Their study further suggested that this effect was possibly due to reduced Akt activation. Our study is the first to demonstrate directly that Akt activation is reduced when SR-BI is knocked down or pharmacologically inhibited. Furthermore, in agreement with the previously mentioned study [30], we showed that proliferation of MDA-MB-231 cells was significantly inhibited by downregulation of SR-BI protein levels and by pharmacological inhibition of SR-BI. In addition, we observed that SR-BI knockdown inhibits migration. This finding may suggest a role for SR-BI in the initiation of metastasis. Finally, we demonstrated that knockdown of SR-BI in MDA-MB-231 cells can lead to reduced tumor growth *in vivo* accompanied by increased activation of Erk1/2 and Akt, and an increase in cellular apoptosis. In MCF7 cells, knockdown of SR-BI also led to reduced xenograft tumor growth.

Previous studies have shown that Akt can inhibit apoptosis through a variety of mechanisms including the phosphorylation of BAD, thus preventing cytochrome c release from mitochondria and the direct inhibition of the caspase activation cascade [39]. Increased cellular cholesterol levels have been shown to increase Akt activation and decrease apoptosis in prostate cancer cells [47]. Consistent with these observations, a recent study has shown that inhibition of xenograft tumor growth could be achieved with colon cancer cells that re-express the ATP-binding cassette transporter A1 (ABCA1) [48]. ABCA1 is a lipid transporter that mediates the efflux of cellular cholesterol to lipid free apolipoprotein A-I [49]. Furthermore, in this study, re-expression of ABCA1 resulted in decreased mitochondrial cholesterol content and increased release of cytochrome c, which ultimately led to increased apoptosis. Our work also indicated that SR-BI knockdown can significantly reduce apoptosis in xenograft tumors as shown by TUNEL staining. Consequently, a reduction in the levels of cellular cholesterol content may be responsible, at least in part, for the decreased apoptosis observed in our model. Taken together, these data also suggest an important role for cholesterol in the regulation of cellular signaling pathways and tumor formation. Importantly, excess cellular cholesterol accumulates in the form of esterified cholesterol. Previous works and ours suggest that the accumulation of esterified cholesterol may lead to a modification of signaling pathways associated with proliferation and migration in tumors. Consistent with this hypothesis, increasing cellular esterified cholesterol levels have been shown to induce cellular proliferation and enhance invasiveness of tumor cell lines [50]. Conversely, the inhibition of cholesterol esterification has been shown to have the reverse effect [51,52].

## Conclusions

In summary, our results suggest that HDL and SR-BI have pro-oncogenic activity and can induce migration and activate signal transduction pathways responsible for cellular proliferation and tumor formation in two breast cancer cell lines. Additionally, knockdown or pharmacological inhibition of SR-BI could attenuate signaling mediated by HDL and inhibit proliferation, migration, and tumor growth. Taken together, these findings identify SR-BI and HDL as potential therapeutic targets for the treatment of breast cancer.

## Abbreviations

ABCA1, ATP binding cassette A1; Akt, Protein Kinase B; BLT1, Blocks lipid transport-1; BSA, Bovine serum albumin; eNOS, endothelial nitric oxide synthase; ER, Estrogen receptor; Erk, Extracellular-signaling related kinase; FBS, fetal bovine serum; HDL, High density lipoprotein; HDL-C, High density lipoprotein-cholesterol; LDL, Low density lipoprotein; PI3K, Phosphatidylinositol 3-kinase; SR-BI, Scavenger receptor Class B, Type I; TN, Triple negative.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

PGF supervised all aspects of this research and the preparation of this manuscript. CD, MPL, IM, and PGF participated in the design of the study. CD, JLGP, IM, and MAM carried out the experimental data acquisition. CD, JLGP, MPL, MAM, and PGF performed data analyses. All authors read, critically revised, and approved the final manuscript.

## Acknowledgements

PGF was supported by grants from the Susan G. Komen Foundation and the National Institutes of Health (R01-CA-098779). This project was funded, in part, under a grant with the Pennsylvania Department of Health (to M.P.L.). The Department specifically disclaims responsibility for any analyses, interpretations or conclusions.

## References

1. Siegel R, Naishadham D, Jemal A: **Cancer statistics, 2012.** *CA Cancer J Clin* 2012, **62**:10–29.
2. Siemianowicz K, Gminski J, Stajszczyk M, Wojakowski W, Goss M, Machalski M, Telega A, Brulinski K, Magiera-Molendowska H: **Serum HDL cholesterol concentration in patients with squamous cell and small cell lung cancer.** *Int J Mol Med* 2000, **6**:307–311.
3. Potischman N, McCulloch CE, Byers T, Houghton L, Nemoto T, Graham S, Campbell TC: **Associations between breast cancer, plasma triglycerides, and cholesterol.** *Nutr Cancer* 1991, **15**:205–215.
4. Furberg AS, Veierod MB, Wilsgaard T, Bernstein L, Thune I: **Serum high-density lipoprotein cholesterol, metabolic profile, and breast cancer risk.** *J Natl Cancer Inst* 2004, **96**:1152–1160.
5. Ray G, Husain SA: **Role of lipids, lipoproteins and vitamins in women with breast cancer.** *Clin Biochem* 2001, **34**:71–76.

6. Kritchevsky SB, Kritchevsky D: **Serum cholesterol and cancer risk: an epidemiologic perspective.** *Annu Rev Nutr* 1992, **12**:391–416.
7. Boyd NF, McGuire V: **Evidence of association between plasma high-density lipoprotein cholesterol and risk factors for breast cancer.** *J Natl Cancer Inst* 1990, **82**:460–468.
8. Ferraroni M, Gerber M, Decarli A, Richardson S, Marubini E, Crastes de Paulet P, Crastes de Paulet A, Pujol H: **HDL-cholesterol and breast cancer: a joint study in northern Italy and southern France.** *Int J Epidemiol* 1993, **22**:772–780.
9. Kitahara CM, Berrington de Gonzalez A, Freedman ND, Huxley R, Mok Y, Jee SH, Samet JM: **Total cholesterol and cancer risk in a large prospective study in Korea.** *J Clin Oncol* 2011, **29**:1592–1598.
10. Alikhani N, Ferguson RD, Novosyadlyy R, Gallagher EJ, Scheinman EJ, Yakar S, Leroith D: **Mammary tumor growth and pulmonary metastasis are enhanced in a hyperlipidemic mouse model.** *Oncogene* 2012, **8**:961–967.
11. Jozan S, Faye JC, Tournier JF, Tauber JP, David JF, Bayard F: **Interaction of estradiol and high density lipoproteins on proliferation of the human breast cancer cell line MCF-7 adapted to grow in serum free conditions.** *Biochem Biophys Res Commun* 1985, **133**:105–112.
12. Llaverias G, Danilo C, Mercier I, Daumer K, Capozza F, Williams TM, Sotgia F, Lisanti MP, Frank PG: **Role of cholesterol in the development and progression of breast cancer.** *Am J Pathol* 2011, **178**:402–412.
13. Pan B, Ren H, Lv X, Zhao Y, Yu B, He Y, Ma Y, Niu C, Kong J, Yu F, *et al*: **Hypochlorite-induced oxidative stress elevates the capability of HDL in promoting breast cancer metastasis.** *J Transl Med* 2012, **10**:65.
14. Pan B, Ren H, Ma Y, Liu D, Yu B, Ji L, Pan L, Li J, Yang L, Lv X, *et al*: **High-density lipoprotein of patients with type 2 diabetes mellitus elevates the capability of promoting migration and invasion of breast cancer cells.** *Int J Cancer* 2011, **131**:70–82.
15. Pussinen PJ, Karten B, Wintersperger A, Reicher H, McLean M, Malle E, Sattler W: **The human breast carcinoma cell line HBL-100 acquires exogenous cholesterol from high-density lipoprotein via CLA-1 (CD-36 and LIMPII analogous 1)-mediated selective cholesteryl ester uptake.** *Biochem J* 2000, **349**:559–566.
16. Rotheneder M, Kostner GM: **Effects of low- and high-density lipoproteins on the proliferation of human breast cancer cells in vitro: differences between hormone-dependent and hormone-independent cell lines.** *Int J Cancer* 1989, **43**:875–879.
17. Pan B, Ren H, He Y, Lv X, Ma Y, Li J, Huang L, Yu B, Kong J, Niu C, *et al*: **HDL of patients with type 2 diabetes mellitus elevates the capability of promoting breast cancer metastasis.** *Clin Cancer Res* 2012, **18**:1246–1256.

18. Simons K, Sampaio JL: **Membrane organization and lipid rafts.** *Cold Spring Harb Perspect Biol* 2011, **3**:a004697.
19. Staubach S, Hanisch FG: **Lipid rafts: signaling and sorting platforms of cells and their roles in cancer.** *Expert Rev Proteomics* 2011, **8**:263–277.
20. Coleman PS, Chen L-C, Sepp-Lorenzino L: **Cholesterol metabolism and tumor cell proliferation.** In *Subcellular Biochemistry. Volume 28*. Edited by Bittman R. New York: Plenum Press; 1997:363–435.
21. Danilo C, Frank PG: **Cholesterol and breast cancer development.** *Current opinion in pharmacology* 2012, **12**:677–682.
22. Grewal T, de Diego I, Kirchhoff MF, Tebar F, Heeren J, Rinninger F, Enrich C: **High density lipoprotein-induced signaling of the MAPK pathway involves scavenger receptor type BI-mediated activation of Ras.** *The Journal of biological chemistry* 2003, **278**:16478–16481.
23. Mineo C, Yuhanna IS, Quon MJ, Shaul PW: **High density lipoprotein-induced endothelial nitric-oxide synthase activation is mediated by Akt and MAP kinases.** *The Journal of biological chemistry* 2003, **278**:9142–9149.
24. Seetharam D, Mineo C, Gormley AK, Gibson LL, Vongpatanasin W, Chambliss KL, Hahner LD, Cummings ML, Kitchens RL, Marcel YL, *et al*: **High-density lipoprotein promotes endothelial cell migration and reendothelialization via scavenger receptor-B type I.** *Circ Res* 2006, **98**:63–72.
25. Al-Jarallah A, Trigatti BL: **A role for the scavenger receptor, class B type I in high density lipoprotein dependent activation of cellular signaling pathways.** *Biochim Biophys Acta* 2010, **1801**:1239–1248.
26. Assanasen C, Mineo C, Seetharam D, Yuhanna IS, Marcel YL, Connelly MA, Williams DL, de la Llera-Moya M, Shaul PW, Silver DL: **Cholesterol binding, efflux, and a PDZ-interacting domain of scavenger receptor-BI mediate HDL-initiated signaling.** *J Clin Invest* 2005, **115**:969–977.
27. Saddar S, Mineo C, Shaul PW: **Signaling by the high-affinity HDL receptor scavenger receptor B type I.** *Arterioscler Thromb Vasc Biol* 2009, **30**:144–150.
28. Krieger M: **Charting the fate of the “good cholesterol”: identification and characterization of the high-density lipoprotein receptor SR-BI.** *Annu Rev Biochem* 1999, **68**:523–558.
29. Twiddy AL, Cox ME, Wasan KM: **Knockdown of scavenger receptor class B type I reduces prostate specific antigen secretion and viability of prostate cancer cells.** *Prostate* 2012, **72**:955–965.
30. Cao WM, Murao K, Imachi H, Yu X, Abe H, Yamauchi A, Niimi M, Miyauchi A, Wong NC, Ishida T: **A mutant high-density lipoprotein receptor inhibits proliferation of human breast cancer cells.** *Cancer Res* 2004, **64**:1515–1521.

31. Bonuccelli G, Tsigos A, Whitaker-Menezes D, Pavlides S, Pestell RG, Chiavarina B, Frank PG, Flomenberg N, Howell A, Martinez-Outschoorn UE, *et al*: **Ketones and lactate “fuel” tumor growth and metastasis: Evidence that epithelial cancer cells use oxidative mitochondrial metabolism.** *Cell Cycle* 2010, **9**:3506–3514.
32. Redgrave TG, Roberts DC, West CE: **Separation of plasma lipoproteins by density-gradient ultracentrifugation.** *Anal Biochem* 1975, **65**:42–49.
33. Williams TM, Sotgia F, Lee H, Hassan G, Di Vizio D, Bonuccelli G, Capozza F, Mercier I, Rui H, Pestell RG, *et al*: **Stromal and epithelial caveolin-1 both confer a protective effect against mammary hyperplasia and tumorigenesis: Caveolin-1 antagonizes cyclin D1 function in mammary epithelial cells.** *Am J Pathol* 2006, **169**:1784–1801.
34. Brabletz T, Jung A, Spaderna S, Hlubek F, Kirchner T: **Opinion: migrating cancer stem cells - an integrated concept of malignant tumour progression.** *Nat Rev Cancer* 2005, **5**:744–749.
35. Sekine Y, Suzuki K, Remaley AT: **HDL and sphingosine-1-phosphate activate stat3 in prostate cancer DU145 cells via ERK1/2 and S1P receptors, and promote cell migration and invasion.** *Prostate* 2011, **71**:690–699.
36. Kellner-Weibel G, de La Llera-Moya M, Connelly MA, Stoudt G, Christian AE, Haynes MP, Williams DL, Rothblat GH: **Expression of scavenger receptor BI in COS-7 cells alters cholesterol content and distribution.** *Biochemistry* 2000, **39**:221–229.
37. Nieland TJ, Penman M, Dori L, Krieger M, Kirchhausen T: **Discovery of chemical inhibitors of the selective transfer of lipids mediated by the HDL receptor SR-BI.** *Proc Natl Acad Sci U S A* 2002, **99**:15422–15427.
38. Solomon KR, Pelton K, Boucher K, Joo J, Tully C, Zurakowski D, Schaffner CP, Kim J, Freeman MR: **Ezetimibe is an inhibitor of tumor angiogenesis.** *Am J Pathol* 2009, **174**:1017–1026.
39. Bellacosa A, Kumar CC, Di Cristofano A, Testa JR: **Activation of AKT kinases in cancer: implications for therapeutic targeting.** *Adv Cancer Res* 2005, **94**:29–86.
40. Lewis GF, Rader DJ: **New insights into the regulation of HDL metabolism and reverse cholesterol transport.** *Circ Res* 2005, **96**:1221–1232.
41. Terasaka N, Wang N, Yvan-Charvet L, Tall AR: **High-density lipoprotein protects macrophages from oxidized low-density lipoprotein-induced apoptosis by promoting efflux of 7-ketocholesterol via ABCG1.** *Proc Natl Acad Sci U S A* 2007, **104**:15093–15098.
42. Deeg MA, Bowen RF, Oram JF, Bierman EL: **High density lipoproteins stimulate mitogen-activated protein kinases in human skin fibroblasts.** *Arterioscler Thromb Vasc Biol* 1997, **17**:1667–1674.
43. Sekine Y, Demosky SJ, Stonik JA, Furuya Y, Koike H, Suzuki K, Remaley AT: **High-density lipoprotein induces proliferation and migration of human prostate androgen-**

**independent cancer cells by an ABCA1-dependent mechanism.** *Mol Cancer Res* 2010, **8**:1284–1294.

44. Whyte J, Bergin O, Bianchi A, McNally S, Martin F: **Mitogen-activated protein kinase signalling in experimental models of breast cancer progression and in mammary gland development.** *Breast Cancer Res* 2009, **11**:209.

45. Vivanco I, Sawyers CL: **The phosphatidylinositol 3-Kinase AKT pathway in human cancer.** *Nat Rev Cancer* 2002, **2**:489–501.

46. Murugesan G, Sa G, Fox PL: **High-density lipoprotein stimulates endothelial cell movement by a mechanism distinct from basic fibroblast growth factor.** *Circ Res* 1994, **74**:1149–1156.

47. Zhuang L, Kim J, Adam RM, Solomon KR, Freeman MR: **Cholesterol targeting alters lipid raft composition and cell survival in prostate cancer cells and xenografts.** *J Clin Invest* 2005, **115**:959–968.

48. Smith B, Land H: **Anticancer Activity of the Cholesterol Exporter ABCA1 Gene.** *Cell Rep* 2012, **2**:580–590.

49. Zannis VI, Chroni A, Krieger M: **Role of apoA-I, ABCA1, LCAT, and SR-BI in the biogenesis of HDL.** *J Mol Med* 2006, **84**:276–294.

50. Paillasse MR, de Medina P, Amouroux G, Mhamdi L, Poirot M, Silvente-Poirot S: **Signaling through cholesterol esterification: a new pathway for the cholecystokinin 2 receptor involved in cell growth and invasion.** *J Lipid Res* 2009, **50**:2203–2211.

51. de Medina P, Boubekour N, Balaguer P, Favre G, Silvente-Poirot S, Poirot M: **The prototypical inhibitor of cholesterol esterification, Sah 58–035 [3-[decyldimethylsilyl]-n-[2-(4-methylphenyl)-1-phenylethyl]propanamide], is an agonist of estrogen receptors.** *The Journal of pharmacology and experimental therapeutics* 2006, **319**:139–149.

52. de Medina P, Genovese S, Paillasse MR, Mazaheri M, Caze-Subra S, Bystricky K, Curini M, Silvente-Poirot S, Epifano F, Poirot M: **Auraptene is an inhibitor of cholesterol esterification and a modulator of estrogen receptors.** *Molecular pharmacology* 2010, **78**:827–836.















Figure 1





**A****Total Cholesterol Content****B****<sup>3</sup>H-Thymidine Incorporation****C****Migration****D****Invasion**




**A**

Figure 4

**B**

**A**

LY294002    -    -    +    +    -    -

U0125    -    -    -    -    +    +


**B**

Figure 5






Figure 6


**A.**

MDA-MB-231  
shCTL

MDA-MB-231  
shSRBI





CD-31



**B.**

MDA-MB-231  
shCTL

MDA-MB-231  
shSRBI



TUNEL




Figure 7