

8-1-2008

Infant mortality in Delaware: the need to improve maternal health and reduce premature births.

David A. Paul, MD

Thomas Jefferson University, DPaul@Christianacare.org

Let us know how access to this document benefits you

Follow this and additional works at: <https://jdc.jefferson.edu/pedsfp> Part of the [Bioethics and Medical Ethics Commons](#), and the [Pediatrics Commons](#)

Recommended Citation

Paul, MD, David A., "Infant mortality in Delaware: the need to improve maternal health and reduce premature births." (2008). *Department of Pediatrics Faculty Papers*. Paper 17.<https://jdc.jefferson.edu/pedsfp/17>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Pediatrics Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

As submitted to:

Delaware medical journal

and later published as:

Delaware medical journal

Volume 80, Issue 8, August 2008, Pages 289-292

PubMed ID: 18795726

**“Infant Mortality in Delaware: The Need to Improve Maternal Health
and Reduce Premature Births”**

David A. Paul, MD

Chair, Delaware Healthy Mother Infant Consortium

Department of Pediatrics, Section of Neonatology, Christiana Care Health Services

Attending Neonatologist Christiana Care Health Services, Attending Neonatologist,

duPont Hospital for Children

Associate Professor of Pediatrics, Thomas Jefferson University

Chair, Delaware Healthy Mother Infant Consortium

Correspondence:

David A. Paul, MD

MAP-1, Suite 217

4745 Ogletown-Stanton Road

Newark, DE 19713

302 733 2410

Infant mortality remains a difficult problem in Delaware. Delaware's infant mortality rate is substantially higher than the overall United States rate, with a recent 16% increase in infant mortality noted from 1995-1997 to 2002-2004¹. During this same time period the United States rate decreased by 7%. Delaware has one of the highest infant mortality rates in the country, with rates similar to rural southern states such as Mississippi and Louisiana, rather than rates comparable to our mid-Atlantic bordering states. This editorial will review some of the potential causes of Delaware's high infant mortality rate as well as some potential solutions.

What do we know about causes of infant mortality in Delaware?

The leading cause of infant mortality in the United States is controversial. Traditionally, birth defects have been considered the leading cause of infant mortality. However, recent research has suggested that premature birth is likely the leading cause². The United States is presently experiencing an epidemic of premature birth with a 20% increase in the number of preterm babies since 1990³. Delaware has a rate of premature birth of 14% compared to 12.7% nationally⁴. The causes of premature birth are multifactorial and include biologic, social, medical and racial factors⁵. There are tremendous racial disparities in rates of premature birth. African American women have a rate of premature birth of 18%, compared to 11.5% in Caucasian women¹. From a medical standpoint, little can be done to stop advanced premature labor. Administration of weekly doses of 17 hydroxy- progesterone has been shown to reduce the risk of preterm birth in women who have had a previous premature infant⁶.

Although many advances have been made in the care of premature infants, recent evidence suggests that there have been little advancement in survival rates of those infants born very low birth weight (<1500 grams)^{7,8}. Despite advances in perinatal care, including the administration of antenatal corticosteroids, antenatal antibiotics, high frequency ventilation, and postnatal surfactant, infants born extremely prematurely remain at high risk of death and/or surviving with a disability⁹. Corresponding to the national increase in premature birth, the number of very low birth weight infants (VLBW, <1500 grams) cared for in the Neonatal Intensive Care Unit at Christiana Hospital increased 55% between 1994 and 2006. This increase in the number of premature infants, at a time corresponding to stagnation of outcomes, has led to an increased number of deaths of premature infants.

In addition to an increased number of premature births, we have recently demonstrated an increase in illness severity in VLBW infants in Delaware¹⁰. The observed increase in illness severity was accompanied by a decrease in stillbirths, suggesting a shift from stillbirths to live births of extremely ill premature infants. In support of this hypothesis, shifts from stillbirth to live premature births have been shown in other populations¹¹, and stillbirths have been shown to be decreasing nationally¹.

Similar to the observed disparities in preterm births, racial disparities in infant mortality rates exist both nationally and in Delaware. Infants born to African American women in Delaware are more than twice as likely to die prior to their first birthday compared to

infants born to Caucasian mothers or Hispanic mothers. The reasons behind this elevated disparity ratio remain elusive but are likely multifactorial. One recent focus of investigation into the causes of infant mortality in Delaware has been racial disparities in delivery of health care. African American mothers delivering VLBW infants are less likely to receive antenatal corticosteroids, tocolytic medications and Cesarean deliveries compared to Caucasian mothers ¹². In other regions of the country, racial disparities in delivery of neonatal intensive care have been associated with increased black infant mortality ¹³.

Delaware may be considered at “high risk” for having an increased infant mortality rate based on the racial demographics of the state. As African American infants have a risk of infant mortality twice that of Caucasian infants, states with a higher percentage of African American population would intuitively have high infant mortality rates. The demographic make-up of Delaware is 20.7% African American compared to 12.4 % nationally ¹⁴. However, our recent analysis indicates that the infant mortality rate in Delaware remains high after adjusting for the population demographics of the state ¹⁵. Furthermore, in this analysis adolescent pregnancy and smoking during pregnancy were identified as factors which are potentially amenable to intervention and could be targeted to decrease the risk of infant mortality ¹⁵.

Social Factors Contributing to Infant Mortality

Infant mortality is as much a social issue as a medical issue. In order to reduce infant mortality in Delaware we will have to make important social gains. With 12% of Delaware residents lacking health insurance, access to basic health care is limited to a large percentage of our population ¹⁶. Employment remains the principle source of health care insurance in our country. Disappointingly, the high school graduation rate in Delaware is only 73% ¹⁶. With education levels linked to employment opportunities, the number of uninsured Delawareans is likely to continue to climb.

Family support is a critical component of child-rearing. Traditionally, research surrounding premature birth and infant mortality have centered on maternal factors. The role of paternal factors and family support networks in Delaware need closer investigation. Fathers play an important role in family economic viability and support. Fathers also play an important role in maternal health through transmission of diseases such as HIV and other sexually transmitted diseases. Births to unmarried mothers have been climbing steadily on a national level ¹. In Delaware, 44.3% of live births were to unmarried mothers in 2005, compared to 36.9% nationally ⁴.

Improved access to health care and strengthened family structures are only two examples of social issues that interface with infant mortality. Neighborhood safety, housing, transportation and racism are examples of other social problems which have been linked to infant mortality in other regions and need to be improved in Delaware.

Preconception health care: a paradigm shift

In order to decrease premature births in Delaware, and subsequent infant mortality, we need to vastly improve the health of women of child-bearing age. This includes better control of conditions such as obesity, diabetes, hypertension and smoking, all of which have been linked to premature birth. In Delaware, 22% of the population smokes and 26% are classified as obese¹⁶, evidence of the poor overall health of the state population. In April, 2006 the CDC published guidelines to improve preconception health in women of childbearing age¹⁷. Preconception health care is not a single visit but a process of care. Preconception care aims to keep a mother healthy prior to pregnancy, not waiting for prenatal care to begin the process of risk reduction. Under the paradigm of preconception health care, any health care encounter with women of childbearing age should be considered an opportunity to improve preconception health.

Many new initiatives to improve preconception health have been funded as part of State of Delaware infant mortality programs. Christiana Care Health Services, St. Francis Hospital, Westside Clinic, La Red, Delmarva Rural Ministries and the University of Delaware all have recently started preconception health programs. These programs, if successful, should improve the future health of women of childbearing age in our state. In addition, Delaware is one of the few states in the nation to have made 17 hydroxy-progesterone available to low-income women at risk for recurrent premature birth.

Summary

Delaware's high infant mortality rate continues to be driven by an increasing number of premature births. Premature birth is a complex problem with both social and medical roots. Based on the CDC road map, programs aimed at improving preconception health of women of childbearing age are blossoming in Delaware. In addition, Delaware's infant mortality rate can only be reduced if we are able to diminish disparities in health care delivery, and improve the social factors influencing infant mortality. These factors include diminishing family support, low high school graduation rate, poor housing, and a lack of health insurance.

REFERENCES

1. Martin JA, Kung HC, Mathews TJ, et al. Annual summary of vital statistics: 2006. *Pediatrics*. 2008;121:788-801.
2. Callaghan WM, MacDorman MF, Rasmussen SA, Qin C, Lackritz EM. The contribution of preterm birth to infant mortality rates in the United States. *Pediatrics*. 2006;118:1566-1573.
3. Hoyert DL, Mathews TJ, Menacker F, Strobino DM, Guyer B. Annual summary of vital statistics: 2004. *Pediatrics*. 2006;117:168-83.
4. Martin JA, Hamilton BE, Sutton PD, et al. Births: Final data for 2005. *National Vital Statistics Reports*. 2007;56:1-103.
5. *Preterm Birth Causes, Consequences and Prevention*. 1st ed. Washington, D.C.: The National Academies Press; 2007.
6. Meis PJ, Society for Maternal-Fetal M. 17 hydroxyprogesterone for the prevention of preterm delivery. *Obstetrics & Gynecology*. 2005;105:1128-35.
7. Horbar JD, Badger GJ, Carpenter JH, et al. Trends in mortality and morbidity for very low birth weight infants, 1991-1999. *Pediatrics*. 2002;110:143-51.

8. Meadow W, Lee G, Lin K, Lantos J. Changes in mortality for extremely low birth weight infants in the 1990s: Implications for treatment decisions and resource use. *Pediatrics*. 2004;113:1223-9.
9. Marlow N, Wolke D, Bracewell MA, Samara M, EPICure Study G. Neurologic and developmental disability at six years of age after extremely preterm birth. *N Engl J Med*. 2005;352:9-19.
10. Paul DA, Leef KH, Locke RG, Bartoshesky L, Walrath J, Stefano JL. Increasing illness severity in very low birth weight infants over a 9-year period. *BMC Pediatrics*. 2006;6:2.
11. Basso O, Rasmussen S, Weinberg CR, Wilcox AJ, Irgens LM, Skjaerven R. Trends in fetal and infant survival following preeclampsia. *JAMA*. 2006;296:1357-62.
12. Paul DA, Locke R, Zook K, Leef KH, Stefano JL, Colmorgen G. Racial differences in prenatal care of mothers delivering very low birth weight infants. *Journal of Perinatology*. 2006;26:74-8.
13. Howell EA, Hebert P, Chatterjee S, Kleinman LC, Chassin MR. Black/white differences in very low birth weight neonatal mortality rates among New York City hospitals. *Pediatrics*. 2008;121:e407-15.
14. United states census bureau. Available at:
<http://factfinder.census.gov/home/saff/main.html>. Accessed May 27, 2008.

15. Paul DA, Mackley A, Locke RG, Stefano JL, Kroelinger C. State infant mortality: An ecologic study to determine modifiable risks and adjusted infant mortality rates. *Matern Child Health J.* 2008: Epub ahead of print.

16. United health foundation. Available at:

www.unitedhealthfoundation.com/ahr2007/states/Delaware.html. Accessed May 28th, 2008.

17. Johnson K, Posner SF, Biermann J, et al. Recommendations to improve preconception health and health care--united states. A report of the CDC/ATSDR preconception care work group and the select panel on preconception care. *Morbidity & Mortality Weekly Report Recommendations & Reports.* 2006;55:1-23.