

4-1-2021

Association of Race With Lung Cancer Risk Among Adults Undergoing Lung Cancer Screening.


Christine S Shusted
Thomas Jefferson University

Nathaniel R Evans
Thomas Jefferson University

Hee-Soon Juon
Thomas Jefferson University

Gregory C Kane
Thomas Jefferson University

Julie A Barta
Follow this and additional works at: <https://jdc.jefferson.edu/pulmcritcarefp>
Thomas Jefferson University

 Part of the [Pulmonology Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Shusted, Christine S; Evans, Nathaniel R; Juon, Hee-Soon; Kane, Gregory C; and Barta, Julie A, "Association of Race With Lung Cancer Risk Among Adults Undergoing Lung Cancer Screening." (2021). *Division of Pulmonary and Critical Care Medicine Faculty Papers*. Paper 18.
<https://jdc.jefferson.edu/pulmcritcarefp/18>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Division of Pulmonary and Critical Care Medicine Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.



Research Letter | Oncology

Association of Race With Lung Cancer Risk Among Adults Undergoing Lung Cancer Screening

Christine S. Shusted, MPH; Nathaniel R. Evans, MD; Hee-Soon Juon, MSN, PhD; Gregory C. Kane, MD; Julie A. Barta, MD

Introduction

African American individuals at high risk of lung cancer may experience greater mortality benefit from annual lung cancer screening (LCS) compared with White individuals.^{1,2} However, African American individuals develop lung cancer with fewer pack-years of smoking and at younger ages than White individuals, and they are less often eligible for LCS when using United States Preventive Services Task Force (USPSTF) criteria.³ One approach to mitigate disparities in LCS eligibility is to use lung cancer risk prediction models to identify the patients with the highest risk.⁴ The Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial modified logistic regression model for lung cancer prediction (PLCOM2012) may improve sensitivity in lung cancer detection, specifically among African American individuals, and potentially reduce underrepresentation in screening cohorts.⁵ This study aimed to identify differences in PLCOM2012 6-year lung cancer risk among USPSTF-eligible LCS patients.

Author affiliations and article information are listed at the end of this article.

Methods

The Thomas Jefferson University institutional review board approved this cross-sectional study and granted a waiver of informed consent because of the minimal risk nature of the research. This study followed the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline.

The Jefferson LCS Program is a centralized program at an urban, academic medical center. Sociodemographic and clinical data from patients screened from January 2018 to September 2020 were extracted from the LCS Registry. This included lung cancer risk using the PLCOM2012 model, calculated at entry into LCS, and diagnosis of screen-detected lung cancer.⁶ After excluding patients who self-reported races other than African American or White, risk quartiles were generated using median PLCOM2012 risk values for the screening and cancer cohorts. Descriptive statistics, independent *t* tests, χ^2 tests, and Mann-Whitney *U* tests were performed using a *P* < .05 significance threshold. Statistical analyses were 2-sided and conducted using SPSS statistical software version 26 (IBM Corp) from October 5 to November 11, 2020.

Results

Among 1276 individuals in the screening cohort, the mean (SD) age was 64.25 (5.81) years, 545 (42.7%) were African American individuals, and 757 (59.3%) were female individuals (**Table**). There were significant differences in gender distribution (African American: 343 female individuals [62.9%] vs White: 414 female individuals [56.6%]; $\chi^2_1 = 5.137$; *P* = .02), smoking history (African American: 348 currently smoking [63.9%] vs White: 358 currently smoking [49.0%]; $\chi^2_1 = 27.967$; *P* < .001), education level (eg, African American: 243 high school graduates [44.6%] and 37 college graduates [6.8%] vs White: 287 high school graduates [39.3%] and 98 college graduates [13.4%]; $\chi^2_6 = 31.500$; *P* < .001), and insurance (eg, African American beneficiaries: 192 Medicaid or dual eligible [35.2%] and 138 private only [25.3%] vs White beneficiaries: 134 Medicaid or dual eligible [18.3%] and 243

Open Access. This is an open access article distributed under the terms of the CC-BY License.

Table. Baseline Characteristics of African American and White Patients Undergoing Lung Cancer Screening

Characteristics	Patients, No. (%)		P value
1276 Patients in full sample	African American (n = 545)	White (n = 731)	
Age, mean (SD), y	64.09 (5.70)	64.37 (5.89)	.40
Gender			
Female	343 (62.9)	414 (56.6)	.02
Male	202 (37.1)	317 (43.4)	
Ethnicity			
Hispanic/Latinx	1 (0.2)	48 (6.6)	<.001
Smoking status			
Current	348 (63.9)	358 (49.0)	<.001
Former	197 (36.1)	373 (51.0)	
Pack-years, mean (SD), y	48.71 (20.56)	57.95 (27.51)	<.001
Cigarettes, mean (SD)	21.62 (8.45)	25.89 (11.79)	<.001
Smoking duration, mean (SD), y	45.25 (7.15)	44.81 (8.18)	.304
Time since quitting, mean (SD), y	5.16 (4.97)	6.69 (5.45)	.001
Time smoked, mean (SD), y	45.25 (7.15)	44.81 (8.18)	.30
Personal history of cancer	83 (15.6)	161 (22.6)	.002
Family history of lung cancer	123 (22.6)	167 (22.8)	.91
COPD	241 (44.2)	292 (39.9)	.13
Body mass index, mean (SD) ^a	30.01 (7.27)	29.11 (6.42)	.02
Education			
Less than high school diploma	77 (14.1)	85 (11.6)	<.001
High school graduate	243 (44.6)	287 (39.3)	
Post high school training	19 (3.5)	28 (3.8)	
Some college	113 (20.7)	119 (16.3)	
College graduate	37 (6.8)	98 (13.4)	
Post graduate/professional degree	19 (3.5)	60 (8.2)	
Unknown	37 (6.8)	54 (7.4)	
Insurance status			
Medicare only	174 (31.9)	240 (32.8)	<.001
Medicaid/dual eligible	192 (35.2)	134 (18.3)	
Private only	138 (25.3)	243 (33.2)	
Medicare with supplement	26 (4.8)	69 (9.4)	
Private with supplement	10 (1.8)	24 (3.3)	
Other/none	5 (0.9)	21 (2.9)	
32 Patients with screen-detected lung cancer			
Characteristics	African American (n = 14)	White (n = 18)	
Age, mean (SD)	64.86 (5.60)	66.44 (5.87)	NA
Gender			
Female	12 (85.7)	11 (61.1)	NA
Male	2 (14.3)	7 (38.9)	NA
Histology			NA
Adenocarcinoma	11 (78.6)	10 (55.6)	NA
Squamous cell carcinoma	1 (7.1)	3 (16.7)	NA
Small cell carcinoma	0 (0.0)	3 (16.7)	NA
Neuroendocrine carcinoma	0 (0.0)	1 (5.6)	NA
Non-small cell carcinoma	2 (14.3)	1 (5.6)	NA
Stage			
I	10 (71.4)	9 (50.0)	NA
II	1 (7.1)	0 (0.0)	NA
III	2 (14.3)	4 (22.2)	NA
IV	1 (7.1)	2 (11.1)	NA
Limited stage small cell	0	1 (5.6)	NA
Extensive stage small cell	0	2 (11.1)	NA

Abbreviation: COPD, chronic obstructive pulmonary disease.

^a Body mass index is calculated as weight in kilograms divided by height in meters squared.

private only [33.2%]; $\chi^2_5 = 58.992$; $P < .001$) among races. Among the 32 patients with screen-detected lung cancer in the cancer cohort, 14 (44%) were African American.


In the screening cohort (median PLCom2012 risk of lung cancer: 4.75%; interquartile range [IQR], 2.63%-8.47%), African American individuals had a significantly higher median risk of lung cancer compared with White individuals (median PLCom2012 risk of lung cancer among African American individuals: 5.81%; IQR, 3.42%-9.79% vs White individuals: 4.10%; IQR, 2.14%-7.26%; $P < .001$) (Figure). When comparing among risk quartiles, 326 of African American individuals (59.8%) undergoing screening had a risk score in quartiles 3 and 4 compared with 311 White individuals (42.5%). However, in the cancer cohort (median PLCom2012 risk: 7.23%; IQR, 2.37%-12.64%), White individuals had a higher median risk compared with African Americans (median PLCom2012 risk of lung cancer among White individuals: 10.55%; IQR, 2.21%-16.16% vs African American individuals: 5.87%; IQR, 2.89%-7.88%; $P = .18$), but this result did not reach statistical significance. Moreover, 11 White individuals (61.1%) with screen-detected lung cancer had a risk score in quartiles 3 and 4, compared with just 5 African American individuals (35.7%).

Discussion

To our knowledge, this study is the first to measure PLCom2012 lung cancer risk among screened individuals and include more than 40% African American participants. We found that lung cancer risk scores were not aligned with lung cancer diagnoses in African American patients. In fact, African American individuals with screen-detected lung cancers were clustered predominantly in the lower risk quartiles. However, among White individuals undergoing screening, higher risk scores were associated with lung cancer diagnoses.

Our findings suggest that we should use caution in applying risk models to diverse populations, given that our current understanding of lung cancer risk is incomplete. Existing models have been derived from screening trials including 5% or fewer African American individuals and may not apply equitably to real-world screening participants.^{2,4} Although our study is limited by its single-center approach and short duration of follow-up, the racially diverse patient population allowed us to identify weaknesses in risk calculation. Determinants of health, including social constructs and environmental factors, may be critical moderators of lung cancer risk among underserved

Figure. Six-Year Lung Cancer Risk Among the Screening and Cancer Cohorts by Risk Quartile



A, Percentage of patients who underwent lung cancer screening in each risk quartile by race. African American and White individuals have overlapping percentages, with 24.9% and 25.2% of each cohort in risk quartile 2. B, Percentage of patients who had a screen-detected lung cancer in each risk quartile by race. The size of each data icon is

proportional to the number of patients in each risk quartile. Risk quartiles are determined by median PLCom2012 risk values (with cutoffs shown in parentheses along the x-axis).

populations. Further research on comprehensive risk prediction for underrepresented racial and ethnic populations should prioritize diversity and focus on additional factors related to socioeconomic status, geographic variables, the environment, and exposure history.

ARTICLE INFORMATION

Accepted for Publication: February 13, 2021.

Published: April 6, 2021. doi:[10.1001/jamanetworkopen.2021.4509](https://doi.org/10.1001/jamanetworkopen.2021.4509)

Open Access: This is an open access article distributed under the terms of the [CC-BY License](#). © 2021 Shusted CS et al. *JAMA Network Open*.

Corresponding Author: Julie A. Barta, MD, The Jane and Leonard Korman Respiratory Institute, Division of Pulmonary and Critical Care Medicine, Department of Medicine, Sidney Kimmel Medical College at Thomas Jefferson University, 834 Walnut St, Ste 650, Philadelphia, PA (Julie.Barta@jefferson.edu).

Author Affiliations: The Jane and Leonard Korman Respiratory Institute, Division of Pulmonary and Critical Care Medicine, Department of Medicine, Sidney Kimmel Medical College at Thomas Jefferson University, Philadelphia, Pennsylvania (Shusted, Kane, Barta); The Jane and Leonard Korman Respiratory Institute, Division of Thoracic Surgery, Department of Surgery, Sidney Kimmel Medical College at Thomas Jefferson University, Philadelphia, Pennsylvania (Evans); Division of Population Science, Department of Medical Oncology, Thomas Jefferson University, Philadelphia, Pennsylvania (Juon).

Author Contributions: Dr Barta had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Shusted, Evans, Kane, Barta.

Acquisition, analysis, or interpretation of data: Shusted, Juon, Barta.

Drafting of the manuscript: Shusted, Evans, Juon, Barta.

Critical revision of the manuscript for important intellectual content: Shusted, Evans, Kane, Barta.

Statistical analysis: Shusted, Juon.

Obtained funding: Evans, Kane.

Administrative, technical, or material support: Evans, Kane, Barta.

Supervision: Evans, Barta.

Conflict of Interest Disclosures: Dr Barta reported receiving grants from the Genentech Health Equity Innovations Fund and the Prevent Cancer Foundation outside the submitted work. No other disclosures were reported.

Funding/Support: This work was supported in part by the Bristol Myers-Squibb Foundation's Specialty Care for Vulnerable Populations initiative.

Role of the Funder/Sponsor: The funder had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Disclaimer: All information and materials in the manuscript are original.

Meeting Presentation: This study premise was presented as an e-poster at the American Thoracic Society International Conference 2020; available online from August 5 to November 10, 2020.

REFERENCES

1. Tanner NT, Gebregziabher M, Hughes Halbert C, Payne E, Egede LE, Silvestri GA. Racial differences in outcomes within the national lung screening trial. implications for widespread implementation. *Am J Respir Crit Care Med*. 2015;192(2):200-208. doi:[10.1164/rccm.201502-0259OC](https://doi.org/10.1164/rccm.201502-0259OC)
2. Aberle DR, Adams AM, Berg CD, et al; National Lung Screening Trial Research Team. Reduced lung-cancer mortality with low-dose computed tomographic screening. *N Engl J Med*. 2011;365(5):395-409. doi:[10.1056/NEJMoa1102873](https://doi.org/10.1056/NEJMoa1102873)
3. Aldrich MC, Mercaldo SF, Sandler KL, Blot WJ, Grogan EL, Blume JD. Evaluation of USPSTF lung cancer screening guidelines among African American adult smokers. *JAMA Oncol*. 2019;5(9):1318-1324. doi:[10.1001/jamaoncol.2019.1402](https://doi.org/10.1001/jamaoncol.2019.1402)
4. Katki HA, Kovalchik SA, Berg CD, Cheung LC, Chaturvedi AK. Development and validation of risk models to select ever-smokers for CT lung cancer screening. *JAMA*. 2016;315(21):2300-2311. doi:[10.1001/jama.2016.6255](https://doi.org/10.1001/jama.2016.6255)

5. Pasquinelli MM, Tammemägi MC, Kovitz KL, et al. Risk prediction model versus United States Preventive Services Task Force lung cancer screening eligibility criteria: reducing race disparities. *J Thorac Oncol*. 2020;15(11):1738-1747. doi:[10.1016/j.jtho.2020.08.006](https://doi.org/10.1016/j.jtho.2020.08.006)
6. Tammemägi MC, Katki HA, Hocking WG, et al. Selection criteria for lung-cancer screening. *N Engl J Med*. 2013;368(8):728-736. doi:[10.1056/NEJMoa1211776](https://doi.org/10.1056/NEJMoa1211776)