

2-2019

# Does Concomitant CABG Influence the Outcomes of Post-Myocardial Infarction Ventricular Septal Defect Repair?

Dylan P. Horan

*Thomas Jefferson University, dylan.horan@jefferson.edu*

Jae Hwan Choi

*Thomas Jefferson University, jae.choi2@jefferson.edu*

Matthew P. Weber

*Thomas Jefferson University, matthew.c.weber@jefferson.edu*

Elizabeth Maynes

*Thomas Jefferson University, elizabeth.maynes@jefferson.edu*

Sinal Patel

*Thomas Jefferson University, sinal.patel@jefferson.edu**See next page for additional authors*

## [Let us know how access to this document benefits you](#)

Follow this and additional works at: [https://jdc.jefferson.edu/si\\_ctr\\_2021\\_phase1](https://jdc.jefferson.edu/si_ctr_2021_phase1) Part of the [Cardiology Commons](#)

---

### Recommended Citation

Horan, Dylan P.; Choi, Jae Hwan; Weber, Matthew P.; Maynes, Elizabeth J.; Patel, Sinal; Challapalli, Jothika; Luc, Jessica G.Y.; Entwistle, John W.; Massey, H. Todd; Morris, Rohinton J.; and Tchantchaleishvili, Vakhtang, "Does Concomitant CABG Influence the Outcomes of Post-Myocardial Infarction Ventricular Septal Defect Repair?" (2019). SKMC JeffMD Scholarly Inquiry, Phase 1, Project 1.

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

---

**Authors**

Dylan P. Horan, Jae Hwan Choi, Matthew P. Weber, Elizabeth Maynes, Sinal Patel, Jothika Challapalli, Jessica G.Y. Luc, John W. Entwistle, H. Todd Massey, Rohinton J. Morris, and Vakhtang Tchantchaleishvili

Dylan Horan  
SKMC Class of 2021  
SI CTR Abstract  
Date 12/14/2018

**Title:** Does Concomitant CABG Influence the Outcomes of Post-Myocardial Infarction  
Ventricular Septal Defect Repair?

**Authors:** Dylan P. Horan, Jae Hwan Choi, Matthew P. Weber, Elizabeth J. Maynes, Sinal Patel,  
Jothika Challapalli, Jessica G.Y. Luc, John W. Entwistle, H. Todd Massey, Rohinton J. Morris,  
Vakhtang Tchantchaleishvili

**Introduction:** Ventricular septal defect (VSD) following myocardial infarction (MI) is a relatively infrequent complication with high mortality. Over time, understanding of the pathology and its management has resulted in improved outcomes; however, controversies remain.

**Objective:** We sought to investigate the effect of concomitant coronary artery bypass graft (CABG) on outcomes following post-MI VSD repair.

**Methods:** Electronic search was performed to identify all relevant studies published from 2000 to 2018. After assessment for inclusion and exclusion criteria, 66 studies were selected for the analysis. Data were extracted and pooled for systematic review and meta-analysis.

**Results:** Average age was 68.7 years (95% CI 67.3-70.1) with 57% (95% CI 54-60) males. Coronary angiogram was available preoperatively in 94% (95% CI 92-96) of patients. Single-vessel disease was most common (47%, 95% CI 42-52) with left anterior descending coronary artery the most commonly involved vessel (55%, 95% CI 46-63). Concomitant CABG was performed in 52% (95% CI 46-57) of patients. Of these, infarcted territory was revascularized in

54% (95% CI 23-82). No significant survival difference was observed between those who had concomitant CABG versus those without CABG at 30 days (65%, 95% CI 58-72) vs (60%, 95% CI 47-72), 1 year (59%, 95% CI 50-68) vs (51%, 95% CI 41-61), and 5 years (46%, 95% CI 38-54) vs (39%, 95% CI 27-52) respectively.

**Discussion:** Overall, concomitant CABG did not have a significant effect on survival following VSD repair, therefore, decision on revascularization should be weighed against the risks associated with prolonged cardiopulmonary bypass.