

12-7-2017

A Rare Case of Hemophagocytic Lymphohistiocytosis of Unknown Etiology

Ida Micaily
Abington Jefferson Health

Marie Helene Gagnon
Abington Jefferson Health

Richard Friedenheim
Abington Jefferson Health

Jason A. Damsker
Abington Hematology and Oncology Associates

Follow this and additional works at: <https://jdc.jefferson.edu/internalfp>

 Part of the [Internal Medicine Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Micaily, Ida; Gagnon, Marie Helene; Friedenheim, Richard; and Damsker, Jason A., "A Rare Case of Hemophagocytic Lymphohistiocytosis of Unknown Etiology" (2017). *Division of Internal Medicine Faculty Papers & Presentations*. Paper 20.

<https://jdc.jefferson.edu/internalfp/20>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Division of Internal Medicine Faculty Papers & Presentations by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

A Rare Case of Hemophagocytic Lymphohistiocytosis of Unknown Etiology
Ida Micaily¹, Marie-Helene Gagnon¹, Richard Friedenheim², Jason Damsker³
¹Department of Medicine, Abington Jefferson Health, Abington, PA, ²Abington
Pulmonary and Critical Care, Abington, PA, ³Abington Hematology and Oncology
Associates, Abington, PA

Abstract:

Hemophagocytic lymphohistiocytosis (HLH) is an uncommon cytokine storm syndrome marked by high mortality. In adults, acquired HLH usually has an underlying infectious, autoimmune or malignant process that triggers the syndrome. In this case report, we present a 64-year old Caucasian male presenting with productive cough, fevers, weight loss and altered mental status who was ultimately found to have HLH of unknown etiology.

Introduction:

According to the 2008 diagnostic criteria, HLH is established by a molecular diagnosis of pathologic mutation or through clinical and laboratory findings of fever, splenomegaly, cytopenias, hypertriglyceridemia and/or hypofibrinogenemia, tissue diagnosis of hemophagocytosis in the bone marrow, spleen or lymph nodes, low or absent natural killer cell activity, elevated ferritin (>500 ng/ mL), and soluble CD 25 > 2400 U/mL. As a cytokine storm syndrome, HLH is often confused for sepsis, macrophage-activation syndrome (MAS) and other multi-organ system failures. Early recognition of HLH can prompt timely treatment and improve morbidity and mortality. Current therapeutic options include immunosuppression, immune modulation, chemotherapy, and biological response modification followed by bone marrow transplant.

Case Report:

A 64-year old Caucasian gentleman presented to the Emergency Department with complaints of lethargy, weakness, progressive dyspnea on exertion, productive cough of whitish sputum and a fever of 102.5 F. His past medical history was significant for non-insulin dependent diabetes, CAD s/p stent 14 years prior, hypertension, and atrial fibrillation. He had a 14 pack year smoking history and had quit smoking 13 years prior. On review of systems, he admitted to a 5 lbs weight loss over the past 3 months, intermittent bouts of confusion and recent leg swelling. Deep venous thrombosis was excluded by his primary care physician, a week prior to admission. While in the ED, a CBC showed pancytopenia, marked by a white blood cell count of 2.7 and anemia with hemoglobin of 10.1. A CXR was performed showing bilateral pulmonary nodules; a CT scan of his chest and abdomen showed extensive pulmonary nodules and moderate splenomegaly. The patient was started on empiric antibiotics for presumed pneumonia with concerns of sepsis.

An exhaustive approach for a possible malignant etiology of the patient's lymphadenopathy was initiated on the second day of hospitalization. A CT-guided lung biopsy and bone marrow biopsy were initially conducted which were negative for any abnormal cells. Additionally, a PET-CT, VATS procedure, and several more

bone marrow biopsies were performed, without any explanation for his symptoms. An extensive rheumatologic work up only showed mild positivity for anti-cardiolipin IgG and beta-2 glycoprotein 1 antibody IgA positivity. Infectious work up for Rickettsia, Histoplasmosis, Blastomycosis, Bartonella, Syphilis, Tuberculosis, Babesiosis, fungal, viral, and other potential infectious etiologies were unrevealing. Of interest, the patient had one AFB positive sample from his VATS procedure after two weeks, however this specimen was concluded to be non-correlative. All repeat AFB cultures and sputum analyses were negative. His hospital course was prolonged and complicated by bouts of altered mental status, continuing fevers and worsening pancytopenia.

Approximately three hospital re-admissions for sepsis-like syndrome, a splenectomy was performed with pathology showing hemophagocytosis. At the time of the splenectomy, an elevated ferritin value of > 70000 was noted, and fibrinogen levels were low. The patient was additionally found to have an elevated IL-2R. Thus, a diagnosis of HLH was made given pancytopenia, ferritin of 70000, elevated triglycerides, hypofibrinogenemia, elevated IL-2R, splenomegaly, hemophagocytosis on splenectomy and fevers. The patient was treated with etoposide and dexamethasone by his oncologist, per protocol, to which his cytopenia, ferritin and fibrinogen improved.

Conclusion:

HLH is a rare disease entity that is often confused for sepsis syndrome. An extensive work up is urgent and necessary to adequately treat patients to reduce morbidity and mortality.

References:

- Scott RB, Robb-Smith AHT. Histiocytic medullary reticulosis. *Lancet*. 1939;234:194–198
- Jordan MB, Filipovich AH. Hematopoietic cell transplantation for hemophagocytic lymphohistiocytosis: a journey of a thousand miles begins with a single (big) step. *Bone Marrow Transplantation* (2008) 42, 433–437
- Janka GE. Hemophagocytic lymphohistiocytosis. *Hematology*. 2005;10(Suppl 1):104–107
- Varghese D, Haseer Koya H, Cherian SV, et al. Hemophagocytic lymphohistiocytosis: an uncommon presentation of enteropathy-associated T-cell lymphoma. *J Clin Oncol*. 2013;31(13):e226–e230
- Kleynberg RL, Schiller GJ. Secondary hemophagocytic lymphohistiocytosis in adults: An update on diagnosis and therapy. *Clin Adv Hematol Oncol*. 2012;10:726–32.
- Shabbir M, Lucas J, Lazarchick J, Shirai K. Secondary hemophagocytic syndrome in adult series of 18 patients in a single institution and a review of literature. *Hematol Oncol*. 2010;29:100–6