
4-30-2015

Conference Poster Season

Daniel G. Kipnis, MSI
Thomas Jefferson University, kipnis007@gmail.com

Follow this and additional works at: <https://jdc.jefferson.edu/jdcnews>



Part of the [Education Commons](#), and the [Medicine and Health Sciences Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Kipnis, MSI, Daniel G., "Conference Poster Season" (2015). *Jefferson Digital Commons News*. Paper 21.

<https://jdc.jefferson.edu/jdcnews/21>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Jefferson Digital Commons News by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Conference Poster Season

The [Jefferson Digital Commons](#) (JDC) has nearly 500 [archived conference posters](#). Don't let **your** conference posters collect dust and sit in hallways where only your colleagues view them on a daily basis. Send them to the JDC to reach a global audience.

Jefferson

When is it Safe for Patients to Drive after Right Total Hip Arthroplasty?

Victor H. Hernandez MD, Alvin C. Ong MD, Fabio Orozco MD, Anne Marie Madden, Zachary D. Post MD.
Investigation performed at the Rothman Institute at Thomas Jefferson University, Philadelphia, PA.

ROTHMAN INSTITUTE

INTRODUCTION

- Driving restrictions after total hip arthroplasty (THA) can be inconvenient and burdensome for patients.
- When patients may safely be allowed to drive remains controversial.
- Most studies recommend 6 weeks but recent advances in surgical approach, pain management and rapid recovery may have changed this time frame.
- The purpose of this study was to prospectively evaluate driving safety after THA through brake reaction time.

MATERIALS AND METHODS

Twenty nine patients who underwent right THA were prospectively evaluated between October 2013 and June 2014. Driving performance was evaluated using the Brake Reaction Test (BRT) measuring brake time after a stimulus (Figure 1). All patients underwent preoperative assessment to establish a baseline. Patients were then re-tested at 2, 4 and 6 weeks post-operatively. Patients were allowed to drive when the post-operative reaction time was equal to or less than the pre-operative baseline. In addition at each testing patients were asked if they felt ready to drive.

General linear repeated measurement was used for analysis.

RESULTS

Of the 38 study patients, 33 (87%) reached their baseline time by 2 weeks. Five patients (13%) reached their baseline at 4 weeks. There were no differences with respect to age, gender, or the use of assistance devices. Of the 33 patients, 24 (73%) stated that they felt ready to drive while 5 (15%) were not sure and 4 (12%) felt they were not ready. Of the 5 patients who returned to baseline at 4 weeks, 2 patients said they thought they were ready while 3 patients did not.

However, even the five patients that did not return to their baseline by two weeks were still under the nationally recommended safe brake time standard of 1.25 seconds which is based upon several scientific studies looking at BRT in all types of patients, surgical and non-surgical alike.

FIGURE 1




TABLE 1

Patient	Age (years)	Gender	T - Pre	T 2-W	T 4-W
1	75	Female	0.703	1.03	0.566
2	52	Male	0.593	0.638	0.567
3	51	Male	0.454	0.478	0.449
4	54	Female	0.598	0.640	0.472
5	64	Female	0.515	0.674	0.455

Table 1 Characteristics of the 5 patients that did not reach their baseline at the 2 weeks.
T-Pre: Pre-operative, 2 weeks

DISCUSSION

Nearly all patients were able to return to normal BRT by 2 weeks after THA. In addition, patient perception of driving ability accurately predicted return of BRT. This represents a substantial improvement from current recommendations.

These findings have allowed us to encourage patients to re-evaluate their driving ability as soon as 2 weeks after THA.

Follow the lead of the Rothman Institute at Thomas Jefferson University. They have just sent the following posters for deposit:

- [Trigger Finger Release Performed Wide Awake: Prospective Comparison of Local Anesthetics](#)
- [Radiation Exposure to the Eye with Mini C-arm Use During Hand Surgery](#)
- [The Patellar Tendon Can Cause External Tibial Component Malrotation in Lateral Unicompartmental Knee Arthroplasty](#)
- [Rotational Alignment in Medial Unicompartmental Knee Arthroplasty: Comparison of Anatomic Landmarks](#)
- [Burden of Periprosthetic Joint Infection and its Treatment on Kidney Function](#)
- [Red Cell Distribution Width: an Unacknowledged Predictor of Mortality and Length of Stay following Revision Arthroplasty](#)
- [When is it Safe for Patients to Drive after Right Total Hip Arthroplasty?](#)
- [Low Dose Aspirin: An Effective Chemoprophylaxis for Preventing Venous Thromboembolic Events](#)
- [Aspirin May Be Adequate for Venous Thromboembolic Event Prophylaxis after Revision Hip and Knee Arthroplasty](#)

Send your conference posters to dan.kipnis@jefferson.edu for inclusion in the **Jefferson Digital Commons**.

Posted online: April 30, 2015, <http://library.jefferson.edu/librarynews/?p=6859>