

2012

Optical Imaging for Determination of Apoptosis Medicated Therapeutic Efficacy

Kaijun Zhang
Thomas Jefferson University

Bishnuhari Paudyal
Thomas Jefferson University

Neil Mehta
Thomas Jefferson University

Brian Gray
Molecular Targeting Technologies, Inc.

Koon Y. Pak
Molecular Targeting Technologies, Inc.
Follow this and additional works at: <https://jdc.jefferson.edu/radiologyfp>

 Part of the [Medical Molecular Biology Commons](#), and the [Radiology Commons](#)
See next page for additional authors

[Let us know how access to this document benefits you](#)

Recommended Citation

Zhang, Kaijun; Paudyal, Bishnuhari; Mehta, Neil; Gray, Brian; Pak, Koon Y.; Wickstrom, Eric; and Thakur, Mathew L, "Optical Imaging for Determination of Apoptosis Medicated Therapeutic Efficacy" (2012). *Department of Radiology Faculty Papers*. Paper 19.
<https://jdc.jefferson.edu/radiologyfp/19>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Radiology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Kaijun Zhang, Bishnuhari Paudyal, Neil Mehta, Brian Gray, Koon Y. Pak, Eric Wickstrom, and Mathew L. Thakur

OPTICAL IMAGING FOR DETERMINATION OF APOPTOSIS MEDIATED THERAPEUTIC EFFICACY










K Zhang, PhD¹, B. Paudyal, PhD¹, N. Mehta¹, B Gray, PhD⁴, C Pak, PhD⁴, E Wickstrom, PhD^{2,3}, M. Thakur, PhD^{1,3}

Departments of: Radiology¹, Biochem. & Mol. Biol.², Kimmel Cancer Center³, Thomas Jefferson University, Philadelphia, PA; Molecular Targeting Technologies, Inc. ⁴, West Chester, PA

Objectives: Targeting apoptosis to assess therapeutic response is uncommon. The purpose was to evaluate the use of two near infrared (NIR) fluorophores to monitor the effectiveness of breast cancer (BC) therapy. Both fluorophores have high affinity for phosphatidylserine (PS) released on the outer leaflet of apoptotic cells mediated by therapeutic intervention. The long term goal is to determine if optical imaging (OI) can play a vital role in determination of therapeutic efficacy noninvasively.

Methods: Two NIR fluorophores linked bis-zinc (II) dipicolylamine coordination complexes, PSVue@643 and PSVue@794, served as the probes. Athymic nude mice bearing human BC tumors BT474, received 16.5 µg of one of the NIR probes i.v. before and after 40 µg of doxorubicin (DOX) given once i.p. OI was performed up to 24 hrs post injection (PSVue@643; Abs. 643 nm; Em. 658 nm; PSVue@794; Abs. 794 nm; Em. 810 nm). Tumor/muscle (T/M) and organ/muscle (O/M) ratios were calculated and plotted.

Results: T/M ratios for PSVue@794 were 200% times higher than for PSVue@643 (P = <0.05). Both probes showed 25%-30% increased T/M ratios post DOX treatment, (P = <0.05) indicative of enhanced apoptosis. With PSVue@643 tumor intensity declined over time, but increased for intestine. With PSVue@794 tumor intensity and T/M ratios increased as a function of time, with a decreased M/O ratios for all organs.



Conclusion: OI of apoptotic BC cells mediated by DOX treatment permits to determine the effectiveness of DOX within 24 hrs. These results are consistent with those in another investigation in which F-18-FDG was used to monitor diminished metabolic activity following DOX treatment. PSVue@794 which eliminates radiation burden to normal organs is a preferable NIR fluorophore for determination of therapeutic effectiveness of BC by OI. **Support:** NIH 1S10 RR026678-01 (MLT).