

1-2020

Predictors of Negative Outcomes and Causes of Loss to Follow Up Among Breast Cancer Patients in Port-au-Prince, Haiti

A. Johnson

Thomas Jefferson University, adrienne.johnson@jefferson.edu

R. R. Henderson

Innovating Health International, Port-au-Prince, Haiti

G. Jones

Innovating Health International, Port-au-Prince, Haiti

Y. Fujii

Innovating Health International, Port-au-Prince, Haiti

P. Haley

Innovating Health International, Port-au-Prince, Haiti

Follow this and additional works at: https://jdc.jefferson.edu/si_phr_2022_phase1



Part of the [Women's Health Commons](#)

See next page for additional authors

[Let us know how access to this document benefits you](#)

Recommended Citation

Johnson, A.; Henderson, R. R.; Jones, G.; Fujii, Y.; Haley, P.; Bernard, J.; and DeGennaro, V., "Predictors of Negative Outcomes and Causes of Loss to Follow Up Among Breast Cancer Patients in Port-au-Prince, Haiti" (2020). *Phase 1*. Paper 30.

https://jdc.jefferson.edu/si_phr_2022_phase1/30

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

A. Johnson, R. R. Henderson, G. Jones, Y. Fujii, P. Haley, J. Bernard, and V. DeGennaro

SKMC Class of 2022

Word Count: 249

Predictors of Negative Outcomes and Causes of Loss to Follow Up Among Breast Cancer Patients in Port-au-Prince, Haiti

A Johnson, R R Henderson, G Jones, Y Fujii, P Haley, J Bernard, V DeGennaro*
Innovating Health International, Port-au-Prince, Haiti

Objective: To determine predictors of negative outcomes and causes of loss to follow-up (LTFU) in patients with breast cancer in Haiti.

Design and Methods: Patients seen by Innovating Health International's (IHI) cancer program were designated as LTFU after at least six months of non-contact (n=606). A cohort of LTFU breast cancer patients (n=101) was compared to the larger population of breast cancer patients (n=939), and a regression model constructed in order to identify risks for death and LTFU. Multiple calls were made to contact LTFU patients, and reasons for LTFU were recorded.

Results: Death and LTFU status were associated with advanced stage ($p < 0.0005$), higher ECOG ($p = 0.011$), and longer care ($p < 0.0005$; $p = .03$); LTFU status was lower with positive family history ($p = 0.022$). 42.7% of LTFU patients were successfully contacted and 37.9% of these were reported deceased. The most common reasons for LTFU to clinic were obtaining care elsewhere and difficulty accessing care (due to distance, unrest, and money).

Conclusions: Understanding causes of LTFU can suggest measures to reduce risk. LTFU was driven by the same factors as mortality, suggesting many "lost" patients may be deceased; this

conclusion is furthermore supported by the low rate of successful recontact and high rate of death in the LTFU population. These conclusions support the need for improved palliative care outreach. Furthermore, LTFU status in Haiti is in part due to difficulties accessing care due to issues of politics, infrastructure, and economics.