

5-6-2014

Permeation Redux: Thermodynamics and Kinetics of Ion Movement through Potassium Channels.

Richard Horn

Department of Molecular Physiology and Biophysics, Jefferson Medical College, richard.horn@jefferson.edu

Benoît Roux

Department of Biochemistry and Molecular Biology, University of Chicago

Johan Aqvist

Department of Cell and Molecular Biology, Uppsala University, Biomedical Center, Uppsala, Sweden

Let us know how access to this document benefits you

Follow this and additional works at: <https://jdc.jefferson.edu/physfp> Part of the [Other Medical Specialties Commons](#)

Recommended Citation

Horn, Richard; Roux, Benoît; and Aqvist, Johan, "Permeation Redux: Thermodynamics and Kinetics of Ion Movement through Potassium Channels." (2014). *Department of Molecular Physiology and Biophysics Faculty Papers*. Paper 14.

<https://jdc.jefferson.edu/physfp/14>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Molecular Physiology and Biophysics Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Permeation Redux: Thermodynamics and Kinetics of Ion Movement through Potassium Channels

Richard Horn,^{†*} Benoît Roux,[‡] and Johan Åqvist[§]

[†]Department of Molecular Physiology and Biophysics, Jefferson Medical College, Philadelphia, Pennsylvania; [‡]Department of Biochemistry and Molecular Biology, The University of Chicago, Chicago, Illinois; and [§]Department of Cell & Molecular Biology, Uppsala University, Biomedical Center, Uppsala, Sweden


ABSTRACT The fundamental biophysics underlying the selective movement of ions through ion channels was launched by George Eisenman in the 1960s, using glass electrodes. This minireview examines the insights from these early studies and the explosive progress made since then.

The recent passing of George Eisenman (December 18, 2013) inspired us to revisit the topic most associated with his passionate input, namely how the membrane proteins known as ion channels control passive movements of ions across biological membranes. Ion permeation has captivated biophysicists for more than half a century, and only now, with the combined advent of atomic-level structures and sophisticated computational wizardry, are the secrets of this amazing process beginning to be revealed. Why “amazing”? For example, because K⁺-selective ion channels can discriminate between K⁺ and Na⁺ ions, which differ in radius by a mere 0.38 Ångstrom, and do so with 1000:1 reliability and at lightning speed near the diffusion limit, the dwell time of an ion in the pore of a channel is as fleeting as ~10⁻⁸ s. Understanding this remarkably-tuned process in K⁺ channels requires attention to two perspectives: the ability of specific channels to discriminate between the ions they might encounter (i.e., selectivity); and the kinetics of ion movement across the channel pore (i.e., conduction).


The classical thermodynamic explanation of ion selectivity is that the relative free energy difference of ions in the pore relative to the bulk solution is the critical quantity to consider (1–4). Some of the earliest insights into thermodynamic selectivity derive from studies of ion binding to aluminosilicate glass electrodes (5,6). Depending on the composition of the glass, these electrodes, originally developed for their proton sensitivity, can exhibit a dramatic range of selectivities among the five alkali metal cations. In rank order, one might expect as many as 5 × 4 × 3 × 2 × 1 = 120 different sequences of selectivities among these five cations. Remarkably, however, in the vast literature of selectivity in biological membranes, typically only 11 sequences are observed (with some exceptions). These became known as the “Eisenman sequences”. The exact same selectivity sequences are observed in glass electrodes of various compositions.

Why are the free energy differences the way they are for a given system? To answer this question, one needs a physical mechanism. For Eisenman, numerical calculations stood as a critical component of the process of better understanding Nature. In other words, proposing a physical mechanism that is qualitatively reasonable is not enough—one must also test it by constructing atomic models leading to actual quantitative predictions (Fig. 1). In the early days, the concept of the anionic field strength of a binding site was formulated and tested with direct calculations based on exceedingly simple atomic hard-sphere models of ions, water molecules, and coordinating ligands such as shown in Fig. 1 A (2,5). Remarkably, these simple calculations led to the Eisenman selectivity sequences. Eisenman was able to account for the limited class of sequences by considering the equilibrium binding of cations to the glass, and the energetic competition between water and glass for the ions. The critical factor that determines the selectivity sequence of a given glass is the anionic field strength of the binding site on the glass. Briefly, the smallest group Ia cation, Li⁺, holds water most tenaciously, so it will only dehydrate and bind in the presence of a strongly negative electrostatic potential.

By contrast, the largest cation, Cs⁺, holds water least tenaciously. It cannot bind readily to a strongly negative site because the site itself greedily clings to water molecules, and thus prevents Cs⁺ binding. However, Cs⁺ is more willing, relative to the smaller cations, to dehydrate and bind in the presence of a weakly negative electrostatic potential. At the extremes, the highest anionic field strength glass shows a selectivity sequence of



and the lowest anionic field strength glass shows a selectivity sequence of



A very simple model, based on the relative Gibbs' free energies of binding and hydration, explains why there are only 11 sequences (5–7). The critical factor underlying the pattern of these selectivity sequences is that the “ion-site

Submitted January 31, 2014, and accepted for publication March 27, 2014.

*Correspondence: richardspikehorn@gmail.com

Editor: Brian Salzberg.

© 2014 by the Biophysical Society
0006-3495/14/05/1859/5 \$2.00






FIGURE 1 Structural models used in theoretical studies of ion selectivity. (A) Simple model used to introduce the concept of field strength leading to 11 cationic selectivity sequences (2,5,6). Ions, water, and ligands are represented by simple hard-spheres with embedded point charges. Selectivity arises from the difference in the interaction energy of the cation with a water molecule (top) and an anionic coordinating ligand (bottom). (B) Ion-selective transfer process is depicted with atomic models incorporating all molecular details in the case of solvation in liquid water (top) and binding to the K^+ -selective ionophore valinomycin (bottom). Such atomic models were used to carry out some of the earliest MD free energy simulations on ion binding selectivity (12,13,15).

interaction energies fall off as a function of cation size as a lower power of the cation radius than do ion-water interaction energies” (5,6). The icing on the cake is that ion selectivity of channels in membranes appears to follow similar principles (7). The thermodynamic principles are evidently analogous. Moreover, Eisenman’s contributions went far beyond the monovalent cation selectivity of potassium channels. His theoretical approach was seminal in understanding both cation and anion selectivity in a diverse range of physical and biological systems (8,9).

The advent of molecular dynamics (MD) simulations around this period was of critical importance to the field. This made it possible to construct increasingly realistic models of proteins (10), including ion channels (11), and examine the ion selectivity of carriers using the alchemical free energy perturbation (FEP) technique (12,13). With no experimental structures yet available for the ion-selective regions of biological K^+ channels, an important step forward was Eisenman’s realization that other ion-selective systems could be used to computationally test the structural basis of his selectivity theory. Both peptidic small ionophores, such as valinomycin and nonactin, and the ion-coordinating fivefold symmetry sites in icosahedral virus structures, thus caught his attention (13). As it turned out, these types of structures were indeed very relevant for the selectivity problem, because K^+ -channel filters were even-

tually shown to be lined likewise by carbonyl groups (14). With the crystallographically determined valinomycin structure at hand, its selectivity could be energetically analyzed by atomistic computer simulations, as illustrated in Fig. 1 B (12,15). The anionic field strength (represented by the carbonyl ligand dipole moment) could then be varied artificially, and the successive progression through the different selectivity sequences, as a function of field strength, directly observed. Likewise, Eisenman and Alvarez (13) made computational predictions for the binding energetics and selectivity of the Ca^{2+} binding site at the fivefold symmetry axis of satellite tobacco necrosis virus, and they subsequently showed experimentally that this binding site had a marked rare-earth ion size selectivity (16). To this day, the general computational FEP/MD framework based on equilibrium thermodynamics used in these studies continues to be a critical tool to understand ion channels (17), transporters (18), and pumps (19).

Despite these early insights, it was always clear to Eisenman that explanations of selectivity solely based on thermodynamic equilibrium were too simple to account for the detailed properties observed in biological systems. Since the halcyon days of equilibrium binding studies on glass electrodes, the permeation landscape presented by the pores of ion channels has emerged as richer than anticipated. One important realization is that binding and

conduction of ions through a channel may act as contradictory processes, because although an ion has to leave the comfort of its hydration shells to selectively enter the mouth of a channel pore, if it binds the channel too tightly, it cannot move rapidly through it. This mini-conundrum is most apparent, perhaps, for K^+ -channels, which attract K^+ ions much more forcefully than Na^+ ions, yet conduct K^+ ions much faster than Na^+ ions.

Another factor evident in early studies of permeation is that ions encounter a series of obstacles (i.e., energy barriers) and binding sites (i.e., energy wells) as they wend their way through the pore. One approach to understanding permeation is to consider that ions hopscotch from one well to the next over a series of barriers. When the number of barriers is rather limited, say <5 , one can use so-called “rate theory” (20) to analyze and formulate the free energy profile experienced by an ion crossing the membrane. Hille (21) proposed that selectivity derives largely from the selectivities of the barriers, not the wells. Eisenman and Horn (7) later considered the possibility that binding sites and barriers within a particular channel might have different selectivity sequences. For example, if a channel presents two barriers, one of which has selectivity sequence I and the other has selectivity sequence XI, the channel as a whole will have an intermediate selectivity sequence that is not an Eisenman sequence at all. Rather, it is a so-called “polarizability sequence” (7). Interestingly, contemporary studies indicate that successive binding sites along K^+ -selective channels display different selectivities (22). Another concept based on Eyring barrier models is that the energy levels for wells and barriers may not be static, and may therefore fluctuate on a timescale relevant to ion permeation (23). Finally, the biophysics of ion permeation and later structural studies show that multiple ions may cohabit the same channel simultaneously, and the interactions among these ions have profound consequences for ion conduction and selectivity.

Fast forward to the 21st century: atomic-level structures and all-atom simulations seem to have blown the permeation field wide open, as suggested by recent reviews (24–27). Once the KcsA channel structure was solved (14), the structural origin of K^+ -ion permeation could finally be addressed by computer simulations of the “real structure” and a number MD simulation studies provided novel insight (22,28–31). Needless to say, George Eisenman took great interest in these simulations even though he had by then retired. Also, in the case of KcsA, the initial work largely revolved around calculations of equilibrium ion binding and selectivity, barrier heights, and energy landscape mapping (22,31), because direct all-atom simulations of spontaneous permeation were not possible. However, the general type of knock-on mechanism with multiion occupancy of the channel selectivity filter, involving key distinct states (22,31), and a surprisingly flat energy landscape (22), appear to be robust features of these channels.

Even with the advent of MD simulations, the concept of field strength has kept its relevance. For example, the selectivity filter in MD simulations of the KcsA channel displayed a range of atomic flexibility that seemed somewhat shocking at the time because a traditional host-guest mechanism of selectivity would require a fairly rigid cavity-size. Yet, free energy computations indicated that this was not strictly necessary to establish the thermodynamic free energy differences needed to support ion selectivity (32). The resilience of Eisenman’s ideas is not entirely surprising because, as foreseen early on by Bertil Hille (21), the concept of field strength remains “useful if the dipoles of the channel are free to move and can be pulled in by small ions and pushed back by large ones”.

Nevertheless, despite the exciting progress, the chapter on ion selectivity in K^+ channels is far from closed. Very recently, a number of studies have revealed some extremely intriguing multiion aspects of selectivity in K^+ channels that appear to stand squarely outside the realm of equilibrium thermodynamics. By examining the properties of MthK (33) and NaK (34) mutants, Liu and Lockless (35) and Sauer et al. (36) showed that the channel becomes K^+ -selective only if there are four consecutive binding sites along the filter. This has culminated more recently with studies of two engineered mutants of the NaK channel, referred to as “NaK2K” and “NaK2CNG”. According to reversal potential measurements from single-channel electrophysiology, the NaK2K construct is K^+ -selective and the NaK2CNG construct is nonselective. Remarkably, despite being nonselective in ion permeation, the NaK2CNG filter displays an equilibrium preference for binding K^+ over Na^+ , as indicated by measurements with isothermal titration calorimetry and concentration-dependent ion replacement within the filter observed through crystallographic titration experiments.

K^+ -selective channels bind two or more K^+ ions in the narrow filter, whereas the nonselective channels bind fewer ions. Based on the crystallographic titration experiments, the NaK2K construct has two high-affinity K^+ sites whereas the NaK2CNG construct has only one K^+ -selective site. These experiments show that both K^+ -selective and nonselective channels select K^+ over Na^+ ions at equilibrium, implying that equilibrium selectivity is insufficient to determine the selectivity of ion permeation (35,36). The data indicate that having multiple K^+ ions bound simultaneously is required for selective K^+ conduction, and that a reduction in the number of bound K^+ ions destroys the multiion selectivity mechanism utilized by K^+ channels. Although these experimental results are intriguing, the underlying microscopic mechanisms remain unclear. The implication is that the multiion character of the permeation process must, somehow, be a critical element for establishing selective ion conduction through K^+ channels.

The progress made, and the challenges that remain, are perhaps best illustrated by returning to computational

studies of the simplest membrane spanning structure known, namely the gramicidin A channel. Before detailed studies of selectivity and conductance of K^+ -channels were launched, computational work on ion conduction through membrane channels was largely focused on this simple channel (37–41). In this case the permeation selectivity was monotonically size-dependent (Eisenman sequence I) and, in this respect, less interesting than K^+ -selective channels. However, from an energetic point of view it was puzzling how this single helical structure could yield free energy barriers low enough to permit high conductivity (7,42). Computer simulations of increasing complexity in this case established that the combined effect of several contributions to ion stabilization along the pore (from the protein, membrane, single-file waters, and bulk solution) indeed results in low barriers to permeation (11,39,40). Furthermore, the most realistic model comes in close agreement with experimental measurements (11,43), although it is clear that work is still needed.

REFERENCES

1. Bezanilla, F., and C. M. Armstrong. 1972. Negative conductance caused by entry of sodium and cesium ions into the K channels of squid axon. *J. Gen. Physiol.* 53:342–347.
2. Eisenman, G., and S. Krasne. 1973. Further considerations on the ion selectivity of carrier molecules and membranes. In IV International Biophysics Congress Symposium on Membrane Structure and Function, Moscow, Russia.
3. Neyton, J., and C. Miller. 1988. Discrete Ba^{2+} block as a probe of ion occupancy and pore structure in the high-conductance Ca^{2+} -activated K^+ channel. *J. Gen. Physiol.* 92:569–586.
4. Piasta, K. N., D. L. Theobald, and C. Miller. 2011. Potassium-selective block of barium permeation through single KcsA channels. *J. Gen. Physiol.* 138:421–436.
5. Eisenman, G. 1962. Cation selective glass electrodes and their mode of operation. *Biophys. J.* 2:259–323.
6. Eisenman, G. 1961. On the Elementary Atomic Origin of Equilibrium Ionic Specificity. Academic Press, New York.
7. Eisenman, G., and R. Horn. 1983. Ionic selectivity revisited: the role of kinetic and equilibrium processes in ion permeation through channels. *J. Membr. Biol.* 76:197–225.
8. Diamond, J. M., and E. M. Wright. 1969. Biological membranes: the physical basis of ion and nonelectrolyte selectivity. *Annu. Rev. Physiol.* 31:581–646.
9. Wright, E. M., and J. M. Diamond. 1977. Anion selectivity in biological systems. *Physiol. Rev.* 57:109–156.
10. McCammon, J. A., B. R. Gelin, and M. Karplus. 1977. Dynamics of folded proteins. *Nature.* 267:585–590.
11. Allen, T. W., O. S. Andersen, and B. Roux. 2004. Energetics of ion conduction through the gramicidin channel. *Proc. Natl. Acad. Sci. USA.* 101:117–122.
12. Åqvist, J., O. Alvarez, and G. Eisenman. 1992. Ion-selective properties of a small ionophore in methanol studied by free energy perturbation simulations. *J. Phys. Chem.* 96:10019–10025.
13. Eisenman, G., and O. Alvarez. 1991. Structure and function of channels and channelogs as studied by computational chemistry. *J. Membr. Biol.* 119:109–132.
14. Doyle, D. A., J. Morais Cabral, ..., R. MacKinnon. 1998. The structure of the potassium channel: molecular basis of K^+ conduction and selectivity. *Science.* 280:69–77.
15. Eisenman, G., J. Åqvist, and O. Alvarez. 1991. Free-energies underlying ion binding and transport in protein channels—free-energy perturbation simulations of ion binding and selectivity for valinomycin. *J. Chem. Soc., Faraday Trans.* 87:2099–2109.
16. Burroughs, S. E., G. Eisenman, and W. D. Horrocks, Jr. 1992. Characterization of the five-fold Ca^{2+} binding site of satellite tobacco necrosis virus using Eu^{3+} luminescence spectroscopy: a marked size-selectivity among rare earth ions. *Biophys. Chem.* 42:249–256.
17. Luzhkov, V. B., and J. Åqvist. 2001. K^+/Na^+ selectivity of the KcsA potassium channel from microscopic free energy perturbation calculations. *Biochim. Biophys. Acta.* 1548:194–202.
18. Noskov, S. Y., and B. Roux. 2008. Control of ion selectivity in LeuT: two Na^+ binding sites with two different mechanisms. *J. Mol. Biol.* 377:804–818.
19. Yu, H., I. M. Ratheal, ..., B. Roux. 2011. Protonation of key acidic residues is critical for the K^+ -selectivity of the Na/K pump. *Nat. Struct. Mol. Biol.* 18:1159–1163.
20. Zwolinski, B. I., H. Eyring, and C. E. Resese. 1949. Diffusion and membrane permeability. *J. Phys. Colloid Chem.* 53:1426–1453.
21. Hille, B. 1975. Ionic selectivity, saturation, and block in sodium channels. A four-barrier model. *J. Gen. Physiol.* 66:535–560.
22. Bernèche, S., and B. Roux. 2001. Energetics of ion conduction through the K^+ channel. *Nature.* 414:73–77.
23. Läger, P., W. Stephan, and E. Frehland. 1980. Fluctuations of barrier structure in ionic channels. *Biochim. Biophys. Acta.* 602:167–180.
24. Roux, B. 2010. Perspectives on: molecular dynamics and computational methods. *J. Gen. Physiol.* 135:547–548.
25. Dror, R. O., M. O. Jensen, ..., D. E. Shaw. 2010. Exploring atomic resolution physiology on a femtosecond to millisecond timescale using molecular dynamics simulations. *J. Gen. Physiol.* 135:555–562.
26. Eisenberg, B. 2013. Interacting ions in biophysics: real is not ideal. *Biophys. J.* 104:1849–1866.
27. Furini, S., and C. Domene. 2013. K^+ and Na^+ conduction in selective and nonselective ion channels via molecular dynamics simulations. *Biophys. J.* 105:1737–1745.
28. Guidoni, L., V. Torre, and P. Carloni. 1999. Potassium and sodium binding to the outer mouth of the K^+ channel. *Biochemistry.* 38:8599–8604.
29. Shrivastava, I. H., and M. S. Sansom. 2000. Simulations of ion permeation through a potassium channel: molecular dynamics of KcsA in a phospholipid bilayer. *Biophys. J.* 78:557–570.
30. Allen, T. W., S. Kuyucak, and S. H. Chung. 1999. Molecular dynamics study of the KcsA potassium channel. *Biophys. J.* 77:2502–2516.
31. Åqvist, J., and V. B. Luzhkov. 2000. Ion permeation mechanism of the potassium channel. *Nature.* 404:881–884.
32. Noskov, S. Y., S. Bernèche, and B. Roux. 2004. Control of ion selectivity in potassium channels by electrostatic and dynamic properties of carbonyl ligands. *Nature.* 431:830–834.
33. Ye, S., Y. Li, and Y. Jiang. 2010. Novel insights into K^+ selectivity from high-resolution structures of an open K^+ channel pore. *Nat. Struct. Mol. Biol.* 17:1019–1023.
34. Derebe, M. G., D. B. Sauer, ..., Y. Jiang. 2011. Tuning the ion selectivity of tetrameric cation channels by changing the number of ion binding sites. *Proc. Natl. Acad. Sci. USA.* 108:598–602.
35. Liu, S., and S. W. Lockless. 2013. Equilibrium selectivity alone does not create K^+ -selective ion conduction in K^+ channels. *Nat. Commun.* 4:2746.
36. Sauer, D. B., W. Zeng, ..., Y. Jiang. 2013. Sodium and potassium competition in potassium-selective and non-selective channels. *Nat. Commun.* 4:2721.
37. Kim, K. S., and E. Clementi. 1985. Energetics and hydration structure of a solvated gramicidin A channel for K^+ and Na^+ cations. *J. Am. Chem. Soc.* 107:5504–5513.
38. Mackay, D. H., P. H. Berens, ..., A. T. Hagler. 1984. Structure and dynamics of ion transport through gramicidin A. *Biophys. J.* 46:229–248.

39. Åqvist, J., and A. Warshel. 1989. Energetics of ion permeation through membrane channels. Solvation of Na^+ by gramicidin A. *Biophys. J.* 56:171–182.
40. Roux, B., and M. Karplus. 1993. Ion-transport in the gramicidin channel—free-energy of the solvated right-handed dimer in a model membrane. *J. Am. Chem. Soc.* 115:3250–3262.
41. Etchebest, C., S. Ranganathan, and A. Pullman. 1984. The gramicidin A channel: comparison of the energy profiles of Na^+ , K^+ and Cs^+ . Influence of the flexibility of the ethanolamine end chain on the profiles. *FEBS Lett.* 173:301–306.
42. Eisenman, G., and J. P. Sandblom. 1984. Modeling the gramicidin channel: interpretation of experimental data using rate theory. *Biophys. J.* 45:88–90.
43. Jensen, M. O., V. Jogini, ..., D. E. Shaw. 2013. Atomic-level simulation of current-voltage relationships in single-file ion channels. *J. Gen. Physiol.* 141:619–632.