

Department of Microbiology and Immunology **Faculty Papers**

Department of Microbiology and Immunology

5-1-2011

Microarray-based analysis of differential gene expression between infective and noninfective larvae of Strongyloides stercoralis.

Roshan Ramanathan

Laboratory of Parasitic Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland

Sudhir Varma Bioinformatics and Computational Biosciences Branch, National Institutes of Health, Bethesda, Maryland

José M C Ribeiro Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases National Hotitwigs of Healthi Bethe son Mary lands://idc.iefferson.edu/mifp

CreathyoGtMay@Eslical Cell Biology Commons, Medical Genetics Commons, and the Medical Immunology Microarray Research Facility, National Institutes of Health, Bethesda, Maryland

Let us know how access to this document benefits you

Department of Pathobiology, School of Veterinary Medicine, University of Pennsylvania, Philadelphia, Pennsylvania Recommended Citation

Ramanathan, Roshan; Varma, Sudhir; Ribeiro, José M C; Myers, Timothy G; Nolan, Thomas J; Abraham, Bayid Lok James Bi and Nytman, Thomas B, "Microarray-based analysis of differential gene expression between infective and noninfective larvae of Strongyloides stercoralis." (2011). Department of Microbiology and Immunology Faculty Papers. Paper 22. https://jdc.jefferson.edu/mifp/22

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's Center for Teaching and Learning (CTL). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Microbiology and Immunology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Roshan Ramanathan, Sudhir Varma, José M C Ribeiro, Timothy G Myers, Thomas J Nolan, David Abraham, James B Lok, and Thomas B Nutman

This article is available at Jefferson Digital Commons: https://jdc.jefferson.edu/mifp/22

Microarray-Based Analysis of Differential Gene Expression between Infective and Noninfective Larvae of *Strongyloides stercoralis*

Roshan Ramanathan¹*, Sudhir Varma², José M. C. Ribeiro³, Timothy G. Myers⁴, Thomas J. Nolan⁵, David Abraham⁶, James B. Lok⁵, Thomas B. Nutman¹

1 Laboratory of Parasitic Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland, United States of America, 2 Bioinformatics and Computational Biosciences Branch, National Institutes of Health, Bethesda, Maryland, United States of America, 3 Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases National Institutes of Health, Bethesda, Maryland, United States of America, 4 Microarray Research Facility, National Institutes of Health, Bethesda, Maryland, United States of America, 5 Department of Pathobiology, School of Veterinary Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, United States of America, 6 Department of Microbiology and Immunology, Thomas Jefferson University, Philadelphia, Pennsylvania, United States of America

Abstract

Background: Differences between noninfective first-stage (L1) and infective third-stage (L3i) larvae of parasitic nematode *Strongyloides stercoralis* at the molecular level are relatively uncharacterized. DNA microarrays were developed and utilized for this purpose.

Methods and Findings: Oligonucleotide hybridization probes for the array were designed to bind 3,571 putative mRNA transcripts predicted by analysis of 11,335 expressed sequence tags (ESTs) obtained as part of the Nematode EST project. RNA obtained from *S. stercoralis* L3i and L1 was co-hybridized to each array after labeling the individual samples with different fluorescent tags. Bioinformatic predictions of gene function were developed using a novel cDNA Annotation System software. We identified 935 differentially expressed genes (469 L3i-biased; 466 L1-biased) having two-fold expression differences or greater and microarray signals with a p value<0.01. Based on a functional analysis, L1 larvae have a larger number of genes putatively involved in transcription (p = 0.004), and L3i larvae have biased expression of putative heat shock proteins (such as *hsp-90*). Genes with products known to be immunoreactive in *S. stercoralis*-infected humans (such as *SsIR* and *NIE*) had L3i biased expression. Abundantly expressed L3i contigs of interest included *S. stercoralis* orthologs of cytochrome oxidase *ucr 2.1* and *hsp-90*, which may be potential chemotherapeutic targets. The *S. stercoralis* ortholog of fatty acid and retinol binding protein-1, successfully used in a vaccine against *Ancylostoma ceylanicum*, was identified among the 25 most highly expressed L3i genes. The sperm-containing glycoprotein domain, utilized in a vaccine against the nematode *Cooperia punctata*, was exclusively found in L3i biased genes and may be a valuable *S. stercoralis* target of interest.

Conclusions: A new DNA microarray tool for the examination of *S. stercoralis* biology has been developed and provides new and valuable insights regarding differences between infective and noninfective *S. stercoralis* larvae. Potential therapeutic and vaccine targets were identified for further study.

Citation: Ramanathan R, Varma S, Ribeiro JMC, Myers TG, Nolan TJ, et al. (2011) Microarray-Based Analysis of Differential Gene Expression between Infective and Noninfective Larvae of *Strongyloides stercoralis*. PLoS Negl Trop Dis 5(5): e1039. doi:10.1371/journal.pntd.0001039

Editor: Elodie Ghedin, University of Pittsburgh, United States of America

Received July 14, 2010; Accepted March 16, 2011; Published May 3, 2011

This is an open-access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the Creative Commons CC0 public domain dedication.

Funding: This work was supported by the Intramural Research Program of the Division of Intramural Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health and by NIH grants AI-050688, AI-022662, AI-082548, and RR02512. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: ramanathanr@niaid.nih.gov

Introduction

Strongyloides stercoralis is a parasitic nematode endemic to the tropics and subtropics that infects an estimated 30–100 million people worldwide. Chronically infected individuals have the potential to develop hyperinfection syndrome or disseminated disease, clinical entities that carry a very high (87–100%) mortality if unrecognized [1].

Free-living *S. stercoralis* infective third stage (L3i) larvae residing in the soil penetrate intact skin and blood vessels, ultimately developing to adults in the small intestine. Adult females, typically residing in the duodenum of the host, produce eggs by mitotic parthenogenesis that develop into first-stage (L1) larvae that are excreted into the stool. L1 larval progeny of parasitic females develop into free-living adults unless triggered by genetic, environmental, or host factors to develop directly into L3i larvae [2,3]. Despite sharing many characteristics, L1 and L3i larvae can be distinguished by their behavior and morphology. L1 larvae have a short, trilobed pharynx and expend much of their energy on feeding and growth [3]. L3i larvae, by contrast, can survive in

Author Summary

Strongyloides stercoralis is a soil-transmitted helminth that affects an estimated 30-100 million people worldwide. Chronically infected persons who are exposed to corticosteroids can develop disseminated disease, which carries a high mortality (87-100%) if untreated. Despite this, little is known about the fundamental biology of this parasite, including the features that enable infection. We developed the first DNA microarray for this parasite and used it to compare infective third-stage larvae (L3i) with noninfective first stage larvae (L1). Using this method, we identified 935 differentially expressed genes. Functional characterization of these genes revealed L3i biased expression of heat shock proteins and genes with products that have previously been shown to be immunoreactive in infected humans. Genes putatively involved in transcription were found to have L1 biased expression. Potential chemotherapeutic and vaccine targets such as far-1, ucr 2.1 and hsp-90 were identified for further study.

harsh environmental conditions, enabled by a comparatively thickened cuticle, constricted gastrointestinal tract, and closed mouth. These larvae are developmentally arrested, non-feeding, stress resistant, and long lived [3–5].

A high degree of specificity between these stages has been suggested by expressed sequence tag (EST) based analysis of free living L1 and L3i larvae for *S. stercoralis* [6–8]. These comparisons, however, are based on short reads of cDNA libraries and assumptions about abundance. There remain many unanswered questions about the basic molecular features underlying the apparent morphologic and behavioral differences between these larval stages. An improved understanding of these differences can provide insights into what defines infectivity and may ultimately prove useful in defining targets for the development of vaccines and therapeutics against this parasite.

In order to answer these questions, a DNA microarray tool for *S. stercoralis* – the species causing the vast majority of human infection worldwide - is needed. Although a DNA microarray has recently been developed for *Strongyloides ratti*, the natural parasite of brown rats (Rattus norvegicus) [9], previous work has suggested little conservation of gene expression profiles between these two species [10], underscoring the need for a DNA microarray specific to this species.

The availability of a S. stercoralis DNA microarray enables comparative analyses across nematodes, which can be utilized to further our understanding of the biologic determinants of parasitism. The free-living, non-parasitic, nematode C. elegans has been used as a model species for comparison with S. stercoralis. C. elegans dauer stage larvae and S. stercoralis L3i larvae share many morphologic and physiologic characteristics. The 'dauer hypothesis' recognizes these similarities and suggests that the same molecular genetic mechanisms control the morphogenesis of these stages [11]. Comparative genomics of gene expression based on EST abundance data for S. stercoralis suggests a higher degree of similarity between S. stercoralis L1 and C. elegans non-dauer expressed genes [6]. By contrast, a robust 'dauer-L3i expression signature' has not been found [6]. A comparative analysis based on microarray expression data for these species could prove useful not only in identifying a 'dauer-L3i expression signature' should it exist, but also in uncovering potentially significant determinants of S. stercoralis L3i infectivity.

The purpose of this study was to: 1) develop and optimize a DNA microarray tool for *S. stercoralis*, 2) utilize this microarray to

examine differences in gene expression between L3i and L1 larvae and 3) perform a comparative microarray analysis between parasitic *S. stercoralis* and non-parasitic *C. elegans* in order to develop further insights into the biologic determinants of parasitism.

Methods

Ethics statement

Animal handling and experimental procedures were undertaken in compliance with the University of Pennsylvania's Institutional Animal Care and Use Committee (IACUC) guidelines. Ethical approval was obtained for the study (protocol number 702342) from IACUC (University of Pennsylvania, Philadelphia, PA).

Parasites

All larvae used in this analysis were obtained from laboratory dogs infected with *S. stercoralis*, UPD strain [12]. Fecal samples from dogs were processed using the charcoal coproculture followed by Baermann funnel technique, as outlined elsewhere [13]. Post parasitic L1 larvae were recovered from freshly deposited stool samples; L3i larvae were recovered after 7 days of stool incubation at 25°C. L3i larvae underwent surface decontamination by migration through low-melting-point agarose. L1 larvae were decontaminated by 3 washes with phosphate buffered saline (PBS) containing an antibiotic cocktail. Decontaminated parasites were subsequently stored in Trizol reagent (Invitrogen, San Diego, CA) at -80° C. Using this method, 30,700 post-parasitic L1 and 50,000 L3i *larvae* were collected.

Isolation of total RNA from larvae

Total RNA was extracted by thawing pooled samples of L1 and L3i larvae at 37° C in a warm water bath and centrifuging the samples at 4° C ($805 \times$ g) for 10 minutes to obtain a pellet. The pellet was frozen in liquid nitrogen, ground thoroughly with an autoclaved mortar and pestle and then purified using an RNeasy mini kit (Qiagen, Valencia, CA) following the manufacturer's protocol. A Nano Drop-1000 spectrophotometer (NanoDrop Products, Wilmington DE) was used to determine the RNA concentration in each sample. RNA was more precisely quantified and quality assessed using the 2100 Bioanalyzer (Agilent, Santa Clara, CA).

Amplification and labeling

RNA samples from L1 and L3i stage larvae were co-hybridized using Cy3 and Cy5 labels to discriminate the relative level of target bound to the microarray probe. Fluorescent-labeled cDNA targets were prepared from total RNA using the Ovations amino-allyl kit (NuGEN, San Carlos, CA) according to the manufacturer's protocol. The kit utilizes an oligo dT primer for selective amplification of mRNA transcripts.

Hybridization procedure

Labeled samples were combined with blocking components poly(dA), yeast tRNA, and human Cot-1, in hybridization buffer composed of 25% formamide/5× saline-sodium citrate (SSC)/ 0.2% (w/v) sodium dodecyl sulfate (SDS) to a total volume of 60 μ l. After heating the sample (95°C for 3 minutes), it was centrifuged (20,000×g) for 3 minutes. Fifty eight μ l of the sample (1.6 μ g of labeled cDNA) was loaded onto the microarray chip. The microarray chips were hybridized overnight at 45°C using the MicroArray User Interface (MAUI) hybridization system (BioMicro Systems, Inc., Salt Lake City, UT). The following day, the chips were washed twice in 1×SSC/0.05% (w/v) SDS buffer

(3 minutes each wash) and twice in $0.1 \times$ SSC buffer (5 minutes each wash).

For the present study, four technical replicate experiments using pooled L1 and L3i larvae were performed, including one dye swap. The microarray chips were imaged using a GenePix 4000 B scanner (Molecular Devices, Sunnyvale, CA). Agilent Feature Extraction software was used for image analysis, protocol GE2-v5 10 Apr08. The data discussed in this publication have been deposited in the National Center for Biotechnical Information (NCBI) Gene Expression Omnibus (GEO) and are accessible through GEO Series accession number GSE24735 (http://www. ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE24735).

Microarray design

ESTs (11,335) were identified from L1 and L3i cDNA libraries created as part of the nematode EST project [6,7]. ESTs were organized into 3,571 contigs by bioinformatics analysis [14]. Oligonucleotide probes designed to hybridize with these contigs were used to develop early versions (V1 and V2) of chips manufactured by Combimatrix (Irvine, CA) based on a variety of algorithms for oligonucleotide design. Versions 1 and 2 were assessed for performance using RNA from L1 and L3i larvae. After testing the performance of these two versions of the arrays, an optimized version (V3) was developed. The best probe for each target was selected based on the average signal intensity for all arrays and the number of arrays with detectable signal. The spot density was 22K spots per array. Of the six oligonucleotides designed per target, one was designed using the Array Designer program (Premier Biosoft International, Palo Alto, CA), two were designed using E-Array (Agilent, Santa Clara, CA) using the "base composition" method (replicated twice), two were designed using E-array "best Tm" method, and the last was a 40-mer designed using Array Designer. Probes were selected to avoid crosshybridization to other sequences in the target (contig) dataset manufactured by Agilent SurePrint. The probes designed to make the V3 microarray are found in Table S1 in Supporting Information Text S1.

Functional annotation

All data were exported into the cDNA Annotation System (dCAS) [14,15]. This tool enabled annotation of each S. stercoralis contig based on Basic Local Alignment Search Tool (BLAST) alignments against multiple databases (NCBI nr protein database (NR), Gene Ontology (GO), euKaryotic Orthologous Groups (KOG), Pfam protein families database (PFAM), Simple Modular Architecture Research Tool (SMART), Wormbase (CELEG), and Saccharomyces genome database (YEAST) and provided the corresponding E-values. The database was also annotated manually with a composite categorization that summarized the findings across databases. The entire annotated database, with hyperlinks to the NIAID exon website, is accessible for download at: http://exon. niaid.nih.gov/transcriptome/S_stercoralis/SS-Supp-Web.zip. Α stand-alone version can also be accessed and downloaded at: http://exon.niaid.nih.gov/transcriptome/S_stercoralis/SS-Supp-StandAlone.zip. Extract the excel file and the links directory to your own computer for browsing the hyperlinks locally.

Statistical analysis

Spot values were calculated using a linear lowess dye normalization. Further, the 50th percentile of a set containing all the ribosomal genes in the array was applied to all spot values. In cases of multiple spots for the same *S. stercoralis* contig, the average of the \log_2 signal was calculated for each array. The mean signal ratio $(\log_2 L_{3i}/L_1)$ was calculated from the signals for all 4 arrays.

No surrogate values were applied. A single group *t*-test analysis was calculated on the data set. Variance shrinkage was not used when calculating p-values for differential expression. Differentially expressed genes were identified using a 'cutoff' of 2 fold expression difference or greater for $\log_2 \text{L3i/L1}$ signal ratios, and p < 0.01 for microarray signal data (false discovery rate (FDR) = 2.5%).

Functional analysis

A functional analysis was performed based on annotations provided by each database (Pfam, SMART, KOG, etc.). The number of genes per functional category (e.g. transcription, cytoskeleton, metabolism, etc.) was compared between L1 and L3i differentially expressed genes (as defined by the above cutoff). To ascertain whether genes belonging to certain functional classes were more likely to be highly expressed in one stage or another, we used a statistical test for one proportion using Normal approximation. Assuming a null proportion of 0.5 (i.e., that there is no difference in the number of genes of that category for the two classes), p values were calculated for deviation from 0.5 using Normal approximation. P values were adjusted for multiple comparisons using the Bonferroni criterion.

Gene-set enrichment analysis

Gene Set Enrichment Analysis (GSEA) is a robust method for analyzing molecular profiling data examines the clustering of a pre-defined group of genes (gene set) across the entire microarray database (all 3,571 contigs) in order to determine whether the gene set has biased expression in one larval stage versus another [16]. GSEA was used in this study to complement our use of single gene methods and determine whether S. stercoralis gene sets grouped according to various putative categories (for example, putative extracellular matrix genes) showed biased expression in either larval stage. For this analysis, the entire list of contigs on the microarray was sorted by mean log₂ L3i/L1 signal ratios. The distribution of genes from an a priori defined gene set throughout this ranked list was then determined using GSEA. Based on this distribution, the expression difference for each gene in the set is aggregated and a p-value for significance of the gene set as a whole is calculated using the Kolmogorov-Smirnoff test.

Gene sets were compiled by first downloading GO categories from Wormbase (www.wormbase.org) for *C. elegans* genes. Definitions for each GO category used can be found at http:// www.wormbase.org/db/ontology/gene. *S. stercoralis* orthologs for *C. elegans* genes were determined by dCAS based on BLAST alignments to the *C. elegans* gene. BLAST matches with E values>0.05 were excluded. Gene sets with fewer than 5 *S. stercoralis* contig matches were excluded from GSEA analysis. Using these criteria, 18 *S. stercoralis* gene sets were created (see Figure 1A). Additional manually compiled gene sets included the group of *S. stercoralis* genes whose products have been shown to be immunoreactive in humans infected with S. *stercoralis* [17–19], and a group of putatively identified heat shock proteins.

Comparative microarray analysis of *S. stercoralis* and *C. elegans*

Microarray expression data for *S. stercoralis* L3i and *C. elegans* dauer larvae were compared using several methods as follows: 1) We defined three gene sets comprising the *S. stercoralis* orthologs of "dauer-enriched" *C. elegans* genes derived from either *C. elegans* microarray expression data alone, both serial analysis of gene expression (SAGE) and microarray expression data or from the Gene Ontology category dauer larval development (Figure 1A) [20,21]. We then used GSEA to determine whether these gene sets

Gene Sets	Definition	ID*	Number of genes in gene set	Enrichment Score	Normalized Enrichment Score	Nominal P Value	FDR**
Immunoreactive	Manually compiled list of genes whose products have been shown to be immunoreactive in sera from S. stercoralis infected patients.	N/A	28	0.86	2.20	0.00	0.00
Heat Shock	List of genes encoding heat shock proteins based on annotation to euKaryotic Orthologous Groups (KOG) database.	N/A	40	0.53	1.40	0.03	0.56
Dauer-enriched (1)	A list of S. stercoralis orthologs of dauer-enriched C. elegans genes identified by microarray analysis [20].	N/A	43	0.50	1.29	0.10	0.77
Dauer-enriched (2)	A list of S. stercoralis orthologs of dauer-enriched C. elegans genes identified by microarray and SAGE analysis [20, 21].	N/A	15	0.41	0.96	0.54	1.00
Dauer larval development	The process whose specific outcome is the progression of the dauer larva over time, through the facultative diapause of the dauer (enduring) larval stage, with specialized traits adopted for dispersal and long-term survival, with elevated stress resistance and without feeding.	GO:0040024	7	0.61	1.24	0.25	0.76
Biosynthetic process	The chemical reactions and pathways resulting in the formation of substances; typically the energy requiring part of metabolism in which simpler substances are transformed into more complex ones.	GO:0009058	8	0.56	1.18	0.29	0.82
Proteolysis	The chemical reactions and pathways resulting in the breakdown of a protein by the destruction of the native, active configuration, with the hydrolysis, of pentide bonds.	GO:0006508	60	0.38	1.02	0.47	1.00
Ubiquitin dependent catabolic process	The chemical reactions and pathways resulting in the breakdown of a protein or peptide by hydrolysis of its peptide bonds, initiated by the covalent attachment of a ubiquitin moiety, or multiple ubiquitin moieties, to the protein.	GO:0008511	18	0.42	1.00	0.51	1.00
Glycolysis	The chemical reactions and pathways resulting in the breakdown of a monosoccharde (generally glucos) into pyruvake with the concomitant production of a small amount of ATP. Pyruvate may be converted to ethanol, lactate, or other small molecules, or fed into the TCA cycle.	GO:0006096	5	0.50	0.96	0.53	1.00
Lipid metabolic process	The chemical reactions and pathways involving lipids, compounds soluble in an organic solvert but not, or spanningly in an aquious solvent. Includes fatty acids, neutral fats, other fatty acid exters, and soaps, long chain (fatty) alcoholes and waxes, sphingolds and other long-chain bases, glycolapids, prosolubilids and sphingolidids, and caroteries, polyprends, sterois, terpenes and other isoprenoids.	GO:0006629	11	0.41	0.93	0.57	1.00
Negative regulation of multicellular organism growth	Any process that stops, prevents or reduces the frequency, rate or extent of growth of an organism to reach its usual body size.	GO:0040015	17	0.36	0.87	0.65	1.00
Oviposition	The deposition of eggs (either fertilized or not) upon a surface or into a medium such as water.	GO:0018991	59	0.30	0.81	0.81	1.00
Metabolic Process	The chemical reactions and pathways, including anabolism and catabolism, by which living organisms transform small molecules, but also include macromolecular processes such as DNA repair and replication, and protein synthesis and degradation.	GO:0008152	102	0.29	0.81	0.89	1.00
Growth	The increase in size or mass of an entire organism, a part of an organism or a cell.	GO:0040007	404	0.28	0.80	0.99	1.00
Defecation	The expulsion of feces from the recturn.	GO:0030421	6	0.39	0.79	0.74	0.96
Positive regulation of growth rate	Any process that increases the rate of growth of all or part of an organism.	GO:0040010	392	0.27	0.77	0.99	0.93
Transcription	The synthesis of either RNA on a template of DNA or DNA on a template of RNA.	GO:0006350	6	0.36	0.72	0.79	0.91
Negative regulation of vulval development	Any process that stops, prevents or reduces the trequency, rate or extent of development of the vulva. Vulval development is the process whose specific outcome is the progression of the ego-laying organ of famale and hermaphrolite nematodes sover time, from its formation to the mature structure. In nematodes, the vulva is formed from vental epidemal cells during larval stages to give rise to a fully formed vulva in the adult.	GO:0040027	24	0.18	0.44	0.98	0.99





Figure 1. Gene Set Enrichment Analysis and enrichment plot. A. Gene sets were compiled by listing the *S. stercoralis* orthologs of *C. elegans* genes assigned to Gene Ontology (GO) categories (downloaded from www.wormbase.org). Some gene sets were manually compiled. Only gene sets with at least 5 *S. stercoralis* contigs were included in this analysis. The results of the GSEA are listed for each gene set. The enrichment score reflects the degree to which each gene set is represented at the top or bottom of the list of 3,571 contigs ranked by fold change (L3i enriched = more positive, L1 enriched = more negative). The normalized enrichment score accounts for differences in gene set size and can be used to compare results across gene sets. The nominal p value estimates the statistical significance of the enrichment for a single gene set and does not correct for gene set size and multiple hypothesis testing. *ID = Gene Ontology Identification. **The False Discovery Rate (FDR) is adjusted for gene set size and multiple hypotheses testing. B. This plot depicts the distribution of individual genes (vertical black lines) encoding immunoreactive antigens recognized by sera from patients infected with *S. stercoralis*. This gene sets was analyzed against a list of 3,571 *S. stercoralis* contigs ranked by fold change of log₂ L3i/L1 mean signal ratios. The clustering of individual genes towards the left side of the list (above the red bar) suggests L3i-biased enrichment of this gene set. These genes are individually listed in Table S8 in Supporting Information Text S1.

showed significant L3i enrichment. 2) We examined whether a correlation exists between *C. elegans* dauer/L1 microarray expression data obtained by Wang and colleagues [20] with our *S. stercoralis* L3i/L1 microarray expression data. The previously

obtained *C. elegans* microarray expression data can be found at http://cmgm.stanford.edu/~kimlab/dauer/ExtraData.htm, Table S1 in Supporting Information Text S1, column "AdjD/ L1_Ratio" which corresponds to the average log₂ expression

. www.plosntds.org

values for *C. elegans* dauer larvae at time 0 relative to L1 larvae [20]. 3) Using these data, we calculated the absolute value of the difference between fold change values for *C. elegans* genes and their *S. stercoralis* orthologs (*C. elegans* dauer/L1 fold change - *S. stercoralis* L3i/L1 fold change). Only those genes with robust microarray expression data (p values<0.01) were included. In order to identify those genes that are expressed differently by *S. stercoralis*-L3i and *C. elegans* dauer larvae, a list was generated of all *S. stercoralis*-C. *elegans* orthologs with the greatest differences in fold change values (absolute value >2). The list was further narrowed to include only those *S. stercoralis*-C. *elegans* gene pairs where gene expression was regulated in opposite directions between the two nematodes (Table 1).

Microarray validation by quantitative real-time polymerase chain reaction (qPCR)

The sequences of L3i biased genes (contigs 24, 25, 65, 243, 2136) and L1 biased genes (contigs 55, 222, 387, 2328) were used to create primer-probe sets designed and manufactured by Applied Biosystems (Foster City, CA). The sequences for these

primer probes are listed in Table S2 in Supporting Information Text S1. The S. stercoralis control genes for qPCR analysis was S. stercoralis glyceraldehyde 3 phosphate dehydrogenase (GAPDH; GenBank accession number BI773092; contig_90; log₂L3i/ L1 = -0.28179). Post-parasitic L1 and L3i larvae (distinct from those hybridized onto the microarray) were collected and total RNA made as described above. Total RNA (1 µg) from L1 and L3i larvae was used to synthesize cDNA. qPCR was performed using all 9 primer probe sets in separate reactions with L1 cDNA and also with L3i cDNA. The reaction was performed using $10 \times$ RT buffer (10 µl), 25 mM MgCl2 (22 µl), dNTP (20 µl), random hexamers (5 μ l), RNase inhibitor (2 μ l), and multiscribe reverse transcriptase (50 U/µl; 6.25 µl) in a microamp 96-well reaction plate (Applied Biosystems). De-ionized, distilled water was added to total volume of 65.25 µl. Cycling conditions were: 25°C for 10 minutes, 37°C for 60 minutes, 95°C for 5 minutes, then 4.0°C. Each experiment was performed in triplicate. The mean negative delta threshold cycle (delta C_T) was calculated for each sample. The data generated by performing qPCR using primer probes for 9 contigs on L1 and L3i cDNA (n = 18) was plotted

Table 1. Differences between 5. stercoralis (L3i/L1) and C. elegans (dauer/L1) gene expression profiles^a.

			Fold change	e direction ^b	Fold chang	je	
<i>C. elegans</i> match	<i>S.stercoralis</i> contig	Putative identification	<i>C. elegans</i> (dauer/L1)	S. stercoralis (L3i/L1)	C. elegans	S. stercoralis	Absolute difference ^c
C13D9.8	2626	псх-9	↑	\downarrow	3.59	0.39	3.20
T14D7.2	2474	protein_id:CAB03365	↑	\downarrow	21.75	0.40	21.36
C06B3.4	2873	stdh-1 estradiol 17 beta-dehydrogenase	↑	\downarrow	3.71	0.44	3.27
T02D1.5	1790	pmp-4 ABC transporters	↑	\downarrow	16.57	0.45	16.12
T19B10.2	504	protein_id:CAA98547	↑	\downarrow	2.56	0.55	2.01
T04B2.5	2003	protein_id:CAA92628	↑	\downarrow	3.97	0.56	3.41
F46C3.1	1269	<i>pek-1</i> eukaryotic translation initiation factor 2 alpha kinase PEK	Ŷ	Ļ	3.61	0.57	3.04
F19H8.1	1652	tps-2 trehalose phosphate synthase	↑	\downarrow	4.22	0.65	3.57
F10B5.3	1998	Zinc finger, C2H2 type	↑	\downarrow	4.60	0.66	3.94
Y57G11C.15	850	protein transport protein SEC61 alpha subunit	↑	\downarrow	2.92	0.76	2.16
F42D1.2	118	tyrosine aminotransferase	Ļ	↑	0.17	2.22	2.05
F42D1.2	117	tyrosine aminotransferase	Ļ	↑	0.17	2.87	2.69
C53B4.5	2200	col-119 collagen	Ļ	↑	0.10	2.98	2.88
F28C1.2	836	egl-10 G-protein beta subunit GPB-2	\downarrow	↑	0.78	2.99	2.21
B0491.5	291	protein_id:CAA90087	\downarrow	1	0.84	3.18	2.34
E02H1.7	3417	nhr-19 Zinc finger, C4 type (two domains)	Ļ	↑	0.35	3.25	2.90
C02F12.7	64	tag-278	\downarrow	↑	0.27	3.89	3.62
C53B4.5	2267	col-119 collagen	\downarrow	↑	0.10	4.11	4.01
E02H1.7	1846	nhr-19 Zinc finger, C4 type (two domains)	\downarrow	1	0.35	4.18	3.83
C03B1.12	785	Imp-1	Ļ	↑	0.70	5.09	4.39
ZK863.2	9	col-37 collagen status	Ļ	↑	0.11	6.74	6.63
B0365.3	220	eat-6 Na(+)/K(+) ATPase alpha subunit	\downarrow	↑	0.71	6.83	6.12
B0495.2	358	CDC2 status	\downarrow	↑	0.71	8.28	7.58
F22B3.4	427	nmy-2 myosin heavy chain status	\downarrow	\uparrow	0.89	8.33	7.44
F42D1.2	116	tyrosine aminotransferase	\downarrow	\uparrow	0.17	9.04	8.87

^aShown only are *S. stercoralis-C. elegans* orthologs with an absolute difference >2. All *S. stercoralis-C. elegans* orthologs are BLAST matches with an E value<0.05. The p value for *S. stercoralis* microarray signal data was <0.01.

^bArrows indicate whether genes had increased () or decreased () expression in *C. elegans* dauer or *S. stercoralis* L3i larvae relative to its respective L1 stage.

^cThe values in this column were calculated by taking the absolute value of [fold change *C. elegans* dauer/L1 - fold change *S. stercoralis* L3i/L1]. *C. elegans* expression data were previously obtained by Wang and colleagues [20].

doi:10.1371/journal.pntd.0001039.t001

against the average L1 and L3i intensity signals for each gene (Figure 2).

Results

Identification of differentially expressed genes

A total of 3,571 distinct contigs were studied by this microarray analysis (Table S3 in Supplemental Information Text S1). Using pre-defined cutoffs, 935 contigs were identified as differentially expressed as shown in the volcano plot (Figure 3). Of these, 466 genes were L1 biased (Table S4 in Supporting Information Text S1) and 469 genes were L3i biased (Table S5 in Supporting Information Text S1). Among the 25 most highly expressed L3i genes were the S. stercoralis orthologs of fatty acid/retinol binding protein-1 (contig 1151; 11 fold expression difference), a ferritin chain homolog (contig 94; 14 fold expression difference), and one of four putative trehalases (contig 68; 14-fold expression difference). Among the 25 most highly expressed L1 genes were electron transport chain proteins such as NADH dehydrogenase (contig 371; 0.13-fold change); cytochrome b (contig 2328; 0.19 fold change) and cytochrome c oxidase subunit 1 (contig 55; 0.29 fold change). The 25 most highly expressed L1 or L3i genes are listed in Table S6 in Supporting Information Text S1.

Functional analysis of L1 and L3i biased genes

A greater number of L1 (n = 40) than L3i biased (n = 18) genes were putatively involved in transcription (p = 0.004, not Bonferroni adjusted; see Figure 4A,B). A complete listing of these genes is shown in Figure 4B. This finding was also noted in an analysis of classifications based on GO categories (p = 0.01 for 'transcription'), and manual annotations (p = 0.007 for 'transcription machinery'), although p values were not <0.05 when Bonferroni-adjusted for multiple comparisons. BLAST matches to SMART and Pfam databases both indicated that the sperm-containing glycoprotein (SCP) domain was found exclusively in the L3i-group (n = 13)



Figure 2. Correlation between microarray signal data and quantitative PCR data. Quantitative PCR was performed using primer probe sets designed from abundantly expressed L1 and L3i S. stercoralis contig sequences (L1 biased contigs 55, 222, 387, 2328; L3i biased contigs_24, 25, 65, 243, 2136) with cDNA synthesized from L1 and L3i larvae (n = 18). Each data point is the calculated negative delta C_T (sample C_T minus control C_T) for the mean of 3 replicates. These data are plotted against the corresponding L1 or L3i average intensity microarray signal. A positive correlation was found (Spearman rank = 0.4778; p = 0.0449).

doi:10.1371/journal.pntd.0001039.g002



Figure 3. Volcano plot used in differential expression analysis. The *x*-axis is log_2 ratio of gene expression levels between two stages; the *y*-axis is adjusted p value based on $-log_{10}$. The colored dots (L1 = green) and right (L3i = red) represent the differentially expressed genes based on p < 0.01 (False Discovery Rate = 2.5%; represented by black horizontal line) and 2-fold expression difference (represented by two black vertical lines). doi:10.1371/journal.pntd.0001039.g003

genes; see Table S7 in Supporting Information Text S1 for the complete list; p value based on matches to Pfam = 0.003, Bonferroni-adjusted for multiple comparisons).

Of the entire 3,571 contigs, 1,351 *S. stercoralis* genes (37.8%) were of unknown function (manual annotation).

GSEA

S. stercoralis orthologs were matched to 35 sets of C. elegans genes grouped by various categories (e.g. negative regulation of vulval induction, oviposition, heat shock proteins, etc.). Eighteen of 35 gene sets queried met criteria for inclusion into the GSEA analysis (based on minimum size of 5 genes; see Figure 1A). Of these 18 gene sets, only 2 gene sets were significantly enriched in the L3i phenotype at nominal p value<5%. The most significantly enriched genes were those with immunoreactive gene products recognized by sera from infected individuals (Figure 1B; nominal p-value<0.0001; FDR<0.0001). Heat shock proteins were the next most highly enriched (nominal p value = 0.034, FDR = 0.56). For an annotated list of the individual genes enriched in each of these categories, refer to Tables S8 and S9 in Supporting Information Text S1. None of the 18 gene sets were enriched in the L1 phenotype.

Comparative microarray analysis of *S. stercoralis* and *C. elegans*

Four hundred and twenty two of 3,571 *S. stercoralis* contigs had *C. elegans* orthologs for which robust microarray signal data were available. When *C. elegans* and *S. stercoralis* microarray signals were plotted against each other, a poor and non-significant correlation was found (Spearman rank = 0.06; p = 0.2444, graph not shown). No significant L3i enrichment of *S. stercoralis* orthologs of *C. elegans* 'dauer enriched' genes was found by GSEA (nominal p-value = 0.10). On the contrary, 25 orthologs expressed in opposite directions by dauer and L3i larvae relative to their respective L1 stage larvae were identified (see Table 1).

Correlation between EST and microarray data

A statistically significant positive correlation was found between microarray expression data and EST abundance data (p < 0.0001; max $R^2 = 0.26$; graph not shown).



B. S. stercoralis L1 and L3i Biased Genes Putatively Involved in Transcription

L1 Biased Genes	
Contig Number	Putative Identification
592	Transcription factor PBX and related HOX domain proteins
	Transcription initiation factor TFIID, subunit
686	BDF1 and related bromodomain proteins
891	Apoptosis antagonizing transcription factor/protein transport protein
1020	RNA polymerase II, large subunit
1036	Transcription factor MEIS1 and related HOX
1000	domain proteins Transcription initiation factor TFIID, subunit
1037	TAF7
1100	Hepatocyte nuclear factor 4 and similar steroid hormone receptors
1262	Dosage compensation complex, subunit MLE
1277	bHLH transcription factor
1351	Thyroid hormone receptor-associated protein
1397	Pterin carbinolamine dehydratase
	Zn2+-binding dehvdrogenase (nuclear
1462	receptor binding factor-1)
1505	RNA polymerase I, second largest subunit
1613	Hepatocyte nuclear factor 4 and similar steroid hormone receptors
1765	Positive cofactor 2 (PC2), subunit of a multiprotein coactivator of RNA polymerase II
1794	Transcription factor
1883	steroid hormone receptors
1905	MOT2 transcription factor
2023	Transcription factor LIM3, contains LIM and HOX domains
2035	C2H2-type Zn-finger protein
2085	CAATT-binding transcription factor/60S ribosomal subunit biogenesis protein
2099	TBP-binding protein, activator of basal
2163	Alpha-1,2 glucosyltransferase/ transcriptional
	Transcription factor NERF and related
2210	proteins, contain ETS domain
2338	superfamily)
2366	Casein kinase II, alpha subunit
2446	Transcription factor TCF20
2460	Nuclear receptors of the nerve growth factor- induced protein B type
2511	Alpha-1,2 glucosyltransferase/ transcriptional activator
2553	Nuclear hormone receptor beta FTZ-F1
2707	Component of histone deacetylase complex (breast carcinoma metastasis suppressor 1 protein in humans)
2757	Transcriptional coactivator
2786	RNA polymerase I, large subunit
2843	TBX2 and related T-box transcription factors
2917	RNA polymerase I, second largest subunit
2943	Activating signal cointegrator 1
3218	RNA Polymerase C (III) 37 kDa subunit
3245	Transcription initiation factor TFIID, subunit TAF6 (also component of historie scabilizant/cransforme, SAC 61)
	MADS how transcription factor
3337	Transcription regulator dachshund, contains
3359	SKI/SNO domain

Contig Number	Putative Identification
156	Transcription regulator XNP/ATRX, DEAD- box superfamily
167	Transcription factor containing NAC and TS N domains
206	RNA polymerase II general transcription factor BTF3 and related proteins
333	Calcium-responsive transcription coactivator
445	Transcriptional regulator
446	Transcriptional regulator
519	Transcription factor GT-2 and related proteins, contains trihelix DNA- binding/SANT domain
770	C2H2-type Zn-finger protein
836	Transcription factor
1388	Ovo and related transcription factors
1478	Predicted forkhead transcription factor
1513	RNA polymerase II, large subunit
1521	Predicted forkhead transcription factor
1719	Transcription factor TMF, TATA element modulatory factor
3091	DNA repair/transcription protein Mms19
3324	Transcriptional coactivator CAPER (RRM superfamily)
3509	Apoptosis antagonizing transcription factor/protein transport protein
3530	bZIP transcription factor NRF1

Figure 4. Functional analysis of differentially expressed *S. stercoralis* L1 and L3i genes. Horizontal bar graph (A) depicting the number of contigs per functional category in each stage (x axis), according to BLAST matches to the KOG database (y-axis). BLAST matches with E values>0.05 were excluded. The strongest difference was found between L1 and L3i genes putatively involved in transcription (p = 0.004). L1 = blue; L3i = red. The table (B) lists the L1 and L3i biased *Strongyloides* contigs putatively classified under "transcription." doi:10.1371/journal.pntd.0001039.g004

Validation of microarray data with qPCR

A positive correlation was found (Spearman rank = 0.4778; p = 0.0449) between average L1 or L3i microarray intensity signals and mean negative delta C_T of qPCR (Figure 2).

Discussion

In this microarray based analysis of differential gene expression between infective and noninfective *S. stercoralis* larvae, we uncovered differences in the expression of genes putatively encoding transcription factors, heat shock proteins and antigens known to be immunoreactive in sera from infected humans. A comparative microarray analysis of our data revealed several differences between *S. stercoralis* L3i and *C. elegans* dauer stage larvae, such as in the expression of genes putatively encoding collagen and myosin. Potential therapeutic and vaccine targets were identified for further study.

L1 larvae appear to be transcriptionally more active

Analogous to their non-dauer C. elegans counterparts, actively growing S. stercoralis L1 larvae are thought to have higher rates of transcription relative to L3i-stage larvae. This supposition is based on comparisons between C. elegans non-dauer biased genes and S. stercoralis L1-biased genes that suggest transcriptional conservation of genes involved in early larval growth [6]. Consistent with this finding, we found L1 biased expression of genes putatively involved in transcription. Among the S. stercoralis L1-biased genes involved in transcription were transcription initiation factors (contigs 3245, 1037, 686), transcription factors (contigs 1905, 1277, 891, 2023, 2446, 1036, 1794, 592, 2210), and subunits of RNA polymerase (contigs 1505, 3218, 1020, 2917). By contrast, the L3i-biased genes involved in transcription though fewer, included transcriptional regulators (contigs 446, 445, 156) as well as transcription factors (contigs 1521, 519, 836, 167, 1478), implying that L3i larvae are not transcriptionally inactive and may regulate transcription differently. This would be consistent with what is known of C. elegans dauer larvae, which express distinct sets of dauer-specific genes at certain time points (dauer exit, for example) [20,21].

L3i biased expression of genes with products that have been shown to be immunoreactive in *S. stercoralis*-infected humans

Not surprisingly, genes encoding *S. stercoralis* antigens known to produce robust antibody responses in infected humans were found to have L3i biased expression by GSEA [17–19]. Two of these genes, IgG immunoreactive antigen (SsIR) and NIE antigen, have been recently employed in serodiagnostic assays with some advantage over crude antigen [19]. The finding that genes with products capable of inducing protective immunity demonstrate stage-biased gene expression supports the further investigation of these genes as vaccine candidates.

Heat shock proteins have been shown to play a critical role in determining parasite survival during stressful conditions because they can bind denatured or misfolded proteins [22,23]. Biased expression of genes encoding heat shock proteins in the *S. stercoralis* L3i relative to L1 larvae, as suggested by GSEA, is consistent with this role. *Hsp-90* in particular has been identified as a parasitism-

central gene based on changes in *S. ratti* gene expression during high immune pressure [22] and is similarly abundantly expressed by *S. stercoralis* L3i larvae.

Sperm containing glycoprotein (SCP) domain exclusively found in L3i larvae

The SCP domain, found exclusively in L3i biased genes, is a conserved domain of unknown function present in a wide range of organisms [24]. Interestingly, it has been found to be present in activation-associated secreted proteins that have been studied as potential vaccine targets in other nematodes [24,25]. Whether overrepresentation of the SCP domain in the L3i group is related to the presence of these secreted proteins is unclear, but activation-associated secreted proteins have been found to be important in many parasitic nematodes in which they have been studied to date.

C. elegans dauer and *S. stercoralis* L3i larvae have distinct characteristics

Consistent with previous findings, a striking L3i-C. elegans 'dauer expression signature' was not uncovered in this comparative microarray analysis [6]. We instead identified genes that are regulated in apparently opposite manners by C. elegans dauer and S. stercoralis L3i larvae which offer useful clues about the biology of S. stercoralis parasitism. L3i biased expression of the putative nmy-2 gene (encoding the myosin heavy chain) is consistent with the highly motile nature of L3i larvae which, unlike their dauer counterparts, seek out and initiate infection in a host. Although dauer and L3i larvae both contain a cuticle that enables survival in the environment, the parasitic cuticle has been associated with the ability of infective stages to evade the immune response of the host, and its structure varies from one species to another [26]. Biased expression of genes putatively encoding particular collagens (col-37, col-119) in the L3i but not the C. elegans dauer, points to differences in the composition of the parasitic cuticle that could potentially have a role in this regard. In fact, a recent microarray based analysis of the response of the S. ratti transcriptome to host immunologic environment notes upregulation of collagen genes by S. ratti which is believed to play a protective role for the parasite [27].

C. elegans dauer and *S. stercoralis* L3i larvae can survive in the environment even in the absence of a steady source of food. One way by which this occurs is by the development of electron-dense intestinal granules that store non-lipid products [11]. The gene *lmp-1* plays an essential role in this regard for dauer larvae as suggested by RNA interference studies [28]. It is likely that L3i larvae similarly utilize these granules while in the environment. The presence of these granules may additionally explain the darkened color of the radially constricted intestines of L3i larvae, an appearance shared by its dauer counterpart.

A key feature shared by dauer and L3i larvae is the ability to extend the lifespan while in the free-living state. In both *C. elegans* and *S. stercoralis*, the forkhead transcription factor DAF-16 plays a role in regulating dauer diapause, longevity and metabolism [11,29,30]. A downstream target of DAF-16, *egl-10*, is known to be negatively regulated by DAF-16 in *C. elegans* [29]. By contrast, this gene was found to have biased L3i larval expression in *S. stercoralis*. Such discordance is consistent with findings from a prior study that failed to detect a transcriptional profile typical of down-regulated

insulin-like signaling in long-lived parasitic females of S. ratti [31]. Although the downstream targets of insulin-like signaling have not been fully elucidated in Strongyloides species, the apparent upregulation of Ss-egl-10 in the L3i potentially highlights adaptations at a molecular level that likely underlie the evolution to parasitism. Such adaptations could include alterations in genes controlling metabolic and developmental functions, adaptations of pre-existing genes to encode new functions, and gene duplication and diversification [32]. The apparent lack of a C. elegans dauer-like transcriptional profile in S. stercoralis L3i is also consistent with published findings on the expression of transcripts encoding the orthologs of DAF-7 in this parasite [33] and in S. ratti and Parastrongyloides trichosuri [34]. DAF-7 is the ligand that activates TGF-β-like signaling and thereby promotes continuous (i.e. non-dauer) development in C. elegans. Its expression is biased towards C. elegans first-stage larvae fated for continuous development rather than dauer third-stage larvae [34,35]. By contrast, messages encoding DAF-7 orthologs in S. stercoralis, S. ratti and P. trichosuri all show biased expression in the L3i, which has been characterized heretofore as dauer-like [33,34]. These facts notwithstanding, outright rejection of the 'dauer hypothesis' of developmental regulation in the L3i of parasitic nematodes on the basic of transcriptional data alone is likely to be premature [36]. It is particularly noteworthy in this regard that key signal transducing elements such as DAF-16 that directly regulate C. elegans dauer development are constitutively transcribed and their functions governed not at the transcriptional level but rather by posttranslational modifications such as phosphorylation [37,38].

The true value in identifying these and other genetic determinants of *S. stercoralis* parasitism lies in whether the products of these genes can induce protective immunity. Indeed, one of the genes identified in our list, the *S. stercoralis* ortholog of *eat-6 Na+k+ATPase*, has already been identified as a potential vaccine candidate based on animal experiments [39].

Additional therapeutic targets and immunodiagnostic genes of significance

Contig 1872, a gene with L3i biased expression, encodes an ortholog of *C. elegans* core subunit of the cytochrome bcl complex, UCR 2.1 (E-value = 1E-014). This subunit has been shown to be a potential target for antiparasitic drugs based on the finding that in *C. elegans*, UCR 2.1 is essential for viability and is less related to mammalian UCR-1 than to mitochondrial processing peptidases from other organisms [40]. *S. stercoralis* transgenesis experiments [41] may prove useful in investigating the question of whether this gene is similarly essential for *S. stercoralis* larval survival.

In our microarray analysis of *S. stercoralis*, we found abundant L3i expression of the *S. stercoralis* ortholog of *hsp-90*, contig_77 (3 fold expression difference). Interestingly, the *hsp-90* inhibitor geldanamycin has been shown to have a macrofilaricidal effect on filarial nematode *Brugia pahangi* [42]. *Hsp-90* has been identified among *S. ratti* parasitism central genes critical for survival and further studies investigating it as a chemotherapeutic target are warranted.

Contig 1151, which was among the 25 most highly biased L3i genes (11-fold expression difference), corresponds to fatty acid and retinol binding protein-1 (FAR-1; E-value = 1E-016). FAR-like proteins are major secreted products of parasitic nematodes that allow the parasite to scavenge essential nutrients from its host [43]. Depletion of host lipids is thought to be necessary for parasite survival and may additionally impair the host immune response [44]. These proteins have additionally demonstrated stage and gender specificity in other nematodes, most notably in the

hookworm Ancylostoma ceylanicum [45]. The immunodiagnostic potential of FAR-like proteins has been assessed in other nematodes, such as Onchocerca volvulus, in a serologic assay based on Ov-20 (FAR-1) [45,46,47]. FAR-1 proteins have been successfully used in a vaccine in animals infected with A.ceylanicum [45]. These microarray data identify S. stercoralis far-1 as an L3i-biased target that may be a potential vaccine candidate or immunodiagnostic antigen.

Limitations

Approximately one-third of S. stercoralis genes are of unknown function. This finding is consistent with a previous EST analysis that revealed a similar percentage (25%) of S. stercoralis clusters with no significant BLAST alignments [8]. This finding is also consistent with functional genomics analyses of the C. elegans and human genomes where significant numbers of genes of unknown function were identified [48,49]. Some of these unknown sequences may derive from 3 untranslated mRNA regions, which are common in polydT-primed libraries [50]. The complete genome sequence of S. stercoralis is not available to date. Inferred functional annotations of an analogous nematode C. elegans, while useful, may not be directly applicable to S. stercoralis, as suggested by interspecies differences uncovered in the present comparative microarray analysis. Because a number of C. elegans genes did not have S. stercoralis orthologs that were also differentially expressed according to our predefined 'cutoffs,' it was difficult to formulate gene lists organized into functional categories with at least 5 contigs. This limited our ability to analyze biochemical or metabolic pathways of potential importance. As our knowledge of the S. stercoralis genome increases, these microarray analyses will likely gain in usefulness and a more direct approach using annotation based on known S. stercoralis gene functions would be even more informative.

Conclusions

DNA microarrays allow for simultaneous analysis of large numbers of genes from two or more biologic conditions. This powerful method of analysis has revolutionized our understanding of the immunopathogenesis of schistosomiasis [51], for example, and has advanced the development of vaccine discovery and therapeutics in parasitology [52,53]. Until now, studies of *S. stercoralis* have been limited to the analysis of ESTs rather than the full genome sequence. Development of a novel DNA microarray tool for the study of *S. stercoralis* represents an exciting step forward in our understanding of this parasite.

Supporting Information

Text S1 This file contains supplemental information regarding microarray probe information (Table S1), primer probe sequences used in real-time PCR analysis (Table S2), all contigs (Table S3), L1 biased contigs (Table S4), L3i biased contigs (Table S5), most highly expressed L1 and L3i contigs (Table S6), L3i biased contigs containing sperm containing glycoprotein domain (Table S7), and results of the GSEA for immunoreactive genes (Table S8) and heat shock proteins (Table S9). For the column marked "Manual Annotation," the following abbreviations were used: em = energy metabolism; extmat = extracellular matrix; cs = cytoskeleton; imm = genes encoding antigens known to be immunoreactive in sera from patients infected with *S. stercoralis*; met = metabolism; nr = nuclear regulation; pe = protein export machinery; pm = protein modification; prot = proteasome machinery; ps = protein synthesis; st = signal transduction; tf = transcription factor;

tm = transcription machinery; tr = transporters and storage proteins; unk = unknown.

Found at: doi:10.1371/journal.pntd.0001039.s001 (8.05 MB XLSX)

Acknowledgments

We thank Guojian Jiang for his expert assistance with microarray hybridizations and NIAID intramural editor Brenda Rae Marshall for assistance with the manuscript preparation.

References

- Ramanathan R, Nutman T (2008) Strongyloides stercoralis infection in the immunocompromised host. Curr Infect Dis Rep 10: 105–110.
- Harvey SC, Gemmill AW, Read AF, Viney ME (2000) The control of morph development in the parasitic nematode *Strongyloides ratti*. Proc R Soc Lond B Biol Sci 267: 2057–2063.
- Speare R (1989) Identification of species of *Strongyloides*. In: Grove DI, ed. Strongyloidiasis a major roundworm infection of man. London: Taylor and Francis. pp 11–84.
- Cassada RC, Russell RL (1975) The dauerlarva, a post-embryonic developmental variant of the nematode *Caenorhabditis elegans*. Dev Biol 46: 326–3242.
- Klass M, Hirsh D (1976) Non-ageing developmental variant of *Caenorhabditis* elegans. Nature 260: 523–525.
- Mitreva M, McCarter JP, Martin J, Dante M, Wylie T, et al. (2004) Comparative genomics of gene expression in the parasitic and free-living nematodes *Strongyloides stercoralis* and *Caenorhabditis elegans*. Genome Res 14: 209–220.
- Moore TA, Ramachandran S, Gam AA, Neva FA, Lu W, et al. (1996) Identification of novel sequences and codon usage in *Strongyloides stercoralis*. Mol Biochem Parasitol 79: 243–248.
- Thompson FJ, Mitreva M, Barker GLA, Martin J, Waterston RH, et al. (2005) An expressed sequence tag analysis of the life-cycle of the parasitic nematode *Strongyloides ratti*. Mol Biochem Parasitol 142: 32–46.
- Evans H, Mello LV, Fang Y, Wit E, Thompson FJ, et al. (2008) Microarray analysis of gender- and parasite-specific gene transcription in *Strongyloides ratti*. Int J Parasitol 38: 1329–1341.
- Thompson FJ, Barker GLA, Hughes L, Wilkes CP, Coghill J, et al. (2006) A microarray analysis of gene expression in the free-living stages of the parasitic nematode *Strongyloides ratti*. BMC Genomics 7: 157–177.
- Castelletto ML, Massey HC, Jr., Lok JB (2009) Morphogenesis of Strongyloides stercoralis infective larvae requires the DAF-16 ortholog FKTF-1. PLoS Pathog 5: e1000370.
- Schad GA, Hellman ME, Muncey DW (1984) Strongyloides stercoralis: hyperinfection in immunosuppressed dogs. Exp Parasitol 57: 287–296.
- Lok JB (2007) Strongyloides stercoralis: a model for translational research on parasitic nematode biology. In: The C. elegans Research Community, Worm-Book, doi/10.1895/wormbook.1.134.1, http://www.wormbook.org.
- Guo Y, Ribeiro JM, Anderson JM, Bour S (2009) dCAS: a desktop application for cDNA sequence annotation. Bioinformatics 25: 1195–1196.
- Ribeiro JM, Topalis P, Louis C (2004) AnoXcel: an Anopheles gambiae protein database. Insect Mol Biol 13: 449–357.
- Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL (2005) Gene set enrichment analysis: a knowledge-based approach for interpreting genomewide expression profiles. Proc Natl Acad Sci USA 102: 15545–15550.
- Ravi V, Ramachandran S, Thompson RW, Andersen JF, Neva FA (2002) Characterization of a recombinant immunodiagnostic antigen (NIE) from Strongyloides stercoralis L3i-stage larvae. Mol Biochem Parasitol 125(1–2): 73–81.
- Ramachandran S, Thompson RW, Gam AA, Neva FA (1998) Recombinant cDNA clones for immunodiagnosis of strongyloidiasis. J Infect Dis 177: 196–203.
- Ramanathan R, Burbelo PD, Groot S, Iadarola MJ, Neva FA, et al. (2008) A luciferase immunoprecipitation systems assay enhances the sensitivity and specificity of diagnosis of *Strongyloides stercoralis* infection. J Infect Dis 198: 444–451.
- Wang J, Kim SK (2003) Global analysis of dauer gene expression in *Caenorhabditis elegans*. Development 130: 1621–1634.
- Jones S, Riddle DL, Pouzyrev AT (2001) Changes in gene expression associated with developmental arrest and longevity in *Caenorhabditis elegans*. Genome Res 11: 1346–1352.
- Thompson FJ, Barker GL, Hughes L, Viney ME (2008) Genes important in the parasitic life of the nematode *Strongyloides ratti*. Mol Biochem Parasitol 158: 112–119.
- Morimoto RI (1998) Regulation of the heat shock transcriptional response: cross talk between a family of heat shock factors, molecular chaperones, and negative regulators. Genes Dev 12: 3788–3796.
- Yatsuda AP, Eysker M, Vieira-Bressan MC, De Vries E (2002) A family of activation-associated secreted protein (ASP) homologues of *Cooperia punctata*. Res Vet Sci 73: 297–306.
- Hawdon JM, Jones BF, Hoffman DR, Hotez PJ (1996) Cloning and characterization of *Ancylostoma*-secreted protein. A novel protein associated with the transition to parasitism by infective hookworm larvae. J Biol Chem 271: 6672–6678.

Author Contributions

Conceived and designed the experiments: DA JBL TBN. Performed the experiments: RR TGM TJN DA JBL TBN. Analyzed the data: RR SV TGM TJN DA JBL TBN. Contributed reagents/materials/analysis tools: SV JMCR TGM TJN DA JBL TBN. Wrote the paper: RR SV TGM TJN JBL TBN.

- Martinez A, De Souza W (1996) A Freeze-Fracture and Deep Etch Study of the Cuticle and Hypodermis of Infective Larvae of *Strongyloides venezuelensis* (Nematoda). Int Journal for Parasitol 27: 289–297.
- O'Meara H, Barber R, Mello LV, Sangaralingam A, Viney ME, Paterson S (2010) Response of the *Strongyloides ratti* transcriptome to host immunological environment. Int J Parasitol;Doi:10.1016/j.ijpara.2010.06.005.
- Kostich M, Fire A, Fambrough DM (2000) Identification and molecular-genetic characterization of a LAMP/CD68-like protein from *Caenorhabditis elegans*. J Cell Sci 113(Pt 14): 595–606.
- Oh SW, Mukhopadhyay A, Dixit BL, Raha TR, Green MR, et al. (2005) Identification of direct DAF-16 targets controlling longevity, metabolism and diapause by chromatin immunoprecipitation. Nat Genet 38: 251–257.
- Massey HC, Jr., Bhopale MK, Li X, Castelletto M, Lok JB (2006) The fork head transcription factor FKTF-1b from *Strongyloides stercoralis* restores DAF-16 developmental function to mutant *Caenorhabditis elegans*. Int J Parasitol 39: 1561–1571.
- Thompson FJ, Barker GL, Nolan T, Gems D, Viney ME, Mech Ageing Dev (2009) Transcript profiles of long and short- lived adults implicate protein synthesis in evolved differences in ageing in the Nematode *Strongyloides ratti*. 130: 167–72.
- Mitreva M, Blaxter M, Bird D, McCarter J (2005) Comparative genomics of nematodes. Trends in Genetics 10: 573–581.
- Massey HC, Castelletto M, Bhopale VM, Schad G, Lok JB (2005) Sst-tgh-1 from Strongyloides stercoralis encodes a proposed ortholog of daf-7 in Caenorhabditis elegans. Molecular and Biochemical Parasitology 142: 116–120.
- Crook M, Thompson FJ, Grant WN, Viney ME (2005) daf-7 and the development of Strongyloides ratti and Parastrongyloides trichosuri. Mol Biochem Parasitol 139: 213–223.
- Ren P, Lim CS, Johnsen R, Albert PS, Pilgrim D, et al. (1996) Control of C. elegans larval development by neuronal expression of a TGF-beta homolong. Science 274: 1389–1391.
- Hotez P, Hawdon JM, Schad GA (1993) Hookworm larval infectivity, arrest and amphiparatenesis: the *Caenorhabditis elegans* daf-c paradigm. Parasitology Today 9: 23–26.
- Lee RY, Hench J, Ruvkun G (2001) Regulation of *C. elegans* DAF-16 and its human ortholog FKHRL1 by the *daf-2* insulin-like signaling pathway. Curr Biol 11: 1950–1957.
- Cahill CM, Tzivion G, Nasrin N, Ogg S, Dore J, et al. (2001) Phosphatidylinositol 3-kinase signaling inhibits DAF-16 DNA binding and function via 14-3-3-dependent and 14-3-3-independent pathways. J Biol Chem 276: 13402–13410.
- Kerepesi LA, Keiser PB, Nolan TJ, Schad GA, Abraham D, et al. (2005) DNA immunization with Na⁺-K⁺ ATPase (Sseat-6) induces protective immunity to larval *Strongyloides stercoralis* in mice. Infect Immun 73: 2298–2305.
- Nomura H, Athauda SB, Wada H, Maruyama Y, Takahashi K, et al. (2006) Identification and reverse genetic analysis of mitochondrial processing peptidase and the core protein of the cytochrome bcl complex of *Caenorhabditis elegans*, a model parasitic nematode. J Biochem (Tokyo) 139: 967–979.
- Lok JB, Artis D (2008) Transgenesis and neuronal ablation in parasitic nematodes: revolutionary new tools to dissect host-parasite interactions. Parasite Immunol 30: 203–214.
- Devaney E, O'Neill K, Harnett W, Whitesell L, Kinnaird JH (2005) Hsp90 is essential in the filarial nematode *Brugia pahangi*. Int J Parasitol 35: 627–636.
- Garofalo A, Rowlinson MC, Amambua NA, Hughes JM, Kelly SM, et al. (2003) The FAR protein family of the nematode *C. elegans*. Differential lipid binding properties, structural characteristics, and developmental regulation. J Biol Chem 278: 8065–8074.
- 44. Basavaraju S, Zhan B, Kennedy MW, Liu Y, Hawdon J, et al. (2003) Ac-FAR-1, a 20 kDa fatty acid- and retinol-binding protein secreted by adult *Ancylostoma caninum* hookworms: gene transcription pattern, ligand binding properties and structural characterisation. Mol Biochem Parasitol 126: 63–71.
- Fairfax KC, Vermeire JJ, Harrison LM, Bungiro RD, Grant W, Husain, et al. (2003) Characterisation of a fatty acid and retinol binding protein orthologue from the hookworm *Ancylostoma ceylanicum*. Med Microbiol Immunol 192: 47–52.
- Kennedy MW, Garside LH, Goodrick LE, McDermott L, Brass A, et al. (1997) The Ov20 protein of the parasitic nematode *Onchoeeca volvulus*. A structurally novel class of small helix-rich retinol-binding proteins. J Biol Chem 272: 29442–29448.
- Burbelo PD, Leahy HP, Iadarola MJ, Nutman TB (2009) Four-antigen mixture for rapid assessment of Onchocerca volvulus infection. PLoS Negl Trop Dis 3: e438.

- 48. C. elegans Sequencing Consortium (1998) Genome sequence of the nematode C. elegans a platform for investigating biology. Science 282: 2012–8.
 49. Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, et al. (2001) The
- sequence of the human genome. Science 291: 1304. 50. Arca B, Lombardo F, Valenzuela JG, Francischetti IM, Marinotti O, et al.
- (2005) An updated catalogue of salivary gland transcripts in the adult female mosquito, *Anopheles gambiae*. J Exp Biol 208: 3971–3986.
- 51. Wynn TA, Thompson RW, Cheever AW, Mentink-Kane MM (2004) Immunopathogenesis of schistosomiasis. Immunol Rev 201: 156-167.
- 52. Driguez P, Doolan DL, Loukas A, Felgner PL, McManus DP (2010)
- Dirguz T, Donal DL, Dokas A, Fegnet TL, Methanis DF (2016) Schistosomiasis vaccine discovery using immunomics. Parasit Vectors 3: 4.
 Gobert GN, Jones MK (2008) Discovering new schistosome drug targets: the role of transcriptomics. Curr Drug Targets 9: 922–930.