

2-2021

Assessment of the Diagnostic Accuracy of Core Needle Biopsy in the Diagnosis of Lymphoma

Christopher A. Gardner

Xiangyun Ye

Catherine Tucker, MD

Austin Redilla

Guldeep Uppal, MD

See next page for additional authors

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2023_phase1



Part of the [Translational Medical Research Commons](#)

[Let us know how access to this document benefits you](#)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Christopher A. Gardner; Xiangyun Ye; Catherine Tucker, MD; Austin Redilla; Guldeep Uppal, MD; and Adam F. Binder



**Sidney Kimmel
Medical College™**
at Thomas Jefferson University

Assessment of the Diagnostic Accuracy of Core Needle Biopsy in the Diagnosis of Lymphoma

Christopher A. Gardner, Xiangyun Ye, Catherine Tucker, MD, Austin Redilla**,
Guldeep Uppal, MD, Adam F. Binder, MD*

(*) indicates primary project advisor

(**) indicates another student who is declaring the same project as primary for SI

Introduction & Objectives

- Excisional biopsy (EB) of an involved lymph node or tissue is the gold standard for diagnosing lymphoma.
- Recent literature suggests that novel diagnostic techniques (e.g. IHC, FC, FISH/Cytogenetics) enhance the accuracy of less invasive diagnostic procedures like Core needle biopsy (CNB)
- At TJUH, CNB has been ordered with increasing frequency
 - from 2016-2018 CNB for diagnosis of lymphoma increased from 19% to 31.6%
- CNB is less invasive, cheaper
- Objective: To determine if CNB are diagnostically adequate compared to EB

Research Question & Hypothesis

- Question: Does CNB with ancillary studies yield similar diagnostic adequacy rates compared to EB in patients with lymphoma at TJUH?
 - Variables: pathologist, class of lymphoma, site, FNA performed, number of passes, gauge size, subsequent biopsy, ancillary studies
- Hypothesis: CNB yields similar diagnostic adequacy compared to excisional biopsies
 - Adequacy determined by reviewing pathologist and whether subsequent biopsy was performed

Approach and Methods

- Retrospective cohort study
- TJUH patients from 01/01/2016 – 12/31/2019 who received a lymphoma dgx
- Intervention: CNB
- Comparison group: EB
- Outcome Measured: Was a diagnosis achieved?
- EPIC, Data collected in REDCap
- Diagnostic odds ratio and confidence interval (Baptista-Pike)



Diagnostic OR and NNH

- A total of 579 biopsies were collected for review, 122 were excluded
- Excisional biopsy adequacy: 96.8% (328)
- CNB adequacy: 56.8% (67)
- Diagnostic odds ratio of CNB was determined to be 0.03583, [95% confidence interval {CI}: 0.01695 to 0.07532] (Baptista-Pike), $p < 0.0001$ (Chi square).
- Number needed to harm: 2.5



Breakdown of Biopsies

	Excisional Biopsy N (%)	Core Needle Biopsy N (%)
Total Samples	339 (74.2%)	118 (25.8%)
Adequacy of sample	328 (96.8%)	67 (56.8%)
Diagnosis		
Benign	154 (45.4%)	50 (42.4%)
Diffuse Large B Cell Lymphoma	30 (8.5%)	12 (10.2%)
Other Large B Cell Lymphoma	13 (3.8%)	6 (5.1%)
Hodgkin Lymphoma	38 (11.2%)	7 (5.9%)
Follicular Lymphoma Grade 1/2/3A	29 (8.6%)	8 (6.8%)
Follicular Lymphoma Grade 3B	4 (1.2%)	0 (0.0%)
Mantle Cell Lymphoma	9 (2.7%)	2 (1.7%)
Marginal Zone Lymphoma	7 (2.1%)	1 (0.85%)
SLL/CLL	7 (2.1%)	3 (2.5%)
Peripheral T Cell Lymphoma, NOS	6 (1.8%)	0 (0.0%)
Angioimmunoblastic T Cell Lymphoma	4 (1.2%)	0 (0.0%)
ATLL	1 (0.3%)	0 (0.0%)
Other T cell Lymphoma	4 (1.2%)	0 (0.0%)
Solid tumor	19 (5.6%)	18 (15.3%)
Other	13 (3.8%)	10 (8.5%)
Lymphoplasmacytic Lymphoma	1 (0.3%)	0 (0.0%)
Low Grade NHL, NOS	0 (0.0%)	1 (0.85%)
Biopsy Location		
Cervical	136 (40.1%)	10 (8.5%)
Supraclavicular	21 (6.2%)	14 (11.9%)
Axillary	58 (17.1%)	41 (34.7%)
Mediastinal	6 (1.8%)	2 (1.7%)
Retroperitoneal	1 (0.3%)	12 (10.2%)
Intra-abdominal	14 (4.1%)	13 (11.0%)
Stomach	1 (0.3%)	0 (0.0%)
Other	17 (5.0%)	6 (5.1%)
Inguinal	79 (23.3%)	17 (14.4%)
Iliac	6 (1.8%)	3 (2.5%)
Reviewing Pathologist		
Hematopathologist	300 (88.5%)	84 (71.2%)
Non-Hematopathologist	39 (11.5%)	34 (28.8%)

Conclusions

- Our diagnostic OR was less than expected, suggesting CNB is not as effective as EB at diagnosing lymphoma
- These findings challenge literature suggesting CNB is non-inferior to EB (Allin D., et al 2017)
- Our findings beg the question why TJUH has been moving away from EB in recent years, as well as the decisions to include or exclude certain ancillary studies with CNB

Future Directions

- Currently analyzing 2020 lymphoma diagnoses
- Better understand the rationale for choosing CNB over EB at easily accessible EB sites
- QI project to reduce the number of CNB for suspected biopsies

Acknowledgements

- Dr. Adam Binder
- Dr. Catherine Tucker
- George Ye
- Austin Redilla