

6-20-2019

Risk Factors for the Elevation of Psychosocial Barriers in the Recovery of Lumbar Pathology

Christopher Keating

Cameron Hui

Gerard Smith

Tingting Zhan

Follow this and additional works at: <https://jdc.jefferson.edu/ptfp>

 Part of the [Physical Therapy Commons](#)

[Let us know how access to this document benefits you](#)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Physical Therapy Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.



Risk Factors for the Elevation of Psychosocial Barriers in the Recovery of Lumbar Pathology

Christopher Keating^{1*}, Cameron Hui², Gerard Smith³ and Tingting Zhan⁴

*Correspondence: christopher.keating@jefferson.edu



CrossMark

Click for updates

¹Assistant Professor, Department of Physical Therapy, Jefferson (Philadelphia University + Thomas Jefferson University), Jefferson - Center City Campus, 901 Walnut Street Room 521 Philadelphia, Pennsylvania, USA.

²Orthopedic Resident, Thomas Jefferson University Orthopedic Physical Therapy Residency Program Christiana Care Health Services, 3506 Kennett Pike Wilmington, Delaware, USA.

³PT Educator, University of Delaware Physical Therapy Department Clinic 540 South College Ave, Suite 160, Newark, Delaware, USA.

⁴Division of Biostatistics, Department of Pharmacology and Experimental Therapeutics Jefferson (Philadelphia University + Thomas Jefferson University) Jefferson - Center City Campus, 1015 Chestnut Street, Suite 520, Philadelphia, USA.

Abstract

Background/Purpose: The prevalence of low back pain (LBP) continues to rise and an appreciation for a biopsychosocial approach, which takes in multiple facets of a patient presentation, may significantly impact this increase. To the best of our knowledge, there are no studies evaluating the relationship between the STarTBack Screening Tool (SBT) and the access point the patient enters the health system. Our hypothesis is that patients that access care through the medical model will exhibit greater psychosocial risk.

Methods: A retrospective analysis of 1404 cases of LBP utilizing past medical history, demographic information, and SBT data. A linear regression model to identify potential predictors was utilized followed by analyses using cumulative logistic regression model with potential predictors.

Results: Referral source had a positive impact on the SBT score and higher probability of high risk classification. Compared to direct access, accessing care through a primary care physician had an odds ratio of 2.802 and specialist an odds ratio of 3.195.

Conclusion: This study provides evidence to the impact that access may have on the psychosocial status of patients with LBP. The approach taken by the healthcare provider may impact long term pain and functional outcomes and costs.

Keywords: Low Back Pain, Direct Access, Health Systems

Introduction

The prevalence of low back pain (LBP) continues to rise, despite the increasing technological advancement in the medical field [1-3]. Following the natural course and history, episodes of LBP increase healthcare utilization and lead more patients into chronic stages [4,5]. A clear association was demonstrated between length of disability and healthcare cost, with 86% of episodes less than one month yielding only 11% of all costs [6]. This reveals the importance of preventing and reducing the frequency of patients who transition from acute to chronic stages [5,7]. A biomedical, pathoanatomical model negatively influence the patient's overall status and those dealing with

chronic low back pain [8,9]. A recent systematic review demonstrates that a multidisciplinary biopsychosocial rehabilitation approach is an effective treatment strategy for low back pain [10]. A biopsychosocial approach takes in multiple facets of a patient presentation. A general outline involves assessing for patients pain, somatic, medical, cognitive, emotional, behavioral, social factors, and motivation for a complete clinical picture of a condition [11].

Most patients will not have a meaningful pathoanatomical diagnosis as imaging findings may be considered abnormal despite being asymptomatic, and the likelihood of abnormalities increase with age [12]. Trends demonstrate earlier medi-

cal imaging led to greater probability towards surgery and healthcare spending [1]. Structural change on imaging in an asymptomatic population continues to support the notion that other factors such as emotion, behavior, cognitive beliefs may contribute to patient's pain perception. Patients' access care differently depending on their cognitive belief system, and will enter a persistent state or recovery influenced by their beliefs [13].

The Start Back Screening Tool (SBT) stratifies patients accordingly to low, medium, high risk for biobehavioral influences on pain, demonstrating good reliability [14-21]. Those classified as high risk may benefit from increased education regarding their condition [2,22-25]. A Cochrane review lends support to the use of a biopsychosocial approach for treatment of sub-acute LBP, compared to usual care [26]. Emerging evidence continues to support use of the biopsychosocial framework, despite low quality in nature [27]. Findings also suggest that this approach may be comparable to other forms other PT interventions, for a chronic population [28,29].

To the best of our knowledge, there are no studies evaluating the relationship of the score on the SBT to the access point the patient enters the health system. Our hypothesis is that patients that access care through the traditional medical model (primary care or specialist) will exhibit greater psychosocial risk. This would be the first study to look at the factors that may elevate psychosocial risk in achieving an outcome to reduce to the likelihood of developing chronic LBP.

Methods

Retrospective analysis of 1404 cases of LBP from September 2015-December 2016 utilizing FOTO (Focus on Therapeutic Outcomes) and Clinicient (electronic medical records system) to gather data from 13 outpatient physical therapy clinics in the Philadelphia, PA area. All patients evaluated during this time, with complete intake information, were included in the study. Patient information was exported from FOTO and integrated with referral source information gathered from the EMR during the intake process at the outpatient clinical sites. Classification of the referral source into primary care, specialist, or direct access was completed by a research assistant with the help from a marketing representative as needed. Utilizing past medical history, demographic information, referral source, and SBT data we performed a linear regression model to identify potential predictors that impact the SBT score. The SBT risk level was analyzed using cumulative logistic regression model with potential predictors.

Results

In 1404 patients, 366 (26.1%) patients were classified as high risk per the SBT (Table 1). Linear regression model revealed a number of significant factors to elevate the SBT score (Table 2). A cumulative logistic regression model with these potential predictors identified the odds ratio to determine how impactful these factors were on probability of risk elevation. Our analysis

Table 1. SStarT Back Screening Tool distribution results.

Risk Level	Value	Percent	95% CI
Low	455	32.4%	(30.0%, 34.9%)
Medium	583	41.5%	(38.9%, 44.1%)
High	366	26.1%	(23.8%, 28.4%)

Table 2. Linear regression model results.

Factor	Mean Difference (SBT raw score)	95% CI	P-Value
Previous Surgery	1.442	(0.464, 2.42)	.004*
BMI	0.013	(.003, .023)	.009*
Osteoarthritis	0.286	(.013, .560)	.040*
COPD	0.994	(.150, 1.837)	.021*
HTN	0.307	(0.030, 0.583)	.030*
Headaches	0.676	(0.350, 1.003)	<.001*
Major Vision	0.581	(0.083, 1.079)	.022*
Previous MVA	0.676	(0.289, 1.062)	<.001*
Anxiety	0.554	(0.171, 0.937)	.005*
Depression	0.545	(0.143, 0.947)	.008*
Sleep Dysfunction	0.547	(0.119, 0.895)	.002*
Referral (PCP vs DA)	1.122	(0.435, 1.808)	.001*
Referral (Specialist vs DA)	1.265	(0.600, 1.930)	<.001*
Osteoporosis	-0.492	(-0.990, 0.006)	.053
Myocardial Infarction	-0.690	(-1.544, 0.164)	.113
Cancer	-0.500	(-1.007, 0.007)	.053

Table 3. Cumulative logistic regression model results.

Factor	Odds Ratio	95% CI
Age	0.984	(0.977, 0.990)
Surgery	4.073	(1.706, 10.504)
BMI	1.017	(1.003, 1.032)
Referral (PCP vs DA)	2.802	(1.521, 5.326)
Referral (Specialist vs DA)	3.195	(1.763, 5.982)
Positive PMHx	1.437	(1.336, 1.547)

found previous surgery (OR 4.073), higher BMI (OR 1.017), OA (mean 0.286), COPD (mean 0.994), HTN (mean 0.307), HA (mean 0.676), major vision disease (mean 0.581), previous accident (mean 0.676), anxiety (mean 0.554), depression (mean 0.545), sleep dysfunction (mean 0.547), and referral source had a positive impact on the SBT score and probability to have high-risk classification. Osteoporosis (mean -0.492), myocardial infarction (mean -0.690), and previous episode of cancer (mean -0.500) had a negative impact on SBT score and lower probability of high-risk classification. Referral source was positive as predicted to increase the SBT risk level. Compared to direct access, accessing care through a primary care physician had

an odds ratio of 2.802 and specialist an odds ratio of 3.195 to present with high risk as measured by the SBT.

Discussion

Positive Predictors

Previous Surgery

Multiple explanations can be used to rationalize prior surgery as a potential predictor of a high SBT score. When assessing the long term outcomes of lumbar surgical intervention, many procedures in current practice have limited outcomes [30,31]. This has especially been noted with surgeries such as the lumbar fusion and discectomy [32-34]. Across the spectrum of 4 year, 7 year, and 20 year follow up, outcomes are similar and not superior to physical therapy and cognitive behavioral therapy. It may be implied from the previous studies that continued disability, along with use of unsuccessful surgical intervention may continue to lead a patient to a lack of understanding regarding the complex nature of lumbar pathology.

Higher BMI

The association between LBP and BMI is conflicting [35]. The heterogeneity of the patient population in this study may result in its modest odds ratio. While chronic low back pain has support to have no relationship, radicular symptoms have been found to be related to higher BMI [36,37]. Given these populations were not separated; it is possible to have a subpopulation that BMI is relevant in the development of increased risk. Non-specific LBP and radicular LBP may be managed differently from healthcare providers leading to this sub classification consideration in future studies.

Osteoarthritis

Level of pain and pain related fear have been shown to be significantly associated with functional limitations in patients with osteoarthritis (OA) [38,39]. Anxiety has also been shown to predict poorer function in people with OA [39,40]. Other studies in the OA population have shown that pain catastrophizing and pain related fear are significantly related to psychological disability and have been shown to explain pain and disability in the OA population [41]. Low levels in catastrophizing and pain related fear have been shown to be related to milder symptoms of knee osteoarthritis [39].

COPD

Chronic obstructive pulmonary disease (COPD) is a condition in which many emotional needs are seen in this population [42,43]. These needs are similar to the domains screened in the SBT (fear and anxiety) [42,43]. Patient understanding the effects of exercise, motivation, and ability to control the condition are all barriers as well contributing to the thoughts and beliefs that lead to avoidance behaviour [42-44]. One barrier to this is the ability of healthcare providers to make referral to pulmonary rehabilitation, know the influence of exercise on the condition, or overcome other barriers to attendance

and completion to program [44]. These barriers correlate to the SBT assessment and suggest that these patients are at risk of a higher psychosocial stratification.

Hypertension (HTN)

Three main barriers exists to the self-management of HTN; medication use, diet, and exercise [45-47]. There are several barriers to this with some not being captured through our data collection (culture, education level, time constraint, financial status) [45,47]. Exercise limitations have been found to be due to BMI >30, female gender, low education level, being employed, psychosocial issues, and belief that MI is likely with current lifestyle [46]. The psychosocial variables found to be predictive of poor exercise adherence were stress, depression, self-worth, and lack of carefulness. These share a relationship with the SBT and therefore, would increase the likelihood of developing a high-risk profile. Identifying patients with poor adherence may subgroup patients to better identify people in need of psychologically informed management of LBP.

Headache

As in chronic pain conditions and low back pain, people who are suffering from recurrent headaches have been shown to display elevated pain related fear and avoidance of the behavior [48,49]. Anxiety sensitivity (AS) has also been shown to have a direct relationship to fear of pain in patients with recurrent headache [49]. As has been shown to be related to higher cognitive anxiety, fear of pain, and avoidance in chronic pain patients [50]. Other research has shown the in people dealing with headaches, higher scores on in fear/avoidance were related to lower pain pressure thresholds on the head [51]. Anxiety and depression have also been shown to be associated with both migraine and non-migrainous headaches [52].

Major vision disease

Major vision disease in a general term focused at capturing all patients with a variety of visual disease that affects their daily life. Little research has been done on the impact vision disease has on exercise and the barriers these individuals have regarding exercise. Haegele et al, have found that exercise does have a positive effect on self-efficacy, but this population is similar to other populations with a disability and are inactive individuals [53,54]. The barriers within this population is significant, but working with healthcare providers to overcome these barriers is important to reduce the likelihood of developing a high-risk profile with lumbar pathology.

Previous accident (MVA)

% of individuals who experience WAD will have persistent pain and disability and a variety of other symptoms including headaches and sleep disturbances among others [55]. The transition from acute to chronic issues in the WAD population has been shown to at least in part be due to cognitive and psychological factors including fear and catastrophizing as

well as hyperalgesia and central sensitization [56-58]. People suffering from chronic WAD can also have symptoms of post-traumatic stress and this can be predictive of development of chronic issues when identified early [57,59].

Anxiety and Depression

The link between anxiety and depression in people suffering with pain is vast in the literature. Depression rates have been shown to be more prevalent in those with pain and vice versa [60]. As pain severity worsens, so does depression [60]. Catastrophizing has also been linked to reports of depression [61]. Anxiety has been previously shown to be related to pain severity and behavior [62,63]. Anxiety can also produce hyperalgesia related to brain function in the hippocampal region [63]. A combination of anxiety and depression with chronic musculoskeletal pain was strongly related to increased pain intensity, an increased pain interference and worse health related quality of life [64].

Sleep Dysfunction

Sleep dysfunction has been identified and is an area to be addressed in people with central sensitization and those suffering from chronic pain conditions and can lead to exacerbation of symptoms [1,65-67]. Individuals with both acute and persistent low back pain have already been shown to have decreased quality of sleep [68-70]. In addition, sleep disturbances are labeled as one of many pain sensitizing factors in individuals with low back pain [71].

Referral Source

The results from this study support the notion that referral sources may have impact on the SBT. When comparing direct access physical therapy versus primary care physicians and specialists, there is an increase risk in SBT risk stratification implicating a change in psychosocial status. This may be due to frequent studies being performed, despite continued recommendation to avoid early imaging of the lumbar spine [72-74]. In addition, physician patient education may lead to an increase risk for patient's psychosocial status. Medical providers may rely heavily on a biomechanical framework for musculoskeletal disorders [34]. The biomechanical framework minimizes the patient's overall experience with the focus on body abnormalities [27,75,76]. Similar to diagnostic imaging, both rely on the belief that abnormalities correlate/associate to patient symptoms which in actuality is false [74,76-78].

Negative Predictors

Osteoporosis

Patients presenting with osteoporosis had an average difference of -0.492 (CI -0.990, 0.006). Several clinical practice guidelines exists to help guide healthcare providers in the management in osteoporosis [79]. While there is a variability in the description of exercise parameters, there is a support for exercise for those with this condition. Patients with osteo-

porosis education have stronger beliefs in the effectiveness and safety of exercise [80]. The reinforcement of the safe effectiveness of exercise may provide the patient with a stronger desire to move and exercise compared to those without this condition.

Myocardial Infarction (MI)

In contrast to the predictors of cardiac events (Hypertension and BMI), patients who have experienced a myocardial infarction had a neutral or negative effect on risk stratification. The mean difference was -0.690 (CI -1.544, 0.0164) providing support for the effect physician encouragement of exercise has on patient belief and expectations regarding exercise. In both the European and American guidelines for coronary cardiovascular health, exercise and its effects on that condition are encouraged [81,82]. This education accompanied with a life-changing event may ensure a more adherent patient and retention of education on exercise.

Previous Cancer (CA)

Cancer survivors had a mean difference of -0.500 (CI -1.007, 0.007). This demonstrates that having a past medical history of CA had either no effect or a negative effect on probability of elevated risk with an episode of lumbar pathology. This can be explained when assessing survivors' belief about expectations of exercise. CA survivors have favorable beliefs about exercise whether or not they believed in the long-term cancer effects [83]. Exercise is often prescribed for a number of comorbidities during oncological treatment [84,85]. This encouraging support for exercise and its effects may carry over into a patient's thoughts and beliefs in regards to lumbar pathology. These factors, as well as, the perspective that CA gives someone when compared to other injuries or conditions.

Clinical Message

- Physician education and seeking behaviour may negatively impact patient psychosocial risk.
- Conditions that support exercising reduce high psychosocial risk.
- Access may have an impact beyond cost of current episode. The approach taken by healthcare providers may impact long term pain and functional perceptions.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Authors' contributions	CK	CH	JS	TZ
Research concept and design	✓	--	✓	--
Collection and/or assembly of data	✓	--	✓	✓
Data analysis and interpretation	✓	✓	✓	✓
Writing the article	✓	✓	✓	--
Critical revision of the article	✓	--	--	--
Final approval of article	✓	✓	✓	✓
Statistical analysis	✓	--	--	✓

Acknowledgments and Funding

Thank you Megan Storey for your support in this project. The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the Department of Physical Therapy at Thomas Jefferson University.

Publication history

Editor: Mohammad H. Hadadzadeh, Wheeling Jesuit University, USA.

Received: 08-May-2019 Final Revised: 07-Jun-2019

Accepted: 11-Jun-2019 Published: 20-Jun-2019

References

1. Fritz JM, Childs JD, Wainner RS and Flynn TW. **Primary care referral of patients with low back pain to physical therapy: impact on future health care utilization and costs.** *Spine (Phila Pa 1976)*. 2012; **37**:2114-21. | [Article](#) | [PubMed](#)
2. Beneciuk JM, Bishop MD, Fritz JM, Robinson ME, Asal NR, Nisenzon AN and George SZ. **The StarT back screening tool and individual psychological measures: evaluation of prognostic capabilities for low back pain clinical outcomes in outpatient physical therapy settings.** *Phys Ther*. 2013; **93**:321-33. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
3. Childs JD, Fritz JM, Wu SS, Flynn TW, Wainner RS, Robertson EK, Kim FS and George SZ. **Implications of early and guideline adherent physical therapy for low back pain on utilization and costs.** *BMC Health Serv Res*. 2015; **15**:150. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
4. Nijs J, Polli A, Willaert W, Malfliet A, Huysmans E and Coppieters I. **Central sensitisation: another label or useful diagnosis?** *Drug Ther Bull*. 2019; **57**:60-63. | [Article](#) | [PubMed](#)
5. Alhowimel A, Alotaibi M, Radford K and Coulson N. **Psychosocial factors associated with change in pain and disability outcomes in chronic low back pain patients treated by physiotherapist: A systematic review.** *SAGE Open Med*. 2018; **6**:2050312118757387. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
6. Dagenais S, Caro J and Haldeman S. **A systematic review of low back pain cost of illness studies in the United States and internationally.** *Spine J*. 2008; **8**:8-20. | [Article](#) | [PubMed](#)
7. Pincus T, Burton AK, Vogel S and Field AP. **A systematic review of psychological factors as predictors of chronicity/disability in prospective cohorts of low back pain.** *Spine (Phila Pa 1976)*. 2002; **27**:E109-20. | [Article](#) | [PubMed](#)
8. Diener I, Kargela M and Louw A. **Listening is therapy: Patient interviewing from a pain science perspective.** *Physiother Theory Pract*. 2016; **32**:356-67. | [Article](#) | [PubMed](#)
9. Nijs J, Clark J, Malfliet A, Ickmans K, Voogt L, Don S, den Bandt H, Goubert D, Kregel J, Coppieters I and Dankaerts W. **In the spine or in the brain? Recent advances in pain neuroscience applied in the intervention for low back pain.** *Clin Exp Rheumatol*. 2017; **35 Suppl 107**:108-115. | [PubMed](#)
10. Salathe CR, Melloh M, Crawford R, Scherrer S, Boos N and Elfering A. **Treatment Efficacy, Clinical Utility, and Cost-Effectiveness of Multidisciplinary Biopsychosocial Rehabilitation Treatments for Persistent Low Back Pain: A Systematic Review.** *Global Spine J*. 2018; **8**:872-886. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
11. Wijma AJ, van Wilgen CP, Meeus M and Nijs J. **Clinical biopsychosocial physiotherapy assessment of patients with chronic pain: The first step in pain neuroscience education.** *Physiother Theory Pract*. 2016; **32**:368-84. | [Article](#) | [PubMed](#)
12. Shreibati JB and Baker LC. **The relationship between low back magnetic resonance imaging, surgery, and spending: impact of physician self-referral status.** *Health Serv Res*. 2011; **46**:1362-81. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
13. Linton SJ, Buer N, Vlaeyen J and Hellsing AL. **Are fear-avoidance beliefs related to the inception of an episode of back pain? A prospective study.** *Psychol Health*. 2000; **14**:1051-9. | [Article](#) | [PubMed](#)
14. Matsudaira K, Oka H, Kikuchi N, Haga Y, Sawada T and Tanaka S. **Psychometric Properties of the Japanese Version of the StarT Back Tool in Patients with Low Back Pain.** *PLoS One*. 2016; **11**:e0152019. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
15. Robinson HS and Dagfinrud H. **Reliability and screening ability of the StarT Back screening tool in patients with low back pain in physiotherapy practice, a cohort study.** *BMC Musculoskelet Disord*. 2017; **18**:232. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
16. Raimundo A, Parraca J, Batalha N, Tomas-Carus P, Branco J, Hill J and Gusi N. **Portuguese translation, cross-cultural adaptation and reliability of the questionnaire <<Start Back Screening Tool>> (SBST).** *Acta Reumatol Port*. 2017; **42**:38-46. | [PubMed](#)
17. Bier JD, Ostelo R, Koes BW and Verhagen AP. **Validity and reproducibility of the modified StarT Back Tool (Dutch version) for patients with neck pain in primary care.** *Musculoskelet Sci Pract*. 2017; **31**:22-29. | [Article](#) | [PubMed](#)
18. Hill JC, Dunn KM, Lewis M, Mullis R, Main CJ, Foster NE and Hay EM. **A primary care back pain screening tool: identifying patient subgroups for initial treatment.** *Arthritis Rheum*. 2008; **59**:632-41. | [Article](#) | [PubMed](#)
19. Hill JC, Whitehurst DG, Lewis M, Bryan S, Dunn KM, Foster NE, Konstantinou K, Main CJ, Mason E, Somerville S, Sowden G, Vohora K and Hay EM. **Comparison of stratified primary care management for low back pain with current best practice (StarT Back): a randomised controlled trial.** *Lancet*. 2011; **378**:1560-71. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
20. Beneciuk JM and George SZ. **Pragmatic Implementation of a Stratified Primary Care Model for Low Back Pain Management in Outpatient Physical Therapy Settings: Two-Phase, Sequential Preliminary Study.** *Phys Ther*. 2015; **95**:1120-34. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
21. Beneciuk JM, Robinson ME and George SZ. **Subgrouping for patients with low back pain: a multidimensional approach incorporating cluster analysis and the StarT Back Screening Tool.** *J Pain*. 2015; **16**:19-30. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
22. Toh I, Chong HC, Suet-Ching Liaw J and Pua YH. **Evaluation of the StarT Back Screening Tool for Prediction of Low Back Pain Intensity in an Outpatient Physical Therapy Setting.** *J Orthop Sports Phys Ther*. 2017; **47**:261-267. | [Article](#) | [PubMed](#)
23. Fritz JM, Beneciuk JM and George SZ. **Relationship between categorization with the StarT Back Screening Tool and prognosis for people receiving physical therapy for low back pain.** *Phys Ther*. 2011; **91**:722-32. | [Article](#) | [PubMed](#)
24. Beneciuk JM, Fritz JM and George SZ. **The StarT Back Screening Tool for prediction of 6-month clinical outcomes: relevance of change patterns in outpatient physical therapy settings.** *J Orthop Sports Phys Ther*. 2014; **44**:656-64. | [Article](#) | [PubMed](#)
25. Magel J, Fritz JM, Greene T, Kjaer P, Marcus RL and Brennan GP. **Outcomes of Patients With Acute Low Back Pain Stratified by the StarT Back Screening Tool: Secondary Analysis of a Randomized Trial.** *Phys Ther*. 2017; **97**:330-337. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
26. Marin TJ, Van Eerd D, Irvin E, Couban R, Koes BW, Malmivaara A, van Tulder MW and Kamper SJ. **Multidisciplinary biopsychosocial rehabilitation for subacute low back pain.** *Cochrane Database Syst Rev*. 2017; **6**:CD002193. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
27. Louw A, Puentedura EJ, Diener I and Peoples RR. **Preoperative therapeutic neuroscience education for lumbar radiculopathy: a single-case fMRI report.** *Physiother Theory Pract*. 2015; **31**:496-508. | [Article](#) | [PubMed](#)
28. Critchley DJ, Ratcliffe J, Noonan S, Jones RH and Hurley MV. **Effectiveness and cost-effectiveness of three types of physiotherapy used to reduce chronic low back pain disability: a pragmatic randomized trial with economic evaluation.** *Spine (Phila Pa 1976)*. 2007; **32**:1474-81. | [Article](#) | [PubMed](#)
29. Moseley GL. **Evidence for a direct relationship between cognitive and physical change during an education intervention in people with chronic low back pain.** *Eur J Pain*. 2004; **8**:39-45. | [Article](#) | [PubMed](#)
30. Atlas SJ, Keller RB, Wu YA, Deyo RA and Singer DE. **Long-term outcomes**

- of surgical and nonsurgical management of sciatica secondary to a lumbar disc herniation: 10 year results from the maine lumbar spine study. *Spine (Phila Pa 1976)*. 2005; **30**:927-35. | [Article](#) | [PubMed](#)
31. Deyo RA and Mirza SK. The case for restraint in spinal surgery: does quality management have a role to play? *Eur Spine J*. 2009; **18** Suppl 3:331-7. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
 32. Brox JI, Nygaard OP, Holm I, Keller A, Ingebrigtsen T and Reikeras O. Four-year follow-up of surgical versus non-surgical therapy for chronic low back pain. *Ann Rheum Dis*. 2010; **69**:1643-8. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
 33. Loupasis GA, Stamos K, Katonis PG, Sapkas G, Korres DS and Hartofilakidis G. Seven- to 20-year outcome of lumbar discectomy. *Spine (Phila Pa 1976)*. 1999; **24**:2313-7. | [Article](#) | [PubMed](#)
 34. Mannion AF, Brox JI and Fairbank JC. Comparison of spinal fusion and nonoperative treatment in patients with chronic low back pain: long-term follow-up of three randomized controlled trials. *Spine J*. 2013; **13**:1438-48. | [Article](#) | [PubMed](#)
 35. Anandacoomarasamy A, Caterson I, Sambrook P, Fransen M and March L. The impact of obesity on the musculoskeletal system. *Int J Obes (Lond)*. 2008; **32**:211-22. | [Article](#) | [PubMed](#)
 36. Brooks C, Siegler JC, Cheema BS and Marshall PW. No relationship between body mass index and changes in pain and disability after exercise rehabilitation for patients with mild to moderate chronic low back pain. *Spine (Phila Pa 1976)*. 2013; **38**:2190-5. | [Article](#) | [PubMed](#)
 37. Fanuele JC, Abdu WA, Hanscom B and Weinstein JN. Association between obesity and functional status in patients with spine disease. *Spine (Phila Pa 1976)*. 2002; **27**:306-12. | [PubMed](#)
 38. Heuts PH, Vlaeyen JW, Roelofs J, de Bie RA, Aretz K, van Weel C and van Schayck OC. Pain-related fear and daily functioning in patients with osteoarthritis. *Pain*. 2004; **110**:228-35. | [Article](#) | [PubMed](#)
 39. Scopaz KA, Piva SR, Wisniewski S and Fitzgerald GK. Relationships of fear, anxiety, and depression with physical function in patients with knee osteoarthritis. *Arch Phys Med Rehabil*. 2009; **90**:1866-73. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
 40. Helminen EE, Sinikallio SH, Valjakka AL, Vaisanen-Rouvali RH and Arokoski JP. Determinants of pain and functioning in knee osteoarthritis: a one-year prospective study. *Clin Rehabil*. 2016; **30**:890-900. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
 41. Somers TJ, Keefe FJ, Pells JJ, Dixon KE, Waters SJ, Riordan PA, Blumenthal JA, McKee DC, LaCaille L, Tucker JM, Schmitt D, Caldwell DS, Kraus VB, Sims EL, Shelby RA and Rice JR. Pain catastrophizing and pain-related fear in osteoarthritis patients: relationships to pain and disability. *J Pain Symptom Manage*. 2009; **37**:863-72. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
 42. Russell S, Ogunbayo OJ, Newham JJ, Heslop-Marshall K, Netts P, Hanratty B, Beyer F and Kaner E. Qualitative systematic review of barriers and facilitators to self-management of chronic obstructive pulmonary disease: views of patients and healthcare professionals. *NPI Prim Care Respir Med*. 2018; **28**:2. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
 43. Kosteli MC, Heneghan NR, Roskell C, Williams SE, Adab P, Dickens AP, Enocson A, Fitzmaurice DA, Jolly K, Jordan R, Greenfield S and Cumming J. Barriers and enablers of physical activity engagement for patients with COPD in primary care. *Int J Chron Obstruct Pulmon Dis*. 2017; **12**:1019-1031. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
 44. Cox NS, Oliveira CC, Lahham A and Holland AE. Pulmonary rehabilitation referral and participation are commonly influenced by environment, knowledge, and beliefs about consequences: a systematic review using the Theoretical Domains Framework. *J Physiother*. 2017; **63**:84-93. | [Article](#) | [PubMed](#)
 45. Adams OP and Carter AO. Diabetes and hypertension guidelines and the primary health care practitioner in Barbados: knowledge, attitudes, practices and barriers--a focus group study. *BMC Fam Pract*. 2010; **11**:96. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
 46. Crowley MJ, Grubber JM, Olsen MK and Bosworth HB. Factors associated with non-adherence to three hypertension self-management behaviors: preliminary data for a new instrument. *J Gen Intern Med*. 2013; **28**:99-106. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
 47. Hekler EB, Lambert J, Leventhal E, Leventhal H, Jahn E and Contrada RJ. Commonsense illness beliefs, adherence behaviors, and hypertension control among African Americans. *J Behav Med*. 2008; **31**:391-400. | [Article](#) | [PubMed](#)
 48. Black AK, Fulwiler JC and Smitherman TA. The role of fear of pain in headache. *Headache*. 2015; **55**:669-79. | [Article](#) | [PubMed](#)
 49. Norton PJ and Asmundson GJ. Anxiety sensitivity, fear, and avoidance behavior in headache pain. *Pain*. 2004; **111**:218-23. | [Article](#) | [PubMed](#)
 50. Asmundson GJ and Norton GR. Anxiety sensitivity in patients with physically unexplained chronic back pain: a preliminary report. *Behav Res Ther*. 1995; **33**:771-7. | [Article](#) | [PubMed](#)
 51. Suhr J and Spickard B. Pain-related fear is associated with cognitive task avoidance: exploration of the cogniphobia construct in a recurrent headache sample. *Clin Neuropsychol*. 2012; **26**:1128-41. | [Article](#) | [PubMed](#)
 52. Zwart JA, Dyb G, Hagen K, Odegard KJ, Dahl AA, Bovim G and Stovner LJ. Depression and anxiety disorders associated with headache frequency. The Nord-Trøndelag Health Study. *Eur J Neurol*. 2003; **10**:147-52. | [PubMed](#)
 53. Haeghele JA, Hodge SR and Kozub FM. Beliefs about physical activity and sedentary behaviors of adults with visual impairments. *Disabil Health J*. 2017; **10**:571-579. | [Article](#) | [PubMed](#)
 54. Haeghele JA, Kirk TN and Zhu X. Self-efficacy and physical activity among adults with visual impairments. *Disabil Health J*. 2018; **11**:324-329. | [Article](#) | [PubMed](#)
 55. Fritz JM, Kim M, Magel JS and Asche CV. Cost-Effectiveness of Primary Care Management With or Without Early Physical Therapy for Acute Low Back Pain: Economic Evaluation of a Randomized Clinical Trial. *Spine (Phila Pa 1976)*. 2017; **42**:285-290. | [Article](#) | [PubMed](#)
 56. Walton DM and Elliott JM. An Integrated Model of Chronic Whiplash-Associated Disorder. *J Orthop Sports Phys Ther*. 2017; **47**:462-471. | [Article](#) | [PubMed](#)
 57. Pedler A and Sterling M. Patients with chronic whiplash can be subgrouped on the basis of symptoms of sensory hypersensitivity and posttraumatic stress. *Pain*. 2013; **154**:1640-8. | [Article](#) | [PubMed](#)
 58. Goldsmith R, Wright C, Bell SF and Rushton A. Cold hyperalgesia as a prognostic factor in whiplash associated disorders: a systematic review. *Man Ther*. 2012; **17**:402-10. | [Article](#) | [PubMed](#)
 59. Sterling M, Jull G, Vicenzino B, Kenardy J and Darnell R. Physical and psychological factors predict outcome following whiplash injury. *Pain*. 2005; **114**:141-8. | [Article](#) | [PubMed](#)
 60. Bair MJ, Robinson RL, Katon W and Kroenke K. Depression and pain comorbidity: a literature review. *Arch Intern Med*. 2003; **163**:2433-45. | [Article](#) | [PubMed](#)
 61. Sullivan MJ and D'Eon JL. Relation between catastrophizing and depression in chronic pain patients. *J Abnorm Psychol*. 1990; **99**:260-3. | [Article](#) | [PubMed](#)
 62. Kain ZN, Sevarino F, Alexander GM, Pincus S and Mayes LC. Preoperative anxiety and postoperative pain in women undergoing hysterectomy. A repeated-measures design. *J Psychosom Res*. 2000; **49**:417-22. | [Article](#) | [PubMed](#)
 63. Ploghaus A, Narain C, Beckmann CF, Clare S, Bantick S, Wise R, Matthews PM, Rawlins JN and Tracey I. Exacerbation of pain by anxiety is associated with activity in a hippocampal network. *J Neurosci*. 2001; **21**:9896-903. | [Article](#) | [PubMed](#)
 64. Bair MJ, Wu J, Damush TM, Sutherland JM and Kroenke K. Association of depression and anxiety alone and in combination with chronic musculoskeletal pain in primary care patients. *Psychosom Med*. 2008; **70**:890-7. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
 65. Nijs J, Van Houdenhove B and Oostendorp RA. Recognition of central sensitization in patients with musculoskeletal pain: Application of pain neurophysiology in manual therapy practice. *Man Ther*. 2010; **15**:135-41. | [Article](#) | [PubMed](#)
 66. Bigatti SM, Hernandez AM, Cronan TA and Rand KL. Sleep disturbances in fibromyalgia syndrome: relationship to pain and depression. *Arthritis Rheum*. 2008; **59**:961-7. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)

67. Choy E. **I137 The Role of Sleep in the Management of Fibromyalgia.** *Rheumatology*. 2016; **55**:i27-i27.
68. Marin R, Cyhan T and Miklos W. **Sleep disturbance in patients with chronic low back pain.** *Am J Phys Med Rehabil*. 2006; **85**:430-5. | [Article](#) | [PubMed](#)
69. Kelly GA, Blake C, Power CK, O'Keeffe D and Fullen BM. **The association between chronic low back pain and sleep: a systematic review.** *Clin J Pain*. 2011; **27**:169-81. | [Article](#) | [PubMed](#)
70. Alsaadi SM, McAuley JH, Hush JM and Maher CG. **Prevalence of sleep disturbance in patients with low back pain.** *Eur Spine J*. 2011; **20**:737-43. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
71. Bunzli S, Smith A, Schutze R, Lin I and O'Sullivan P. **Making Sense of Low Back Pain and Pain-Related Fear.** *J Orthop Sports Phys Ther*. 2017; **47**:628-636. | [Article](#) | [PubMed](#)
72. McCullough BJ, Johnson GR, Martin BI and Jarvik JG. **Lumbar MR imaging and reporting epidemiology: do epidemiologic data in reports affect clinical management?** *Radiology*. 2012; **262**:941-6. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
73. Ojha HA, Snyder RS and Davenport TE. **Direct access compared with referred physical therapy episodes of care: a systematic review.** *Phys Ther*. 2014; **94**:14-30. | [Article](#) | [PubMed](#)
74. Fritz JM, Brennan GP and Hunter SJ. **Physical Therapy or Advanced Imaging as First Management Strategy Following a New Consultation for Low Back Pain in Primary Care: Associations with Future Health Care Utilization and Charges.** *Health Serv Res*. 2015; **50**:1927-40. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
75. Marcum JA. **Biomechanical and phenomenological models of the body, the meaning of illness and quality of care.** *Med Health Care Philos*. 2004; **7**:311-20. | [PubMed](#)
76. Tousignant-Laflamme Y, Martel MO, Joshi AB and Cook CE. **Rehabilitation management of low back pain - it's time to pull it all together!** *J Pain Res*. 2017; **10**:2373-2385. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
77. Karel YH, Verkerk K, Endenburg S, Metselaar S and Verhagen AP. **Effect of routine diagnostic imaging for patients with musculoskeletal disorders: A meta-analysis.** *Eur J Intern Med*. 2015; **26**:585-95. | [Article](#) | [PubMed](#)
78. Schmidt SG. **Recognizing potential barriers to setting and achieving effective rehabilitation goals for patients with persistent pain.** *Physiother Theory Pract*. 2016; **32**:415-26. | [Article](#) | [PubMed](#)
79. Armstrong JJ, Rodrigues IB, Wasiuta T and MacDermid JC. **Quality assessment of osteoporosis clinical practice guidelines for physical activity and safe movement: an AGREE II appraisal.** *Arch Osteoporos*. 2016; **11**:6. | [Article](#) | [PubMed](#)
80. Resnick B, Nahm ES, Zhu S, Brown C, An M, Park B and Brown J. **The impact of osteoporosis, falls, fear of falling, and efficacy expectations on exercise among community-dwelling older adults.** *Orthop Nurs*. 2014; **33**:277-86. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
81. Smith SC, Jr., Allen J, Blair SN, Bonow RO, Brass LM, Fonarow GC, Grundy SM, Hiratzka L, Jones D, Krumholz HM, Mosca L, Pasternak RC, Pearson T, Pfeffer MA and Taubert KA. **AHA/ACC guidelines for secondary prevention for patients with coronary and other atherosclerotic vascular disease: 2006 update: endorsed by the National Heart, Lung, and Blood Institute.** *Circulation*. 2006; **113**:2363-72. | [Article](#) | [PubMed](#)
82. Perk J, De Backer G, Gohlke H, Graham I, Reiner Z, Verschuren M, Albus C, Benlian P, Boysen G, Cifkova R, Deaton C, Ebrahim S, Fisher M, Germano G, Hobbs R, Hoes A, Karadeniz S, Mezzani A, Prescott E, Ryden L, Scherer M, Syvanne M, Scholte op Reimer WJ, Vrints C, Wood D, Zamorano JL and Zannad F. **European Guidelines on cardiovascular disease prevention in clinical practice (version 2012). The Fifth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of nine societies and by invited experts).** *Eur Heart J*. 2012; **33**:1635-701. | [Article](#) | [PubMed](#)
83. Hirschey R, Docherty SL, Pan W and Lipkus I. **Exploration of Exercise Outcome Expectations Among Breast Cancer Survivors.** *Cancer Nurs*. 2017; **40**:E39-E46. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)
84. Bergenthal N, Will A, Streckmann F, Wolkewitz KD, Monsef I, Engert A, Elter T and Skoetz N. **Aerobic physical exercise for adult patients with haematological malignancies.** *Cochrane Database Syst Rev*. 2014; CD009075. | [Article](#) | [PubMed](#)
85. Ferrer RA, Huedo-Medina TB, Johnson BT, Ryan S and Pescatello LS. **Exercise interventions for cancer survivors: a meta-analysis of quality of life outcomes.** *Ann Behav Med*. 2011; **41**:32-47. | [Article](#) | [PubMed Abstract](#) | [PubMed FullText](#)

Citation:

Keating C, Hui C, Smith G and Zhan T. **Risk Factors for the Elevation of Psychosocial Barriers in the Recovery of Lumbar Pathology.** *Phys Ther Rehabil*. 2019; **6**:9.
http://dx.doi.org/10.7243/2055-2386-6-9