

5-31-2017


Improving Resulted Hemoglobin A1c Rates: A Feasibility Study for Point-of-Care Hemoglobin A1c Testing at an Urban Family Medicine Office

Phoebe Askie, MD, MPH
Thomas Jefferson University Hospital

Daniel Chung, MD
Thomas Jefferson University

Laura Parente, MD
Thomas Jefferson University

Yury Parra, MD
Follow this and additional works at: <https://jdc.jefferson.edu/patientsafetyposters>
Thomas Jefferson University

 Part of the [Medicine and Health Sciences Commons](#)

Grace Amadi, MD
Thomas Jefferson University

[Let us know how access to this document benefits you](#)

Recommended Citation

See next page for additional authors

Askie, MD, MPH, Phoebe; Chung, MD, Daniel; Parente, MD, Laura; Parra, MD, Yury; Amadi, MD, Grace; Kuperman, MD, MPH, Mariana R.; Reaves, MD, Bruce; Aragon, MD, Miranda; Bardet, MD, Kyle; Lai, MD, Sunny; Woods, MD, Anna; and Altshuler, MD, Marc, "Improving Resulted Hemoglobin A1c Rates: A Feasibility Study for Point-of-Care Hemoglobin A1c Testing at an Urban Family Medicine Office" (2017). *House Staff Quality Improvement and Patient Safety Conference (2016-2019)*. Poster 71.
<https://jdc.jefferson.edu/patientsafetyposters/71>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in House Staff Quality Improvement and Patient Safety Conference (2016-2019) by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Phoebe Askie, MD, MPH; Daniel Chung, MD; Laura Parente, MD; Yury Parra, MD; Grace Amadi, MD; Mariana R. Kuperman, MD, MPH; Bruce Reaves, MD; Miranda Aragon, MD; Kyle Bardet, MD; Sunny Lai, MD; Anna Woods, MD; and Marc Altshuler, MD



Improving Resulted Hemoglobin A1c Rates: A Feasibility Study for Point-of-Care Hemoglobin A1c Testing at an Urban Family Medicine Office

Pheobe Askie, MD, MPH; Daniel Chung, MD; Laura Parente, MD; Yury Parra, MD; Grace Amadi, MD; Mariana Kuperman, MD, MPH; Bruce C. Reaves, MD; Miranda Aragon, MD; Kyle Bardet, MD; Sunny Lai, MD; Anna Woods, MD; Marc Altshuler MD
Thomas Jefferson University Hospital, Department of Family And Community Medicine, Philadelphia

Background and Introduction

- Hemoglobin A1c is the gold standard for evaluating glycemic control in patients with diabetes mellitus. It is an index measure of the patient's average blood glucose level over the preceding 3 months.
- In a meta-analysis of diabetes management, rapid availability of testing (point-of-care versus routine lab) resulted in more frequent intensification of therapy and lowered A1c levels.
- Over the course of 12 months (September 2015- August 2016) only 62.0% of Jefferson Family Medicine Associates (JFMA) patients had a A1c reported

Study Aims

- Our practice's goal is to increase the number of up-to-date hemoglobin A1c for diabetic patients seen at JFMA in order to help improve glycemic control.
- The aim of this study is to see if point-of-care (POC) hemoglobin A1C is a feasible way to increase the number of up-to-date hemoglobin A1C. We looked at various factors including timing, training, and flow.

Methods


Sample Selection:

Adult patients with the diagnosis of diabetes or at risk at diabetes who had not had an A1c in the last 3 months who visited Jefferson Family Medicine Associates' Team One on two selected dates.

Process:

- Identify patients with diabetes mellitus or a risk of diabetes via chart review of Team 1 provider schedules.
- Observe medical assistants (MA) from start (MA collects blood tube) to finish (MA documents data in computer).
- Document the MA workflow through mapping and obtain qualitative feedback on the process from MA.
- Record the start and finish times.
- Analysis mapping and time measurements; as well as qualitative feedback from medical assistants.

Mapping and Data



MA collects tubes needed	MA Enters room	MA Leaves Room	Enters A1c Room	A1c Resulted	A1c Entered into Computer	Total Time (minutes)
1:24	1:27	1:29	1:29	1:33	1:36 (gave up)	9
1:42	1:45	1:47	1:47	1:51		9
1:53	1:56	2:01	2:01	2:06	2:07	14
2:08	2:10	patient left				n/a
2:35	2:38	2:39	2:40	2:44		*new diagnosis* 9
2:41	2:46	2:48	2:48	(came back 28 s after result) 2:52		13
2:55	3:02	3:03	3:03	(came back @ 3:06 after flu shot) 3:07		13
3:35	3:37	3:39	3:39	3:43	3:45	10
4:26	4:30	4:31	4:32	4:36	4:38	12
1:15	1:17	1:22	1:23	1:27	gave up	n/a

Result Analysis

- An average of 10.2 additional minutes were spent by the MAs performing the test.
- MAs were able to incorporate operating the A1c machine into their workflow and could consecutively perform tasks such as urinalysis and administering vaccines while the A1c sample was being processed.
- MAs reported that additional training is needed.
- Feedback from the medical assistants:

"The instructions don't actually tell you how to do it - the details like "green stuff should be facing the wall, put in the patient identifiers first"

"I'm on a mission!" "Because everyone gets in my way!"

"My patient is waiting for this...I only have 2 rooms. I feel bad."

Conclusions

- It is feasible to have an A1c in our practice to implement POC A1C testing
- Successful implementation would require buy-in from MAs and providers as well as potential reconstructing the geography of the office/clinic

Challenges and Limitations

- Small sample size that may not be representative of JFMA patient's population
- We did not look at how collection of POC A1C affected patient outcomes or clinical management
- Lack of comparison group to determine if similar delays exist with other POC testing and interventions (ie EKGs, vaccines, INR, etc.)
- Cost of testing: \$8/A1c slide

Potential Future Directions

- Improve training for medical assistants on POC A1C testing
- Increase provider awareness of POC A1C testing
- Implement study of POC A1C testing across the practice over a longer period of time
- Compare diabetes outcomes of patients who received POC A1c testing vs traditional laboratory testing
- Look at the effect of adding more POC A1C machines on office flow and ability to obtain A1C.

References

- Al-Ansary L, Farmer A, Hirst J, et al. "Point-of-Care Testing for HbA1c in the Management of Diabetes: A Systematic Review and Metaanalysis." *Clinical Chemistry*. 57.4 (2011):568-76.
- Delamater AM. "Clinical Use of Hemoglobin A1c to Improve Diabetes Management." *Clinical Diabetes*. 24.1(2006):6-8.
- Diabetes Control and Complications Research Group. "The Effect of Intensive Treatment of Diabetes on the Development and Progression of Long-Term Complications in Insulin-Dependent Diabetes Mellitus." *New England Journal of Medicine*. 329.14(1993):977-86.
- Kennedy L, Herman WH, Strang P, et al. "Impact of Active Versus Usual Algorithmic Titration of Basal Insulin and Point-Of-Care Versus Laboratory Measurement of HbA1c on Glycemic Control in Patient With Type 2 Diabetes." *Diabetes Care*. 29.1 (2006):1-8.
- Miller CD, Barnes CS, Phillips LS, Ziemer DC, et al. "Rapid A1c Availability Improves Clinical Decision-Making in an Urban Primary Care Clinic." *Diabetes Care*. 26.4 (2003): 1158-63.
- Stratton IM, Adler AI, Neil HAW, et al. "Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes mellitus (UKPDS 35): prospective observational study." *BMJ*. 321 (2000): 405-12.

Acknowledgements

- Team One medical assistants and nurse
- Team One Providers
- Thomas Jefferson University Hospital, Department of Family and Community Medicine