

1-15-2021

## Myeloid-derived suppressor cells in gastrointestinal cancers: A systemic review.

Maham Farshidpour  
*Banner University of Medical Center*

Monjur Ahmed  
*Thomas Jefferson University*

Shilpa Junna  
*Banner University of Medical Center*

Juanita L. Merchant  
*Banner University of Medical Center*

Follow this and additional works at: <https://jdc.jefferson.edu/internalfp>

 Part of the [Internal Medicine Commons](#)

[Let us know how access to this document benefits you](#)

### Recommended Citation

Farshidpour, Maham; Ahmed, Monjur; Junna, Shilpa; and Merchant, Juanita L., "Myeloid-derived suppressor cells in gastrointestinal cancers: A systemic review." (2021). *Division of Internal Medicine Faculty Papers & Presentations*. Paper 38.

<https://jdc.jefferson.edu/internalfp/38>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Division of Internal Medicine Faculty Papers & Presentations by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

# World Journal of *Gastrointestinal Oncology*

*World J Gastrointest Oncol* 2021 January 15; 13(1): 1-91



### THERAPEUTIC AND DIAGNOSTIC GUIDELINES

- 1 Myeloid-derived suppressor cells in gastrointestinal cancers: A systemic review  
*Farshidpour M, Ahmed M, Junna S, Merchant JL*

### ORIGINAL ARTICLE

#### Clinical and Translational Research

- 12 Laparoscopy-assisted transanal total mesorectal excision for lower rectal cancer: A feasible and innovative technique  
*Li YJ, Wang L, Sun TT, Wu AW*

#### Retrospective Study

- 24 Survival outcomes and prognostic indicators for gastric cancer patients with positive peritoneal wash cytology but no peritoneal metastasis after radical gastrectomy  
*Kang WZ, Zhong YX, Ma FH, Xue LY, Xiong JP, Ma S, Li Y, Xie YB, Quan X, Tian YT*

#### Observational Study

- 37 Mining The Cancer Genome Atlas database for tumor mutation burden and its clinical implications in gastric cancer  
*Zhao DY, Sun XZ, Yao SK*
- 58 Diagnostic performance of narrow-band imaging international colorectal endoscopic and Japanese narrow-band imaging expert team classification systems for colorectal cancer and precancerous lesions  
*Wang Y, Li WK, Wang YD, Liu KL, Wu J*

### META-ANALYSIS

- 69 Efficacy and safety of intraoperative radiotherapy in rectal cancer: A systematic review and meta-analysis  
*Liu B, Ge L, Wang J, Chen YQ, Ma SX, Ma PL, Zhang YQ, Yang KH, Cai H*

### CASE REPORT

- 87 Internal hemorrhoid harboring adenocarcinoma: A case report  
*Caparelli ML, Batey JC, Tailor A, Braverman T, Barrat C*

**ABOUT COVER**

Editorial board member of *World Journal of Gastrointestinal Oncology*, Dr. Una Cidon is a medical oncologist at Dorset University Hospitals NHS Foundation Trust, United Kingdom. She received her Bachelor's degree in medicine from Salamanca's University (Spain) and undertook postgraduate training at the Asturias Central University Hospital in Oviedo, receiving title of Specialist in Medical Oncology in 2004. In 2009, she obtained her PhD from the Clinical University Hospital of Valladolid. She then became Associate Professor of Oncology at the University of Valladolid and obtained a Master's degree in Molecular Oncology from the Spanish Centre for Cancer Research. Her ongoing research interests involve the design and conduct of clinical investigations to improve quality of life of patients receiving antineoplastic treatments and disseminating educational information. (L-Editor: Filipodia)

**AIMS AND SCOPE**

The primary aim of *World Journal of Gastrointestinal Oncology* (WJGO, *World J Gastrointest Oncol*) is to provide scholars and readers from various fields of gastrointestinal oncology with a platform to publish high-quality basic and clinical research articles and communicate their research findings online.

WJGO mainly publishes articles reporting research results and findings obtained in the field of gastrointestinal oncology and covering a wide range of topics including liver cell adenoma, gastric neoplasms, appendiceal neoplasms, biliary tract neoplasms, hepatocellular carcinoma, pancreatic carcinoma, cecal neoplasms, colonic neoplasms, colorectal neoplasms, duodenal neoplasms, esophageal neoplasms, gallbladder neoplasms, etc.

**INDEXING/ABSTRACTING**

The WJGO is now indexed in Science Citation Index Expanded (also known as SciSearch®), PubMed, and PubMed Central. The 2020 edition of Journal Citation Reports® cites the 2019 impact factor (IF) for WJGO as 2.898; IF without journal self cites: 2.880; 5-year IF: 3.316; Ranking: 143 among 244 journals in oncology; Quartile category: Q3; Ranking: 55 among 88 journals in gastroenterology and hepatology; and Quartile category: Q3.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Jia-Hui Li; Production Department Director: Xiang Li; Editorial Office Director: Ya-Juan Ma.

**NAME OF JOURNAL**

*World Journal of Gastrointestinal Oncology*

**ISSN**

ISSN 1948-5204 (online)

**LAUNCH DATE**

February 15, 2009

**FREQUENCY**

Monthly

**EDITORS-IN-CHIEF**

Rosa M Jimenez Rodriguez, Pashtoon Kasi, Monjur Ahmed

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/1948-5204/editorialboard.htm>

**PUBLICATION DATE**

January 15, 2021

**COPYRIGHT**

© 2021 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>



## Myeloid-derived suppressor cells in gastrointestinal cancers: A systemic review

Maham Farshidpour, Monjur Ahmed, Shilpa Junna, Juanita L Merchant

**ORCID number:** Maham Farshidpour 0000-0001-6282-6148; Monjur Ahmed 0000-0003-0515-9224; Shilpa Junna 0000-0001-8312-986X; Juanita L Merchant 0000-0002-6559-8184.

**Author contributions:** Farshidpour M and Merchant JL conceived of and designed the study, and collected the data; Farshidpour M, Ahmed M, Junna S, and Merchant JL drafted the manuscript; all authors reviewed the results and approved the final version of the manuscript.

**Conflict-of-interest statement:** The authors declare that they have no conflicts of interest.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Invited manuscript

**Maham Farshidpour**, Inpatient Medicine, Banner University of Medical Center, Tucson, AZ 85724, United States

**Monjur Ahmed**, Division of Gastroenterology and Hepatology, Department of Internal Medicine, Thomas Jefferson University, Philadelphia, PA 19107, United States

**Shilpa Junna, Juanita L Merchant**, Division of Gastroenterology and Hepatology, Banner University of Medical Center, Tucson, AZ 85724, United States

**Corresponding author:** Maham Farshidpour, MD, Doctor, Inpatient Medicine, Banner University of Medical Center, 1501 N Campbell Ave, Tucson, AZ 85724, United States. [mfarshidpour@email.arizona.edu](mailto:mfarshidpour@email.arizona.edu)

### Abstract

Gastrointestinal (GI) cancers are one of the most common malignancies worldwide, with high rates of morbidity and mortality. Myeloid-derived suppressor cells (MDSCs) are major components of the tumor microenvironment (TME). MDSCs facilitate the transformation of premalignant cells and play roles in tumor growth and metastasis. Moreover, in patients with GI malignancies, MDSCs can lead to the suppression of T cells and natural killer cells. Accordingly, a better understanding of the role and mechanism of action of MDSCs in the TME will aid in the development of novel immune-targeted therapies.

**Key Words:** Myeloid-derived suppressor cells; Gastrointestinal cancers; Immune checkpoint inhibitors; Tumor progression

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** In patients with cancer, the levels of myeloid-derived suppressor cells (MDSCs) are presumed to be of prognostic and predictive value. Recent studies have shown that MDSCs appear to be independent prognostic factors in gastrointestinal cancer. In addition, therapeutics that target MDSCs have been shown to enhance anti-tumor immune responses in animal models. Consequently, a better understanding of the role and mechanism of action of MDSCs in the tumor microenvironment may aid in the development of novel immune-targeted therapies.



**Specialty type:** Oncology**Country/Territory of origin:** United States**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): 0

Grade C (Good): C

Grade D (Fair): D

Grade E (Poor): 0

**Received:** August 25, 2020**Peer-review started:** August 25, 2020**First decision:** November 16, 2020**Revised:** December 1, 2020**Accepted:** December 16, 2020**Article in press:** December 16, 2020**Published online:** January 15, 2021**P-Reviewer:** Gao F, Mortazavian AM**S-Editor:** Huang P**L-Editor:** A**P-Editor:** Li JH**Citation:** Farshidpour M, Ahmed M, Junna S, Merchant JL. Myeloid-derived suppressor cells in gastrointestinal cancers: A systemic review. *World J Gastrointest Oncol* 2021; 13(1): 1-11**URL:** <https://www.wjgnet.com/1948-5204/full/v13/i1/1.htm>**DOI:** <https://dx.doi.org/10.4251/wjgo.v13.i1.1>

## INTRODUCTION

In 2018, 4.8 million new diagnoses of gastrointestinal (GI) cancer and 3.4 million related deaths were reported globally. The incidence of GI cancer is 26% worldwide, accounting for 35% of all cancer-associated mortalities<sup>[1]</sup>. Numerous stromal and immune cells and soluble markers are related to the immunosuppressive network in the tumor microenvironment (TME)<sup>[2]</sup>. This network is involved in tumor cell growth and the blockade of anti-tumor immune responses, which subsequently promote the progression and invasion of tumor cells<sup>[3]</sup>. Macrophages, monocytes, and dendritic cells (DCs) represent a subgroup of leukocytes called myeloid cells, which are generated from polymorphonuclear granulocytes<sup>[4]</sup>. Myeloid-derived suppressor cells (MDSCs) are a heterogeneous population of immature myeloid cells that arise from myeloid progenitor cells<sup>[5]</sup>. They play a key role in tumor-associated immune evasion, angiogenesis, and tumor metastasis<sup>[6,7]</sup>.

In patients with cancer, the levels of MDSCs are thought to have prognostic and predictive significance<sup>[8]</sup>. Recent studies have examined the role of MDSCs in solid tumors, and discovered that they appear to be independent prognostic factors in GI cancer<sup>[8,9]</sup>. In a meta-analysis of 17 studies with 1115 patients with GI malignancies, patients with a higher number of MDSCs at tumor sites and peripheral blood had higher mortality rates (hazard ratio: 3.35, 95% confidence interval: 1.46-7.68;  $P = 0.0004$ ), risk of relapse, and tumor progression. The authors concluded that MDSC levels have prognostic and predictive value in cancer patients<sup>[10]</sup>. Additionally, higher levels of MDSCs in patients with cancer are associated with advanced tumor stage and a poor clinical prognosis<sup>[11]</sup>. Shibata *et al*<sup>[12]</sup> evaluated 123 patients with advanced GI malignancy, including 62 with colorectal cancer (CRC), 43 with gastric cancer (GC), and 18 with esophageal malignancies, and found that overall survival (OS) was significantly shorter in stage IV GI cancer patients with high MDSC levels than in those with low MDSC levels ( $P < 0.05$ ). Because MDSCs have a significant role in modulating cancer progression and metastasis by inhibiting the anti-tumor reactivity of T cells and natural killer (NK) cells, targeting MDSCs with immune checkpoint inhibitors (ICIs) can alleviate their pro-tumorigenic functions. Therefore, in this systematic review, we summarize the characteristics and proposed function of MDSCs in the TME and their relationship to prognosis in patients with GI cancers.

## MATERIALS AND METHODS

PubMed/MEDLINE databases were explored with search strategies using search keywords "MDSCs," "gastrointestinal cancers," "prognosis," "tumor progression," and "mortality rate," to categorize studies published between 2006 and 2020. A total of 128 articles were reviewed by the authors for relevance to MDSCs and GI cancers, including retrospective, cross-sectional, case reports, and cohort studies, of which 85 papers were selected that met our selection criteria.

## MDSC MECHANISM OF ACTION

MDSCs are myeloid-derived heterogeneous cells with potent immune regulatory functions. They are derived from the myeloid lineage of immune cells that give rise to macrophages, granulocytes, and immature DCs<sup>[13]</sup>. Monocytic MDSCs (M-MDSCs) and polymorphonuclear MDSCs (PMN-MDSCs) are the two major myeloid subsets of MDSCs<sup>[14]</sup>. Phenotypically and morphologically, they are equivalent to monocytes and neutrophils, respectively (Table 1)<sup>[15]</sup>.

The process of myelopoiesis is driven by granulocyte-macrophage colony-stimulating factor (GM-CSF) in normal physiological conditions. Ultimately, GM-CSF and M-CSF induce the differentiation of granulocytes and macrophages from a common myeloid precursor that transforms into a common myeloblast<sup>[16]</sup>. Myeloid

**Table 1** Two main categories of myeloid-derived suppressor cells and their immunosuppressive functions<sup>[59]</sup>

Type of MDSC	Markers in humans	Immunosuppression mediator	Mechanism of immunosuppression
PMN-MDSCs	CD11b <sup>+</sup> CD14 <sup>-</sup> CD15 <sup>+</sup> HLADR <sup>-</sup> or CD11b <sup>+</sup> CD14 <sup>-</sup> CD66b <sup>+</sup> or LOX-1 <sup>+</sup>	ARG1, ROS	Suppressing immune responses mainly in an antigen-specific manner; ROS production
M-MDSCs	CD11b <sup>+</sup> CD14 <sup>+</sup> CD15 <sup>-</sup> HLADR <sup>low/-</sup>	NO, ARG1, and cytokines such as TGF- $\beta$ and IL-10	Suppressing T cell responses, both in antigen-specific and non-specific manners; production of NO and cytokines

ARG1: Arginase 1; IL-10: Interleukin-10; M-MDSCs: Monocytic myeloid-derived suppressor cells; NO: Nitric oxide; PMN-MDSCs: Polymorphonuclear myeloid-derived suppressor cells; ROS: Reactive oxygen species; TGF- $\beta$ : Transforming growth factor-beta; MDSC: Myeloid-derived suppressor cell.


DCs eventually arise from monocytic as opposed to granulocytic lineages<sup>[17]</sup>. However, the hypersecretion of these mediator factors during chronic inflammation and cancer leads to the generation of MDSCs<sup>[18]</sup>. For example, inflammatory cytokines such as interleukin 6 (IL-6), IL-1 $\beta$ , and IL-3 and C-X-C chemokine receptor type 4 (CXCR4) and CXCL12 can lead to the induction and proliferation of MDSCs in peripheral blood and tumor sites in cancer patients<sup>[7,19]</sup>. The most important function of MDSCs is immunosuppression, mainly of target T cells (Figure 1)<sup>[14]</sup>.

Several *in vitro* and *in vivo* studies have documented the mechanisms underlying the immunosuppressive actions of MDSCs. Arginase 1 (ARG1), inducible nitric oxide (iNOS), reactive oxygen species, and reactive nitrogen species are important suppressive factors produced by MDSCs<sup>[20,21]</sup>. Cao *et al*<sup>[22]</sup> reported that PMN-MDSCs store ARG1 and secrete it to the TME. ARG1 and NOS activities lead to cellular depletion of L-arginine (referred to as L-arg), which is an essential substrate for T cell proliferation<sup>[22,23]</sup>. Similar to T cells, depletion of L-arg also impairs the function of NK cells<sup>[24]</sup>.

In addition to immunosuppressive factors, MDSCs can overpower T cell functions by directly engaging with T cell inhibitory and apoptotic receptors. Activated MDSCs express high levels of Fas ligand (referred to as Fas L), programmed death-ligand 1 (PD-L1), and galectin-9. Subsequently, the interaction between these ligands with their receptors on T cells leads to T cell exhaustion *via* PD-L1/programmed cell death protein 1 (PD-1) or T cell apoptosis through the Fas L/Fas and galectin-9/T cell immunoglobulin and mucin domain-3 pathways<sup>[25]</sup>. Generally, M-MDSCs have more suppressive effects than PMN-MDSCs<sup>[26]</sup>. Moreover, MDSCs can stimulate and recruit regulatory T cells (Tregs) to the TME<sup>[27]</sup>. Tregs suppress anti-tumor immunity, and the interaction between MDSCs and Tregs create a strong blockade preventing cytotoxic immune cells from mounting an anti-tumor attack<sup>[28]</sup>. Elevated levels of Tregs are associated with poor survival in patients with hepatocellular carcinoma (HCC) and pancreatic cancer<sup>[29,30]</sup>. Regarding metastasis, MDSCs can promote angiogenesis by secreting IL-28 [interferon lambda (IFN- $\lambda$ )] and matrix metalloproteinase (MMP)-9, promoting the invasion and migration of tumor cells<sup>[31]</sup>.

## MARKERS OF MDSCS IN PERIPHERAL BLOOD

MDSCs are categorized according to their phenotype, which includes several recognized surface markers, such as cluster of differentiation 33 (CD33), CD11b, or human leukocyte antigen-DR isotype (HLA-DR), as well as by the lack of expression of markers distinctive of mature lymphoid cells, such as CD3, CD19, and CD56<sup>[32]</sup>. Typically, flow cytometry is performed to isolate MDSCs from peripheral mononuclear blood cells (PBMCs)<sup>[33]</sup>. Fluorescent-labeled monoclonal antibodies are used to distinguish M-MDSCs and granulocytic MDSCs (G-MDSCs). M-MDSCs are identified as CD11b<sup>+</sup>CD14<sup>+</sup>CD33<sup>high</sup>HLA-DR<sup>low</sup> and CD66b, whereas G-MDSCs are recognized as CD11b<sup>+</sup>CD14<sup>-</sup>CD33<sup>low</sup>HLA-DR<sup>-</sup>CD66b<sup>+</sup><sup>[34]</sup>. MDSCs can be distinguished from other immune suppressor cells within the myeloid lineage, *e.g.*, tumor-associated macrophages and macrophage type 2, by other specific surface markers, including CD163 and F4/80<sup>[35]</sup>.



**Figure 1 Chronic inflammation activates myeloid-derived suppressor cell generation, migration, and immunosuppression in the tumor microenvironment.** Several cytokines and stimulator factors secreted by stroma and tumor cells (e.g., VEGF, granulocyte-macrophage colony-stimulating factor, IL-1, IL-6, HIF-1 $\alpha$ , TGF- $\beta$ , COX-2) trigger myeloid-derived suppressor cell (MDSC) generation and migration. Cytokines (e.g., CCL2, CCL3, CCL4, CCL5, CXCL1) stimulate the migration of MDSCs into the tumor microenvironment. At the tumor site, MDSCs undergo activation (via TNF, IL-10, IL-1, IL-6, IFN- $\gamma$ , COX-2, HIF-1 $\alpha$ , etc.) and suppress the anti-tumor reactivity of T and natural killer cells. Cross-talk between MDSCs and dendritic cells (DCs) impairs DC function and promotes tumor progression<sup>[81]</sup>. GM-CSF: Granulocyte-macrophage colony-stimulating factor; PMN-MDSC: Polymorphonuclear myeloid-derived suppressor cells; M-MDSC: Monocytic myeloid-derived suppressor cells; MDSC: Myeloid-derived suppressor cell; DC: Dendritic cell; ARG-1: Arginase 1; NOS: Nitric oxide; PD-L1: Programmed death-ligand 1; NK: Natural killer; ROS: Reactive oxygen species.

## MDSCS IN GI MALIGNANCIES

### Esophageal cancer

High levels of circulating MDSCs in esophageal cancer are associated with a poor prognosis<sup>[5]</sup>. Elevated MDSCs in the blood are correlated with elevated numbers of immunosuppressive cells, including Tregs<sup>[5]</sup>. Jiao *et al*<sup>[36]</sup> evaluated 31 esophageal cancer patients and 26 healthy controls (HCs), and found that MDSC numbers in the peripheral blood were increased 15-fold in esophageal cancer patients compared to HCs. The authors also showed that the plasma levels of ARG1 were 3-fold higher in cancer patients than in HCs. Xu *et al*<sup>[37]</sup> showed that 178 patients with esophageal cancer had a high level of G-MDSCs (> 82.5%), which were correlated with high morbidity due to the development of sepsis postoperatively after esophageal cancer surgery. The authors suggested that the level of G-MDSCs may be used to determine the incidence of sepsis in preoperative esophageal cancer patients postoperatively, and could improve the mortality of cancer-associated sepsis by targeting the level of MDSCs.

Additionally, a study by Chen *et al*<sup>[38]</sup> found that the levels of IL-6 and MDSCs predicted the prognosis and treatment response in mice with esophageal squamous cell carcinoma (SCC). The levels of MDSCs induced by IL-6 were linked to tumor growth and a poor prognosis. The authors concluded that targeted therapy against IL-6 with rapamycin or casein kinase 2 inhibitors might be a potential treatment modality for esophageal SCC<sup>[38,39]</sup>.

### GC

According to the Global Cancer Observatory (GLOBOCAN) 2018 database, GC is the fifth most common cancer and third most deadly cancer worldwide, with an estimated



783000 deaths in 2018<sup>[40]</sup>, Li *et al*<sup>[41]</sup> documented the levels of MDSCs in the peripheral blood of 21 GC patients who had not previously received treatment, and noted that the levels of MDSCs in these patients were about 4-fold higher than in the control groups. The authors concluded that cancer cell differentiation and lymph node metastasis are mostly related to the presence of M-MDSCs. They also showed that treatment with epirubicin and paclitaxel regimens can reduce the level of MDSCs in these patients, potentially leading to better outcomes for patients due to inhibition of cancer progression. Moreover, in 29 patients with GC and 18 HCs, MDSCs were increased in stage IV patients compared with HCs, and the 2-year survival rate of patients with higher levels of MDSCs was significantly poorer (median OS: 498 d *vs* 473 d;  $P = 0.048$ ), but no significant difference was observed in survival among patients with stage I, II, and III GC<sup>[42]</sup>. Previously, we reported that schlafen (SLFN) 4-expressing myeloid cells recruited to the stomach during *Helicobacter* infection undergo a phenotypic shift to G-MDSCs under the influence of damage-associated molecular pattern (DAMP) signaling and the production of IFN- $\alpha$ <sup>[43,44]</sup>. SLFN4 is a myeloid cell differentiation factor that controls myelopoiesis<sup>[45]</sup>. These SLFN-expressing MDSCs secrete factors including microRNAs, which can be detected in the peripheral blood as a biomarker and promote epithelial cell growth. This sustained immune dysregulation creates a microenvironment capable of supporting GC development<sup>[46]</sup>.

### HCC

HCC is a leading cause of death in cirrhotic patients. Per the GLOBOCAN 2018 database, 841000 new cases of primary liver cancer and 782000 deaths due to HCC occurred that year<sup>[40]</sup>. In a prospective case-control study, Elwan *et al*<sup>[47]</sup> demonstrated a higher number of MDSCs in the peripheral blood of cirrhotic groups without HCC than in patients with cirrhosis and HCC compared to patients in control groups. They showed that mean MDSC counts in the peripheral blood of cirrhotics without HCC group and cirrhotics with HCC group were about 3.5-fold and 5-fold higher compared to the control groups, respectively. Although not statistically significant, the authors reported a low number of MDSCs in the ascitic fluid of patients with both cirrhosis and HCC. Additionally, they investigated the correlation of levels of IFN- $\gamma$  and alpha-fetoprotein with MDSC level. Their data showed that alpha-fetoprotein was positively and INF- $\gamma$  was negatively correlated with MDSC count in the HCC group<sup>[47]</sup>. A high frequency of MDSCs in the PBMCs of patients with HCC has been linked to more aggressive forms of HCC and poor clinical outcomes following local ablation, hepatectomy, or hepatic arterial infusion chemotherapy<sup>[48,49]</sup>. A cohort study by Bayik *et al*<sup>[50]</sup> showed an upsurge in circulating MDSC frequency in 114 patients with a secondary liver cancer, including CRC with liver metastases and neuroendocrine tumors, compared to individuals with benign lesions.

Data from animal models have shown that myeloid cells secrete MMPs, serine proteases, and cysteine cathepsins, which facilitate tumor cell invasion and metastasis by disrupting cell adhesions<sup>[51]</sup>. Tumor angiogenesis in HCC can be promoted by MDSCs by producing high levels of MMP-9 in HCC<sup>[52]</sup>.

### Pancreatic cancer

The incidence of pancreatic ductal adenocarcinoma has significantly increased worldwide over the last 30 years, with a 5-year survival time less than 8%<sup>[53]</sup>. Khaled *et al*<sup>[54]</sup> demonstrated that G-MDSCs, but not M-MDSCs, are much higher in circulation and in the tumor tissue of patients with pancreatic cancer compared to HCs or those with chronic pancreatitis. These results suggest that the high level of G-MDSCs in pancreatic cancer plays a key factor in tumor development and progression. A cohort study reported that the percentages of all subpopulations of MDSCs were higher in patients with intraductal papillary mucinous neoplasm (IPMN) than in HCs, and were even higher in those with pancreatic adenocarcinoma. Although there was a trend towards higher MDSC levels in pancreatic cancer *vs* IPMN, it was not statistically significant ( $P = 0.33$ )<sup>[55]</sup>.

### CRC

A recent study reported that CRC cells induce an increase in the number of MDSCs by producing inflammatory factors, such as transforming growth factor-beta, IL-10, and ARG1<sup>[56,57]</sup>. Consequently, T cell proliferation can be suppressed by tumor-derived MDSCs and promote tumor cell growth *via* oxidative metabolism. Previously, it was shown that the numbers of circulating Tregs and MDSCs are significantly reduced following tumor resection in patients with CRC. These data indicate that immunosuppression can be mitigated by reducing the number of MDSCs and Tregs in

patients with CRC after reducing the tumor burden<sup>[57]</sup>. Tada *et al*<sup>[58]</sup> showed that patients with unresectable metastatic CRC with high M-MDSC, low CD4<sup>+</sup>, or low CD8<sup>+</sup> effector memory T cell levels had significantly shorter progression-free survival.

## MDSC-TARGETED THERAPY

Many studies have examined MDSCs as the core of targeted therapeutic strategies to improve tumor control in experimental animal models. These targeted therapies could be achieved by reducing MDSC numbers, hindering their trafficking and migration, or inhibiting their immunosuppressive function (Table 2)<sup>[59]</sup>.

Because of variances in immunophenotype and mechanisms of suppression in the TME and diverse nature of human MDSCs, it is challenging to target human MDSCs<sup>[60]</sup>. Wang *et al*<sup>[61]</sup> treated pancreatic cancer patients with cytokine-induced killer (CIK) cell immunotherapy, CIK plus gemcitabine, and 5-fluorouracil (5-FU), and analyzed the levels of MDSCs in the peripheral blood pre- and post-treatment. The OS of metastatic pancreatic patients was increased with the combination of CIK and chemotherapy (gemcitabine and 5-FU) compared to patients treated with only CIK. Also, Jiang *et al*<sup>[62]</sup> reported that the quality of life and 2-year survival rate improved in patients with advanced GC following combining chemotherapy (5-FU and oxaliplatin) with CIK cell treatment compared to treatment with chemotherapy alone. Tadalafil, the Federal Drug Administration-approved phosphodiesterase-5 inhibitor, can suppress MDSCs through downregulation of ARG1 and iNOS activities in several preclinical models<sup>[63-65]</sup>. Rawat *et al*<sup>[66]</sup> showed in aflatoxin-induced HCC rats, that tadalafil reduced the level of glutamic oxaloacetic transaminase, an important enzyme that facilitates carbohydrate and protein metabolism in cancer cells. A previous study showed that treatment with tyrosine kinase inhibitors, such as sunitinib, reduced the number of MDSCs and Tregs in animals with intrahepatic colorectal metastases<sup>[67]</sup>. The authors also showed that the number of MDSCs was significantly reduced from 53.9% in phosphate-buffered saline-treated mice to 39% in sunitinib-treated mice. Sunitinib has established efficacy against advanced GI stromal tumors<sup>[68]</sup>.

Treatment of esophageal SCC with 1 $\alpha$ ,25-dihydroxyvitamin D3 (calcitriol) has been shown to inhibit MDSC proliferation induced by IL-6 stimulation in C57 mice. Therefore, it has been proposed that this treatment may be a promising strategy for the prevention and treatment of esophageal SCC<sup>[69]</sup>.

ICIs such as cytotoxic T-lymphocyte-associated protein 4 (ipilimumab and tremelimumab), PD-1 (pembrolizumab and nivolumab), and PD-L1 (atezolizumab, avelumab, and durvalumab) are promising treatment strategies that can be applied across numerous solid tumors<sup>[70]</sup>. Pembrolizumab is a therapeutic antibody that blocks PD-1, and has shown promising anti-tumor activity in advanced GC<sup>[71]</sup>. Although ICIs show great therapeutic benefits, substantial GI side effects such as colitis and GI bleeding can limit their use<sup>[72]</sup>. However, due to immunosuppression, which is regulated by MDSCs, some patients with cancer may develop resistance to ICIs<sup>[73]</sup>. Therefore, it is important to inhibit MDSC proliferation and migration to the TME by different strategies, such as anti-CXCR2 monoclonal antibody, to enhance PD-1 efficacy<sup>[74]</sup>.

## CORONAVIRUS-19 AND MDSCS

In this age of coronavirus-19 (COVID-19), we would be remiss not to address what is currently known about activation of the host's immune response, in particular MDSCs, by severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2). Plasmacytoid DCs (pDCs) are the major source of tissue-derived type 1 IFNs in response to tissue antigens and activation of DAMPs. Typically, intracellular Toll-like receptors (TLRs) found on endosomes that mediate this pathway evolve to defend the cell against viral pathogens<sup>[75]</sup>. Thus, we queried whether coronavirus infections might be modulated by pDCs residing in the GI tract. Apparently, SARS-CoV-2 induces a massive anti-viral response by secretion of IFN- $\alpha$  from pDCs *via* TLR7<sup>[76]</sup>. The severity of COVID-19 infection might correlate with the activation of endosomal TLRs on pDCs in the GI tract and increased myeloid cell polarization to MDSCs. If this occurs, the cellular immune response to the virus could be rendered ineffective, suggesting that those with severe disease exhibit higher levels of MDSCs than those with mild disease. Indeed, the limited studies available of COVID-19 patients have shown that the MDSC population expands in those with severe disease<sup>[77,78]</sup>. Although T cell exhaustion from

**Table 2 Potential therapeutic strategy for targeting myeloid-derived suppressor cells**

Strategy	Agents
Blocking TDFs from being produced or from reaching the bone marrow Key cytokines, such as IL-6 or S100A8/A9, could be directly targeted <sup>[82,84]</sup>	Targeting the IL-6 receptor (tocilizumab) <sup>[83]</sup>
Inhibiting generation of MDSCs from bone marrow progenitors or inducing apoptosis of circulating MDSCs <sup>[6]</sup>	Gemcitabine, 5-fluorouracil, sunitinib, and zoledronate <sup>[84]</sup>
Preventing trafficking of myeloid cells from the marrow to peripheral lymphoid organs or to the tumor microenvironment <sup>[6]</sup>	Drugs targeting chemokines CXCR2, CXCR4, and CSF1R <sup>[14]</sup>
Directly blocking MDSC suppression of T cells <sup>[85]</sup>	Phosphodiesterase type 5 inhibitors, <i>e.g.</i> , sildenafil and tadalafil, or cyclooxygenase 2 inhibitors <sup>[63]</sup>
Drugs that would promote differentiation of MDSCs into proficient antigen-presenting cells that can stimulate tumor-specific T cells and/or into mature leukocytes <sup>[85]</sup>	All-trans retinoic acid, vitamin D3, and the DNA-methylating agent 5-azacytidine <sup>[85]</sup>

MDSCs: Myeloid-derived suppressor cells; CSF1R: Colony-stimulating factor 1 receptor; CXCR2: C-X-C motif receptor 2; CXCR4: C-X-C motif receptor 4; IL-6: Interleukin-6; TDFs: Tumor-derived factors.

the cytokine storm that COVID-19 patients display is the leading cause for severe disease, massive production of immune suppressor cells also explains the lymphopenia occurring in many of these patients<sup>[79,80]</sup>. A better understanding of what controls MDSCs will facilitate not only therapeutic treatments but may ultimately help to predict who will respond to vaccination. Whether patients who recover from these infections are predisposed to chronic disorders, such as autoimmune diseases and cancer, will require long-term follow up of these patients over decades.

## CONCLUSION

This review article provides a better understanding of the role and mechanism of action of MDSCs in GI malignancies. MDSCs are one of the most important elements in the TME. In patients with GI cancer, MDSCs can lead to immunosuppression, and they play an important role in premalignant cell transformation, tumor growth, and metastasis. A higher number of MDSCs at tumor sites and peripheral blood is correlated with higher mortality rates, risk of relapse, and tumor progression. Therefore, monitoring circulating MDSC levels might have prognostic and predictive value in patients with GI malignancies. The benefit of targeting treatment against MDSCs as a combination therapy has been shown. Consequently, a better comprehension of the role and mechanism of action of MDSCs in the TME may aid in the development of novel immune-targeted therapies. Further prospective studies are needed to understand the characterization and clinical value of MDSCs and more selective anti-MDSC therapies with improved therapeutic outcomes.

## REFERENCES

1. **Arnold M**, Abnet CC, Neale RE, Vignat J, Giovannucci EL, McGlynn KA, Bray F. Global Burden of 5 Major Types of Gastrointestinal Cancer. *Gastroenterology* 2020; **159**: 335-349. e15 [PMID: 32247694 DOI: 10.1053/j.gastro.2020.02.068]
2. **Duffy A**, Zhao F, Haile L, Gamrekashvili J, Fioravanti S, Ma C, Kapanadze T, Compton K, Figg WD, Greten TF. Comparative analysis of monocytic and granulocytic myeloid-derived suppressor cell subsets in patients with gastrointestinal malignancies. *Cancer Immunol Immunother* 2013; **62**: 299-307 [PMID: 23011590 DOI: 10.1007/s00262-012-1332-3]
3. **Lindau D**, Gielen P, Kroesen M, Wesseling P, Adema GJ. The immunosuppressive tumour network: myeloid-derived suppressor cells, regulatory T cells and natural killer T cells. *Immunology* 2013; **138**: 105-115 [PMID: 23216602 DOI: 10.1111/imm.12036]
4. **Wilcox RA**. Cancer-associated myeloproliferation: old association, new therapeutic target. *Mayo Clin Proc* 2010; **85**: 656-663 [PMID: 20592171 DOI: 10.4065/mcp.2010.0077]
5. **Gabitass RF**, Annels NE, Stocken DD, Pandha HA, Middleton GW. Elevated myeloid-derived suppressor cells in pancreatic, esophageal and gastric cancer are an independent prognostic factor and are associated with significant elevation of the Th2 cytokine interleukin-13. *Cancer Immunol Immunother* 2011; **60**: 1419-1430 [PMID: 21644036 DOI: 10.1007/s00262-011-1028-0]
6. **Gabrilovich DI**. Myeloid-Derived Suppressor Cells. *Cancer Immunol Res* 2017; **5**: 3-8 [PMID: 28111111 DOI: 10.1158/2156-8474.CCR16-0105]

- 28052991 DOI: [10.1158/2326-6066.CIR-16-0297](https://doi.org/10.1158/2326-6066.CIR-16-0297)]
- 7 **Ostrand-Rosenberg S**, Sinha P. Myeloid-derived suppressor cells: linking inflammation and cancer. *J Immunol* 2009; **182**: 4499-4506 [PMID: [19342621](https://pubmed.ncbi.nlm.nih.gov/19342621/) DOI: [10.4049/jimmunol.0802740](https://doi.org/10.4049/jimmunol.0802740)]
- 8 **Wang L**, Chang EW, Wong SC, Ong SM, Chong DQ, Ling KL. Increased myeloid-derived suppressor cells in gastric cancer correlate with cancer stage and plasma S100A8/A9 proinflammatory proteins. *J Immunol* 2013; **190**: 794-804 [PMID: [23248262](https://pubmed.ncbi.nlm.nih.gov/23248262/) DOI: [10.4049/jimmunol.1202088](https://doi.org/10.4049/jimmunol.1202088)]
- 9 **Gao XH**, Tian L, Wu J, Ma XL, Zhang CY, Zhou Y, Sun YF, Hu B, Qiu SJ, Zhou J, Fan J, Guo W, Yang XR. Circulating CD14<sup>+</sup> HLA-DR<sup>-low</sup> myeloid-derived suppressor cells predicted early recurrence of hepatocellular carcinoma after surgery. *Hepatol Res* 2017; **47**: 1061-1071 [PMID: [27764536](https://pubmed.ncbi.nlm.nih.gov/27764536/) DOI: [10.1111/hepr.12831](https://doi.org/10.1111/hepr.12831)]
- 10 **Hirbod-Mobarakeh A**, Mirghorbani M, Hajiju F, Marvi M, Bashiri K, Rezaei N. Myeloid-derived suppressor cells in gastrointestinal cancers: A systematic review. *J Gastroenterol Hepatol* 2016; **31**: 1246-1256 [PMID: [26729006](https://pubmed.ncbi.nlm.nih.gov/26729006/) DOI: [10.1111/jgh.13284](https://doi.org/10.1111/jgh.13284)]
- 11 **Wang PF**, Song SY, Wang TJ, Ji WJ, Li SW, Liu N, Yan CX. Prognostic role of pretreatment circulating MDSCs in patients with solid malignancies: A meta-analysis of 40 studies. *Oncoimmunology* 2018; **7**: e1494113 [PMID: [30288362](https://pubmed.ncbi.nlm.nih.gov/30288362/) DOI: [10.1080/2162402X.2018.1494113](https://doi.org/10.1080/2162402X.2018.1494113)]
- 12 **Shibata M**, Gonda K, Nakajima T, Matsumoto Y, Nakamura I, Ohki S, Ohtake T, Kumamoto K, Shimura T, Takenoshita S, Abe N, Momma T. Pretreatment serum levels of circulating myeloid-derived suppressor cells (MDSC) as a prognostic indicator in patients with gastrointestinal cancer. In ASCO Annual Meeting 2014; Chicago
- 13 **Veglia F**, Perego M, Gabrilovich D. Myeloid-derived suppressor cells coming of age. *Nat Immunol* 2018; **19**: 108-119 [PMID: [29348500](https://pubmed.ncbi.nlm.nih.gov/29348500/) DOI: [10.1038/s41590-017-0022-x](https://doi.org/10.1038/s41590-017-0022-x)]
- 14 **Gabrilovich DI**, Ostrand-Rosenberg S, Bronte V. Coordinated regulation of myeloid cells by tumours. *Nat Rev Immunol* 2012; **12**: 253-268 [PMID: [22437938](https://pubmed.ncbi.nlm.nih.gov/22437938/) DOI: [10.1038/nri3175](https://doi.org/10.1038/nri3175)]
- 15 **Dumitru CA**, Moses K, Trellakis S, Lang S, Brandau S. Neutrophils and granulocytic myeloid-derived suppressor cells: immunophenotyping, cell biology and clinical relevance in human oncology. *Cancer Immunol Immunother* 2012; **61**: 1155-1167 [PMID: [22692756](https://pubmed.ncbi.nlm.nih.gov/22692756/) DOI: [10.1007/s00262-012-1294-5](https://doi.org/10.1007/s00262-012-1294-5)]
- 16 **Barreda DR**, Hanington PC, Belosevic M. Regulation of myeloid development and function by colony stimulating factors. *Dev Comp Immunol* 2004; **28**: 509-554 [PMID: [15062647](https://pubmed.ncbi.nlm.nih.gov/15062647/) DOI: [10.1016/j.jci.2003.09.010](https://doi.org/10.1016/j.jci.2003.09.010)]
- 17 **Merad M**, Sathe P, Helft J, Miller J, Mortha A. The dendritic cell lineage: ontogeny and function of dendritic cells and their subsets in the steady state and the inflamed setting. *Annu Rev Immunol* 2013; **31**: 563-604 [PMID: [23516985](https://pubmed.ncbi.nlm.nih.gov/23516985/) DOI: [10.1146/annurev-immunol-020711-074950](https://doi.org/10.1146/annurev-immunol-020711-074950)]
- 18 **Marvel D**, Gabrilovich DI. Myeloid-derived suppressor cells in the tumor microenvironment: expect the unexpected. *J Clin Invest* 2015; **125**: 3356-3364 [PMID: [26168215](https://pubmed.ncbi.nlm.nih.gov/26168215/) DOI: [10.1172/JCI80005](https://doi.org/10.1172/JCI80005)]
- 19 **Chornoguz O**, Grmai L, Sinha P, Artemenko KA, Zubarev RA, Ostrand-Rosenberg S. Proteomic pathway analysis reveals inflammation increases myeloid-derived suppressor cell resistance to apoptosis. *Mol Cell Proteomics* 2011; **10**: M110. 2980; 00 [PMID: [21191032](https://pubmed.ncbi.nlm.nih.gov/21191032/) DOI: [10.1074/mcp.M110.002980](https://doi.org/10.1074/mcp.M110.002980)]
- 20 **Talmadge JE**, Gabrilovich DI. History of myeloid-derived suppressor cells. *Nat Rev Cancer* 2013; **13**: 739-752 [PMID: [24060865](https://pubmed.ncbi.nlm.nih.gov/24060865/) DOI: [10.1038/nrc3581](https://doi.org/10.1038/nrc3581)]
- 21 **Khaled YS**, Ammori BJ, Elkord E. Myeloid-derived suppressor cells in cancer: recent progress and prospects. *Immunol Cell Biol* 2013; **91**: 493-502 [PMID: [23797066](https://pubmed.ncbi.nlm.nih.gov/23797066/) DOI: [10.1038/icb.2013.29](https://doi.org/10.1038/icb.2013.29)]
- 22 **Cao Y**, Feng Y, Zhang Y, Zhu X, Jin F. L-Arginine supplementation inhibits the growth of breast cancer by enhancing innate and adaptive immune responses mediated by suppression of MDSCs in vivo. *BMC Cancer* 2016; **16**: 343 [PMID: [27246354](https://pubmed.ncbi.nlm.nih.gov/27246354/) DOI: [10.1186/s12885-016-2376-0](https://doi.org/10.1186/s12885-016-2376-0)]
- 23 **Rodriguez PC**, Ernstoff MS, Hernandez C, Atkins M, Zabaleta J, Sierra R, Ochoa AC. Arginase I-producing myeloid-derived suppressor cells in renal cell carcinoma are a subpopulation of activated granulocytes. *Cancer Res* 2009; **69**: 1553-1560 [PMID: [19201693](https://pubmed.ncbi.nlm.nih.gov/19201693/) DOI: [10.1158/0008-5472.CAN-08-1921](https://doi.org/10.1158/0008-5472.CAN-08-1921)]
- 24 **Goh CC**, Roggerson KM, Lee HC, Golden-Mason L, Rosen HR, Hahn YS. Hepatitis C Virus-Induced Myeloid-Derived Suppressor Cells Suppress NK Cell IFN- $\gamma$  Production by Altering Cellular Metabolism via Arginase-1. *J Immunol* 2016; **196**: 2283-2292 [PMID: [26826241](https://pubmed.ncbi.nlm.nih.gov/26826241/) DOI: [10.4049/jimmunol.1501881](https://doi.org/10.4049/jimmunol.1501881)]
- 25 **Sinha P**, Chornoguz O, Clements VK, Artemenko KA, Zubarev RA, Ostrand-Rosenberg S. Myeloid-derived suppressor cells express the death receptor Fas and apoptose in response to T cell-expressed FasL. *Blood* 2011; **117**: 5381-5390 [PMID: [21450901](https://pubmed.ncbi.nlm.nih.gov/21450901/) DOI: [10.1182/blood-2010-11-321752](https://doi.org/10.1182/blood-2010-11-321752)]
- 26 **Movahedi K**, Guillems M, Van den Bossche J, Van den Bergh R, Gysemans C, Beschijn A, De Baetselier P, Van Ginderachter JA. Identification of discrete tumor-induced myeloid-derived suppressor cell subpopulations with distinct T cell-suppressive activity. *Blood* 2008; **111**: 4233-4244 [PMID: [18272812](https://pubmed.ncbi.nlm.nih.gov/18272812/) DOI: [10.1182/blood-2007-07-099226](https://doi.org/10.1182/blood-2007-07-099226)]
- 27 **Huang B**, Pan PY, Li Q, Sato AI, Levy DE, Bromberg J, Divino CM, Chen SH. Gr-1+CD115+ immature myeloid suppressor cells mediate the development of tumor-induced T regulatory cells and T-cell anergy in tumor-bearing host. *Cancer Res* 2006; **66**: 1123-1131 [PMID: [16424049](https://pubmed.ncbi.nlm.nih.gov/16424049/) DOI: [10.1158/0008-5472.CAN-05-1299](https://doi.org/10.1158/0008-5472.CAN-05-1299)]
- 28 **Goedeghebuere P**, Mitchem JB, Porembka MR, Tan MC, Belt BA, Wang-Gillam A, Gillanders WE, Hawkins WG, Linehan DC. Myeloid-derived suppressor cells: general characteristics and relevance to clinical management of pancreatic cancer. *Curr Cancer Drug Targets* 2011; **11**: 734-751 [PMID: [21191032](https://pubmed.ncbi.nlm.nih.gov/21191032/) DOI: [10.1089/cdr.2010.0005](https://doi.org/10.1089/cdr.2010.0005)]



- 21599634 DOI: [10.2174/156800911796191024](https://doi.org/10.2174/156800911796191024)]
- 29 **Fu J**, Xu D, Liu Z, Shi M, Zhao P, Fu B, Zhang Z, Yang H, Zhang H, Zhou C, Yao J, Jin L, Wang H, Yang Y, Fu YX, Wang FS. Increased regulatory T cells correlate with CD8 T-cell impairment and poor survival in hepatocellular carcinoma patients. *Gastroenterology* 2007; **132**: 2328-2339 [PMID: [17570208](https://pubmed.ncbi.nlm.nih.gov/17570208/) DOI: [10.1053/j.gastro.2007.03.102](https://doi.org/10.1053/j.gastro.2007.03.102)]
  - 30 **Hiraoka N**, Onozato K, Kosuge T, Hirohashi S. Prevalence of FOXP3<sup>+</sup> regulatory T cells increases during the progression of pancreatic ductal adenocarcinoma and its premalignant lesions. *Clin Cancer Res* 2006; **12**: 5423-5434 [PMID: [17000676](https://pubmed.ncbi.nlm.nih.gov/17000676/) DOI: [10.1158/1078-0432.CCR-06-0369](https://doi.org/10.1158/1078-0432.CCR-06-0369)]
  - 31 **Mucha J**, Majchrzak K, Taciak B, Hellmén E, Król M. MDSCs mediate angiogenesis and predispose canine mammary tumor cells for metastasis via IL-28/IL-28RA (IFN- $\lambda$ ) signaling. *PLoS One* 2014; **9**: e103249 [PMID: [25075523](https://pubmed.ncbi.nlm.nih.gov/25075523/) DOI: [10.1371/journal.pone.0103249](https://doi.org/10.1371/journal.pone.0103249)]
  - 32 **Nagaraj S**, Gabrilovich DI. Myeloid-derived suppressor cells in human cancer. *Cancer J* 2010; **16**: 348-353 [PMID: [20693846](https://pubmed.ncbi.nlm.nih.gov/20693846/) DOI: [10.1097/PP0.0b013e3181eb3358](https://doi.org/10.1097/PP0.0b013e3181eb3358)]
  - 33 **Kotsakis A**, Harasymczuk M, Schilling B, Georgoulas V, Argiris A, Whiteside TL. Myeloid-derived suppressor cell measurements in fresh and cryopreserved blood samples. *J Immunol Methods* 2012; **381**: 14-22 [PMID: [22522114](https://pubmed.ncbi.nlm.nih.gov/22522114/) DOI: [10.1016/j.jim.2012.04.004](https://doi.org/10.1016/j.jim.2012.04.004)]
  - 34 **Marini O**, Spina C, Mimiola E, Cassaro A, Malerba G, Todeschini G, Perbellini O, Scupoli M, Carli G, Facchinelli D, Cassatella M, Scapini P, Tecchio C. Identification of granulocytic myeloid-derived suppressor cells (G-MDSCs) in the peripheral blood of Hodgkin and non-Hodgkin lymphoma patients. *Oncotarget* 2016; **7**: 27676-27688 [PMID: [27050283](https://pubmed.ncbi.nlm.nih.gov/27050283/) DOI: [10.18632/oncotarget.8507](https://doi.org/10.18632/oncotarget.8507)]
  - 35 **Broz ML**, Krummel MF. The emerging understanding of myeloid cells as partners and targets in tumor rejection. *Cancer Immunol Res* 2015; **3**: 313-319 [PMID: [25847968](https://pubmed.ncbi.nlm.nih.gov/25847968/) DOI: [10.1158/2326-6066.CIR-15-0041](https://doi.org/10.1158/2326-6066.CIR-15-0041)]
  - 36 **Jiao ZJ**, Gao JJ, Hua SH, Chen DY, Wang WH, Wang H, Wang XH, Xu HX. Correlation between circulating myeloid-derived suppressor cells and Th17 cells in esophageal cancer. *World J Gastroenterol* 2012; **18**: 5454-5461 [PMID: [23082063](https://pubmed.ncbi.nlm.nih.gov/23082063/) DOI: [10.3748/wjg.v18.i38.5454](https://doi.org/10.3748/wjg.v18.i38.5454)]
  - 37 **Xu J**, Peng Y, Yang M, Guo N, Liu H, Gao H, Niu F, Wang R, Wang C, Yu K. Increased levels of myeloid-derived suppressor cells in esophageal cancer patients is associated with the complication of sepsis. *Biomed Pharmacother* 2020; **125**: 109864 [PMID: [32007915](https://pubmed.ncbi.nlm.nih.gov/32007915/) DOI: [10.1016/j.biopha.2020.109864](https://doi.org/10.1016/j.biopha.2020.109864)]
  - 38 **Chen MF**, Kuan FC, Yen TC, Lu MS, Lin PY, Chung YH, Chen WC, Lee KD. IL-6-stimulated CD11b<sup>+</sup> CD14<sup>+</sup> HLA-DR<sup>+</sup> myeloid-derived suppressor cells, are associated with progression and poor prognosis in squamous cell carcinoma of the esophagus. *Oncotarget* 2014; **5**: 8716-8728 [PMID: [25238263](https://pubmed.ncbi.nlm.nih.gov/25238263/) DOI: [10.18632/oncotarget.2368](https://doi.org/10.18632/oncotarget.2368)]
  - 39 **Aparicio-Siegmund S**, Sommer J, Monhasery N, Schwanbeck R, Keil E, Finkstädt D, Pfeffer K, Rose-John S, Scheller J, Garbers C. Inhibition of protein kinase II (CK2) prevents induced signal transducer and activator of transcription (STAT) 1/3 and constitutive STAT3 activation. *Oncotarget* 2014; **5**: 2131-2148 [PMID: [24742922](https://pubmed.ncbi.nlm.nih.gov/24742922/) DOI: [10.18632/oncotarget.1852](https://doi.org/10.18632/oncotarget.1852)]
  - 40 **Bray F**, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018; **68**: 394-424 [PMID: [30207593](https://pubmed.ncbi.nlm.nih.gov/30207593/) DOI: [10.3322/caac.21492](https://doi.org/10.3322/caac.21492)]
  - 41 **Li N**, Han D, Sun J, Li Y, Zhang J, Zhang Y, Liu M, Peng R, Wang H, Zhang Z, Wang J, Liu Z, Ma J. Subtypes of MDSCs in mechanisms and prognosis of gastric cancer and are inhibited by epirubicin and paclitaxel. *Discov Med* 2018; **25**: 99-112 [PMID: [29641972](https://pubmed.ncbi.nlm.nih.gov/29641972/)]
  - 42 **Gonda K**, Kanke Y, Yazawa T, Monma T, Nakamura SS, Ohki S, Shimura T, Ohto H, Takenoshita S. Circulating myeloid-derived suppressor cells (MDSC) and correlation to poor prognosis, Th2-polarization, inflammation, and nutritional damages in patients with gastric cancer. *J Clin Oncol* 2013; **31**: 3063-3063
  - 43 **Ding L**, Hayes MM, Photenhauer A, Eaton KA, Li Q, Ocadiz-Ruiz R, Merchant JL. Schlafen 4-expressing myeloid-derived suppressor cells are induced during murine gastric metaplasia. *J Clin Invest* 2016; **126**: 2867-2880 [PMID: [27427984](https://pubmed.ncbi.nlm.nih.gov/27427984/) DOI: [10.1172/JCI82529](https://doi.org/10.1172/JCI82529)]
  - 44 **Merchant JL**, Ding L. Hedgehog Signaling Links Chronic Inflammation to Gastric Cancer Precursor Lesions. *Cell Mol Gastroenterol Hepatol* 2017; **3**: 201-210 [PMID: [28275687](https://pubmed.ncbi.nlm.nih.gov/28275687/) DOI: [10.1016/j.jcmgh.2017.01.004](https://doi.org/10.1016/j.jcmgh.2017.01.004)]
  - 45 **Companioni Nápoles O**, Tsao AC, Sanz-Anquela JM, Sala N, Bonet C, Pardo ML, Ding L, Simo O, Saqui-Salces M, Blanco VP, Gonzalez CA, Merchant JL. SCHLAFEN 5 expression correlates with intestinal metaplasia that progresses to gastric cancer. *J Gastroenterol* 2017; **52**: 39-49 [PMID: [27032393](https://pubmed.ncbi.nlm.nih.gov/27032393/) DOI: [10.1007/s00535-016-1202-4](https://doi.org/10.1007/s00535-016-1202-4)]
  - 46 **Ding L**, Li Q, Chakrabarti J, Munoz A, Faure-Kumar E, Ocadiz-Ruiz R, Razumilava N, Zhang G, Hayes MH, Sontz RA, Mendoza ZE, Mahurkar S, Greenson JK, Perez-Perez G, Hanh NTH, Zavros Y, Samuelson LC, Iliopoulos D, Merchant JL. MiR130b from Schlafen4<sup>+</sup> MDSCs stimulates epithelial proliferation and correlates with preneoplastic changes prior to gastric cancer. *Gut* 2020; **69**: 1750-1761 [PMID: [31980446](https://pubmed.ncbi.nlm.nih.gov/31980446/) DOI: [10.1136/gutjnl-2019-318817](https://doi.org/10.1136/gutjnl-2019-318817)]
  - 47 **Elwan N**, Salem ML, Kobtan A, El-Kalla F, Mansour L, Yousef M, Al-Sabbagh A, Zidan AA, Abd-El salam S. High numbers of myeloid derived suppressor cells in peripheral blood and ascitic fluid of cirrhotic and HCC patients. *Immunol Invest* 2018; **47**: 169-180 [PMID: [29182438](https://pubmed.ncbi.nlm.nih.gov/29182438/) DOI: [10.1080/08820139.2017.1407787](https://doi.org/10.1080/08820139.2017.1407787)]
  - 48 **Arihara F**, Mizukoshi E, Kitahara M, Takata Y, Arai K, Yamashita T, Nakamoto Y, Kaneko S. Increase in CD14<sup>+</sup>HLA-DR<sup>+</sup>/low myeloid-derived suppressor cells in hepatocellular carcinoma



- patients and its impact on prognosis. *Cancer Immunol Immunother* 2013; **62**: 1421-1430 [PMID: 23764929 DOI: 10.1007/s00262-013-1447-1]
- 49 **Mizukoshi E**, Yamashita T, Arai K, Terashima T, Kitahara M, Nakagawa H, Iida N, Fushimi K, Kaneko S. Myeloid-derived suppressor cells correlate with patient outcomes in hepatic arterial infusion chemotherapy for hepatocellular carcinoma. *Cancer Immunol Immunother* 2016; **65**: 715-725 [PMID: 27083166 DOI: 10.1007/s00262-016-1837-2]
- 50 **Bayik D LA**, Roversi GA, Serbinowski E, Acevedo-Moreno L, Lanigan C, Orujov M, Lo A, Alban TJ, Silver DJ, Brown JM, Allende DS, Aucejo FN, Lathia JD. Cholangiocarcinoma presents a distinct myeloid-derived suppressor cell signature compared to other hepatobiliary cancers. *bioRxiv* 2019 [DOI: 10.1101/554600]
- 51 **Aggarwal N**, Sloane BF. Cathepsin B: multiple roles in cancer. *Proteomics Clin Appl* 2014; **8**: 427-437 [PMID: 24677670 DOI: 10.1002/prca.201300105]
- 52 **Roderfeld M**, Rath T, Lammert F, Dierkes C, Graf J, Roeb E. Innovative immunohistochemistry identifies MMP-9 expressing macrophages at the invasive front of murine HCC. *World J Hepatol* 2010; **2**: 175-179 [PMID: 21160992 DOI: 10.4254/wjh.v2.i5.175]
- 53 **Groot VP**, Rezaee N, Wu W, Cameron JL, Fishman EK, Hruban RH, Weiss MJ, Zheng L, Wolfgang CL, He J. Patterns, Timing, and Predictors of Recurrence Following Pancreatectomy for Pancreatic Ductal Adenocarcinoma. *Ann Surg* 2018; **267**: 936-945 [PMID: 28338509 DOI: 10.1097/SLA.0000000000002234]
- 54 **Khaled YS**, Ammori BJ, Elkord E. Increased levels of granulocytic myeloid-derived suppressor cells in peripheral blood and tumour tissue of pancreatic cancer patients. *J Immunol Res* 2014; **2014**: 879897 [PMID: 24741628 DOI: 10.1155/2014/879897]
- 55 **Ma P**, Beatty PL, McKolanis J, Brand R, Schoen RE, Finn OJ. Circulating Myeloid Derived Suppressor Cells (MDSC) That Accumulate in Premalignancy Share Phenotypic and Functional Characteristics With MDSC in Cancer. *Front Immunol* 2019; **10**: 1401 [PMID: 31275327 DOI: 10.3389/fimmu.2019.01401]
- 56 **Mace TA**, Ameen Z, Collins A, Wojcik S, Mair M, Young GS, Fuchs JR, Eubank TD, Frankel WL, Bekaii-Saab T, Bloomston M, Lesinski GB. Pancreatic cancer-associated stellate cells promote differentiation of myeloid-derived suppressor cells in a STAT3-dependent manner. *Cancer Res* 2013; **73**: 3007-3018 [PMID: 23514705 DOI: 10.1158/0008-5472.CAN-12-4601]
- 57 **OuYang LY**, Wu XJ, Ye SB, Zhang RX, Li ZL, Liao W, Pan ZZ, Zheng LM, Zhang XS, Wang Z, Li Q, Ma G, Li J. Tumor-induced myeloid-derived suppressor cells promote tumor progression through oxidative metabolism in human colorectal cancer. *J Transl Med* 2015; **13**: 47 [PMID: 25638150 DOI: 10.1186/s12967-015-0410-7]
- 58 **Tada K**, Kitano S, Shoji H, Nishimura T, Shimada Y, Nagashima K, Aoki K, Hiraoka N, Honma Y, Iwasa S, Okita N, Takashima A, Kato K, Yamada Y, Katayama N, Boku N, Heike Y, Hamaguchi T. Pretreatment Immune Status Correlates with Progression-Free Survival in Chemotherapy-Treated Metastatic Colorectal Cancer Patients. *Cancer Immunol Res* 2016; **4**: 592-599 [PMID: 27197061 DOI: 10.1158/2326-6066.CIR-15-0298]
- 59 **Lu LC**, Chang CJ, Hsu CH. Targeting myeloid-derived suppressor cells in the treatment of hepatocellular carcinoma: current state and future perspectives. *J Hepatocell Carcinoma* 2019; **6**: 71-84 [PMID: 31123667 DOI: 10.2147/JHC.S159693]
- 60 **Fultang L**, Panetti S, Ng M, Collins P, Graef S, Rizkalla N, Booth S, Lenton R, Noyvert B, Shannon-Lowe C, Middleton G, Mussai F, De Santo C. MDSC targeting with Gemtuzumab ozogamicin restores T cell immunity and immunotherapy against cancers. *EBioMedicine* 2019; **47**: 235-246 [PMID: 31462392 DOI: 10.1016/j.ebiom.2019.08.025]
- 61 **Wang Z**, Liu Y, Zhang Y, Shang Y, Gao Q. MDSC-decreasing chemotherapy increases the efficacy of cytokine-induced killer cell immunotherapy in metastatic renal cell carcinoma and pancreatic cancer. *Oncotarget* 2016; **7**: 4760-4769 [PMID: 26716894 DOI: 10.18632/oncotarget.6734]
- 62 **Jiang J**, Xu N, Wu C, Deng H, Lu M, Li M, Xu B, Wu J, Wang R, Xu J, Nilsson-Ehle P. Treatment of advanced gastric cancer by chemotherapy combined with autologous cytokine-induced killer cells. *Anticancer Res* 2006; **26**: 2237-2242 [PMID: 16821594]
- 63 **Serafini P**, Meckel K, Kelso M, Noonan K, Califano J, Koch W, Dolcetti L, Bronte V, Borrello I. Phosphodiesterase-5 inhibition augments endogenous antitumor immunity by reducing myeloid-derived suppressor cell function. *J Exp Med* 2006; **203**: 2691-2702 [PMID: 17101732 DOI: 10.1084/jem.20061104]
- 64 **Lin S**, Wang J, Wang L, Wen J, Guo Y, Qiao W, Zhou J, Xu G, Zhi F. Phosphodiesterase-5 inhibition suppresses colonic inflammation-induced tumorigenesis via blocking the recruitment of MDSC. *Am J Cancer Res* 2017; **7**: 41-52 [PMID: 28123846]
- 65 **Yu SJ**, Ma C, Heinrich B, Brown ZJ, Sandhu M, Zhang Q, Fu Q, Agdashian D, Rosato U, Korangy F, Greten TF. Targeting the crosstalk between cytokine-induced killer cells and myeloid-derived suppressor cells in hepatocellular carcinoma. *J Hepatol* 2019; **70**: 449-457 [PMID: 30414862 DOI: 10.1016/j.jhep.2018.10.040]
- 66 **Rawat D**, Koiri RK. Tadalafil inhibits elevated glutamic oxaloacetic transaminase during alcohol aflatoxin induced hepatocellular carcinoma in rats. *Int J Immunother Cancer Res* 2020; **6**: 10-13 [DOI: 10.17352/2455-8591.000022]
- 67 **Ozao-Choy J**, Ma G, Kao J, Wang GX, Meseck M, Sung M, Schwartz M, Divino CM, Pan PY, Chen SH. The novel role of tyrosine kinase inhibitor in the reversal of immune suppression and modulation of tumor microenvironment for immune-based cancer therapies. *Cancer Res* 2009; **69**: 2514-2522

- [PMID: 19276342 DOI: 10.1158/0008-5472.CAN-08-4709]
- 68 **Mulet-Margalef N**, Garcia-Del-Muro X. Sunitinib in the treatment of gastrointestinal stromal tumor: patient selection and perspectives. *Onco Targets Ther* 2016; **9**: 7573-7582 [PMID: 28008275 DOI: 10.2147/OTT.S101385]
  - 69 **Chen PT**, Hsieh CC, Wu CT, Yen TC, Lin PY, Chen WC, Chen MF. 1 $\alpha$ ,25-Dihydroxyvitamin D3 Inhibits Esophageal Squamous Cell Carcinoma Progression by Reducing IL6 Signaling. *Mol Cancer Ther* 2015; **14**: 1365-1375 [PMID: 25824337 DOI: 10.1158/1535-7163.MCT-14-0952]
  - 70 **Albertini MR**. The age of enlightenment in melanoma immunotherapy. *J Immunother Cancer* 2018; **6**: 80 [PMID: 30134977 DOI: 10.1186/s40425-018-0397-8]
  - 71 **Muro K**, Chung HC, Shankaran V, Geva R, Catenacci D, Gupta S, Eder JP, Golan T, Le DT, Burtneess B, McRee AJ, Lin CC, Pathiraja K, Luncceford J, Emancipator K, Juco J, Koshiji M, Bang YJ. Pembrolizumab for patients with PD-L1-positive advanced gastric cancer (KEYNOTE-012): a multicentre, open-label, phase 1b trial. *Lancet Oncol* 2016; **17**: 717-726 [PMID: 27157491 DOI: 10.1016/S1470-2045(16)00175-3]
  - 72 **Wang ZH**, Shen L. Management of gastrointestinal adverse events induced by immune-checkpoint inhibitors. *Chronic Dis Transl Med* 2018; **4**: 1-7 [PMID: 29756118 DOI: 10.1016/j.cdtm.2017.12.001]
  - 73 **Weber R**, Fleming V, Hu X, Nagibin V, Groth C, Altevogt P, Utikal J, Umansky V. Myeloid-Derived Suppressor Cells Hinder the Anti-Cancer Activity of Immune Checkpoint Inhibitors. *Front Immunol* 2018; **9**: 1310 [PMID: 29942309 DOI: 10.3389/fimmu.2018.01310]
  - 74 **Highfill SL**, Cui Y, Giles AJ, Smith JP, Zhang H, Morse E, Kaplan RN, Mackall CL. Disruption of CXCR2-mediated MDSC tumor trafficking enhances anti-PD1 efficacy. *Sci Transl Med* 2014; **6**: 237ra67 [PMID: 24848257 DOI: 10.1126/scitranslmed.3007974]
  - 75 **Gilliet M**, Cao W, Liu YJ. Plasmacytoid dendritic cells: sensing nucleic acids in viral infection and autoimmune diseases. *Nat Rev Immunol* 2008; **8**: 594-606 [PMID: 18641647 DOI: 10.1038/nri2358]
  - 76 **Cervantes-Barragan L**, Züst R, Weber F, Spiegel M, Lang KS, Akira S, Thiel V, Ludewig B. Control of coronavirus infection through plasmacytoid dendritic-cell-derived type I interferon. *Blood* 2007; **109**: 1131-1137 [PMID: 16985170 DOI: 10.1182/blood-2006-05-023770]
  - 77 **Agrati C**, Sacchi A, Bordoni V, Cimini E, Notari S, Grassi G, Casetti R, Tartaglia E, Lalle E, D'Abramo A, Castilletti C, Marchioni L, Shi Y, Mariano A, Song JW, Zhang JY, Wang FS, Zhang C, Fimia GM, Capobianchi MR, Piacentini M, Antinori A, Nicastrì E, Maeurer M, Zumla A, Ippolito G. Expansion of myeloid-derived suppressor cells in patients with severe coronavirus disease (COVID-19). *Cell Death Differ* 2020; **27**: 3196-3207 [PMID: 32514047 DOI: 10.1038/s41418-020-0572-6]
  - 78 **Bordoni V**, Sacchi A, Cimini E, Notari S, Grassi G, Tartaglia E, Casetti R, Giancola ML, Bevilacqua N, Maeurer M, Zumla A, Locatelli F, De Benedetti F, Palmieri F, Marchioni L, Capobianchi MR, D'Offizi G, Petrosillo N, Antinori A, Nicastrì E, Ippolito G, Agrati C. An Inflammatory Profile Correlates With Decreased Frequency of Cytotoxic Cells in Coronavirus Disease 2019. *Clin Infect Dis* 2020; **71**: 2272-2275 [PMID: 32407466 DOI: 10.1093/cid/ciaa577]
  - 79 **Fathi N**, Rezaei N. Lymphopenia in COVID-19: Therapeutic opportunities. *Cell Biol Int* 2020; **44**: 1792-1797 [PMID: 32458561 DOI: 10.1002/cbin.11403]
  - 80 **Liu J**, Li S, Liu J, Liang B, Wang X, Wang H, Li W, Tong Q, Yi J, Zhao L, Xiong L, Guo C, Tian J, Luo J, Yao J, Pang R, Shen H, Peng C, Liu T, Zhang Q, Wu J, Xu L, Lu S, Wang B, Weng Z, Han C, Zhu H, Zhou R, Zhou H, Chen X, Ye P, Zhu B, Wang L, Zhou W, He S, He Y, Jie S, Wei P, Zhang J, Lu Y, Wang W, Zhang L, Li L, Zhou F, Wang J, Dittmer U, Lu M, Hu Y, Yang D, Zheng X. Longitudinal characteristics of lymphocyte responses and cytokine profiles in the peripheral blood of SARS-CoV-2 infected patients. *EBioMedicine* 2020; **55**: 102763 [PMID: 32361250 DOI: 10.1016/j.ebiom.2020.102763]
  - 81 **Umansky V**, Blattner C, Gebhardt C, Utikal J. The Role of Myeloid-Derived Suppressor Cells (MDSC) in Cancer Progression. *Vaccines (Basel)* 2016; **4**: 36 [PMID: 27827871 DOI: 10.3390/vaccines4040036]
  - 82 **Diaz-Montero CM**, Finke J, Montero AJ. Myeloid-derived suppressor cells in cancer: therapeutic, predictive, and prognostic implications. *Semin Oncol* 2014; **41**: 174-184 [PMID: 24787291 DOI: 10.1053/j.seminoncol.2014.02.003]
  - 83 **Tobin RP**, Jordan KR, Kapoor P, Sponberg E, Davis D, Vorwald VM, Coutts KL, Gao D, Smith DE, Borgers JSW, Robinson S, Amato C, Gonzalez R, Lewis KD, Robinson WA, Borges VF, McCarter MD. IL-6 and IL-8 Are Linked With Myeloid-Derived Suppressor Cell Accumulation and Correlate With Poor Clinical Outcomes in Melanoma Patients. *Front Oncol* 2019; **9**: 1223 [PMID: 31781510 DOI: 10.3389/fonc.2019.01223]
  - 84 **Najjar YG**, Finke JH. Clinical perspectives on targeting of myeloid derived suppressor cells in the treatment of cancer. *Front Oncol* 2013; **3**: 49 [PMID: 23508517 DOI: 10.3389/fonc.2013.00049]
  - 85 **Daurkin I**, Eruslanov E, Vieweg J, Kusmartsev S. Generation of antigen-presenting cells from tumor-infiltrated CD11b myeloid cells with DNA demethylating agent 5-aza-2'-deoxycytidine. *Cancer Immunol Immunother* 2010; **59**: 697-706 [PMID: 19882154 DOI: 10.1007/s00262-009-0786-4]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

