

2-2019

Factors Associated with Stage at Presentation Among Patients with Rare Pancreatic Tumors: An Exploratory Analysis of the SEER Database

J. Subramoney

Thomas Jefferson University, jay.subramoney@jefferson.edu

D. Delgado

Thomas Jefferson University

V. Nguyen

Thomas Jefferson University

R. Denny

Thomas Jefferson University, rachel.denny@jefferson.edu

B. George

*Thomas Jefferson University**See next page for additional authors*

[Let us know how access to this document benefits you](#)

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2021_phase1

 Part of the [Oncology Commons](#)

Recommended Citation

Subramoney, J.; Delgado, D.; Nguyen, V.; Denny, R.; George, B.; and Mitchell, E., "Factors Associated with Stage at Presentation Among Patients with Rare Pancreatic Tumors: An Exploratory Analysis of the SEER Database" (2019). SKMC JeffMD Scholarly Inquiry, Phase 1, Project 1.

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

J. Subramoney; D. Delgado; V. Nguyen; R. Denny; B. George; and E. Mitchell, MD

Factors associated with Stage at Presentation among Patients with Rare Pancreatic Tumors: An Exploratory Analysis of the SEER Database

Authors: J. Subramoney, D. Delgado, V. Nguyen, R. Denny, B. George, E. Mitchell MD
Thomas Jefferson University, Philadelphia, PA, USA

Introduction

Rare pancreatic carcinomas (PC) account for roughly 15% of all pancreatic cancers and approximately 8,300 new cases within the US annually. Little research compares initial stage at diagnosis among rare vs. common pancreatic tumors. The aim of this study was to examine associations between tumor histology and initial stage at diagnosis among cases of rare pancreatic cancers.

Methods

Retrospective cohort study, using the NCI's Surveillance, Epidemiology, and End Results (SEER) 1990-2015 database. We included patients diagnosed with pancreatic cancer aged 18+. Histology was classified into four categories (ductal, carcinoid, mucinous adenocarcinoma, and undetermined) and stage was dichotomized (locoregional versus distant). Multivariate logistic regression was used to describe the association between tumor histology with initial stage at diagnosis, controlling for patient and tumor characteristics.

Results

90,764 PC patients were analyzed: 9.60%(ductal carcinomas), 4.47%(carcinoid), 3.78%(mucinous), and 15.81%(undetermined); 54.74% of cases were distantly metastasized at diagnosis. In multivariate analysis, PC patients with mucinous histology had greater odds of presenting with late-stage disease compared to those with adenocarcinoma (OR=1.41, CI=1.31, 1.52%; carcinoid and undetermined neoplasm findings were not significant (OR=0.99, CI=0.92, 1.06 and OR=1.01, CI=0.97, 1.05 respectively). Other factors associated with distant PC included age 40-59, male sex, and African American background.

Discussion

We observed that mucinous histology has a greater risk of late-stage diagnosis while carcinoid and undetermined neoplasms had a similar risk of late-stage diagnosis vs. adenocarcinoma.

