

Thomas Jefferson University Jefferson Digital Commons

Phase 1

Class of 2022

1-2020

Investigation of the Superman Stretch on Posterior Capsule Range of Motion

Brian Gibbs Thomas Jefferson University, brian.gibbs@jefferson.edu

Christopher M. Aland, MD

Joshua Hornstein, MD

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2022_phase1

Part of the Orthopedics Commons, and the Translational Medical Research Commons Let us know how access to this document benefits you

Recommended Citation

Gibbs, Brian; Aland, MD, Christopher M.; and Hornstein, MD, Joshua, "Investigation of the Superman Stretch on Posterior Capsule Range of Motion" (2020). *Phase 1.* Paper 65. https://jdc.jefferson.edu/si_ctr_2022_phase1/65

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's Center for Teaching and Learning (CTL). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson Scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

SI/CTR Abstract

Word count: 250 words

Investigation of the Superman Stretch on Posterior Capsule Range of Motion Brian Gibbs, Christopher M. Aland, MD*, Joshua Hornstein, MD

Introduction: Shoulder pain is a common clinical entity affecting all age groups and can frequently be attributed to pathological internal impingement. A primary contributing factor to labrum tears and internal impingement is posterior capsule tightness. In order to improve range of motion (R.O.M.), prior studies have demonstrated that stretching is an important tool for increasing R.O.M. at many joints. We hypothesized that the superman stretch would lead to greater increases in IR than the traditional sleeper stretch.

Methods: A level II, prospective study in collegiate-level athletes from The College of New Jersey (TCNJ) was performed. Shoulder internal rotation (IR), shoulder external rotation (ER), and glenohumeral (GH) horizontal adduction were measured. Students were randomized to perform the sleeper stretch (control) or superman stretch (experimental). Measurements were taken with a digital goniometer. P-values were calculated with a two-sample z-test.

Results: Both stretches produced significant differences in pre- and post-stretch R.O.M. across all outcomes measured (IR, ER, adduction). The superman stretch appears to provide an increase in IR (p-value 0.196) and in horizontal adduction (p-value 0.0731), whereas the sleeper stretch appears to provide an increase in ER (p-value 0.343).

Discussion: Though not statistically significant, our results indicate that the superman stretch may provide an increase in IR and horizontal adduction when compared to the sleeper stretch. A larger sample size is required to determine more significant values. Demonstration of significant improvements in shoulder R.O.M. would allow us to begin to investigate the effects of posterior capsule stretching on overhead athletes experiencing shoulder pain.