

2-1-2016

Evaluation of a PACAP Peptide Analogue Labeled with (68)Ga Using Two Different Chelating Agents.

Pardeep Kumar

Thomas Jefferson University, pardeep.kumar@jefferson.edu

Sushil Kumar Tripathi

Thomas Jefferson University, sushil.tripathi@jefferson.edu

Chang-Po Chen

Henan Normal University

Neil Mehta

Thomas Jefferson University, Neil.Mehta@jefferson.edu

Bishnuhari Paudyal

*Thomas Jefferson University, Bishnuhari.Paudyal@jefferson.edu**See next page for additional authors*

[Let us know how access to this document benefits you](#)

Follow this and additional works at: <https://jdc.jefferson.edu/radiologyfp> Part of the [Radiology Commons](#)

Recommended Citation

Kumar, Pardeep; Tripathi, Sushil Kumar; Chen, Chang-Po; Mehta, Neil; Paudyal, Bishnuhari; Wickstrom, Eric; and Thakur, Mathew L., "Evaluation of a PACAP Peptide Analogue Labeled with (68)Ga Using Two Different Chelating Agents." (2016). *Department of Radiology Faculty Papers*. Paper 44.

<https://jdc.jefferson.edu/radiologyfp/44>

Authors

Pardeep Kumar, Sushil Kumar Tripathi, Chang-Po Chen, Neil Mehta, Bishnuhari Paudyal, Eric Wickstrom, and Mathew L. Thakur

Evaluation of a PACAP peptide analogue labeled with ⁶⁸Ga using two different chelating agents.

Pardeep Kumar¹, Sushil Tripathi¹, Chang-Po Chen², Neil Mehta¹, Bishnuhari paudyal¹, Eric Wickstrom^{3,4}, Mathew L. Thakur^{*1,4,5}.

ABSTRACT

Objective: The authors have conjugated chelating agents (DOTA and NODAGA) with a peptide (PACAP analogue) which has a high affinity for VPAC1 receptors expressed on cancer cells. In order to determine a suitable chelating agent for labeling with ⁶⁸Ga, we have compared the labeling kinetics and stability of these peptide conjugates.

Methods: For labeling, ⁶⁸GaCl₃ was eluted in 0.1 M HCl from a [⁶⁸Ge-⁶⁸Ga] generator. The influences of peptide concentration, pH and temperature on the radiolabeling efficiency were studied. The stability was evaluated in saline, human serum, DTPA, Transferrin and metallic ions (FeCl₃, CaCl₂ and ZnCl₂). Cell binding assay was performed using human breast cancer cells (T47D). Tissue biodistribution was studied in normal athymic nude mice.

Results: Optimal radiolabeling (> 95.0%) of the DOTA-peptide conjugates required higher (50°C–90°C) temperature and 10 min incubation at pH 2–5. The NODAGA-peptide conjugate needed incubation only at 25°C for 10 min. Both radio-complexes were stable in saline, serum as well as against transchelation and transmetallation. Cell binding at 37°C for 15 min incubation with ⁶⁸Ga-NODAGA-peptide was 34.0% as compared to 24.5% for ⁶⁸Ga-DOTA-peptide. Tissue biodistribution at 1 hr post-injection of both ⁶⁸Ga labeled peptide conjugates showed clearance through the kidneys.

Conclusions: NODAGA-peptide showed more convenient radiolabeling features than that of DOTA-peptide.

Key words: Breast cancer, Prostate Cancer, chelating agent, peptides, radiopharmaceuticals.

¹Department of Radiology, Thomas Jefferson University, Philadelphia, Pennsylvania. ² School of Chemistry and Chemical Engineering, Henan Normal University, Xinxiang, China. ³Department of Biochemistry and Molecular Biology, ⁴Kimmel Cancer Center, ⁵Department of radiation oncology, Thomas Jefferson University, Philadelphia, Pennsylvania.

Introduction

Positron emission tomography (PET) has become a prominent imaging modality in the field of oncology.¹ PET has several technical advantages over the single photon emission computed tomography (SPECT) like attenuation correction, and higher spatial resolution.²⁻³ Small molecules such as sugars, amino acids, nucleic acids, or receptor-binding ligands are labeled with positron-emitting radionuclides for PET imaging to study *in vivo* visualization of physiological processes on a molecular level.⁴⁻⁷ There are several positron emitting radionuclides like fluorine-18 [¹⁸F] ($t_{1/2} \approx 110$ min), copper-64 [⁶⁴Cu] ($t_{1/2} \approx 12.7$ hr), carbon-11 [¹¹C] ($t_{1/2} \approx 20$ min), and oxygen-15 [¹⁵O] ($t_{1/2} \approx 2$ min). However, their production requires in-house cyclotron, not available at every academic institution and hospital.

In recent years, Gallium-68 [⁶⁸Ga] - a positron emitting radionuclide conveniently available in the form of Germanium-68 ($t_{1/2} \approx 270$ days)-Gallium-68 ($t_{1/2} \approx 68$ min) [⁶⁸Ge \rightarrow ⁶⁸Ga] generator has drawn considerable attention for oncologic diagnostic applications. The short physical half-life of ⁶⁸Ga induces low radiation burden on patients and makes it an ideal radionuclide for diagnostic use.⁸⁻¹⁰ The labeling of biomolecules with ⁶⁸Ga is achieved by using bi-functional chelators (BFC) such as DTPA (diethylene triamine pentaacetic acid), DOTA (1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid), NOTA (1,4,7-triazacyclononane-1,4,7-triacetic acid), or NODAGA (1,4,7-triazacyclononane-1-glutamic acid-4,7-diacetic acid). BFCs differ from each other in their physio-chemical properties which can influence radiolabeling kinetics. They can also influence overall charge of the complex, biodistribution and more importantly uptake in the tumor site.¹¹⁻¹³

In recent years, there has been increasing interest in the receptor targeted scintigraphy in which peptides have emerged as promising biomolecules.¹⁴ Their properties such as fast clearance, rapid tissue penetration, low antigenicity, and relatively easy synthesis have added to their applicability.¹⁵ In past few years, our laboratory has designed, synthesized and radiolabeled peptide conjugates which have high affinity for VPAC-1 receptors, over-expressed in many malignant tumors including those of the breast and prostate.¹⁶⁻²⁰ The peptides are analogues of Pituitary adenylate cyclase-activating peptide (PACAP) - a 27 amino acid peptide and vasoactive intestinal peptide (VIP)- 28 amino acid. Both have similar biochemical properties and bind to VPAC1 receptors expressed in high density on the surface of certain cancers such as of the breast, prostate and urinary bladder (100%), colon (96%), pancreas (65%), lung (58%), stomach (54%), and liver (49%).²¹⁻²⁴

The present study was aimed to investigate the influence of these two BFC agents on the PACAP analogue, which differs in terms of their radiolabeling kinetics, biological activity and stability of the resultant product.

Methods and materials

Reagents

Fmoc amino acid, solvents and reagents for peptide chelator synthesis, was purchased from Fluka Chemicals (St. Louis, MO). The BFC DOTA was purchased from Macrocyclics (Dallas, TX) and NODAGA was purchased from Chematech (Dijon, France). Sodium acetate, diethylenetriamine pentaacetic acid (DTPA), calcium chloride (CaCl₂), zinc chloride (ZnCl₂), Ferric chloride (FeCl₃), hydrochloric acid solutions were purchased from Fisher Scientific, Inc. (Waltham, MA). Transferrin was purchased from Sigma-Aldrich (St. Louis, MO). A 370MBq [⁶⁸Ge-⁶⁸Ga] generator was purchased from Eckert and Zeigler (Berlin, Germany). Sodium chloride solution (0.9%) was prepared in this laboratory using deionized water. Human serum was prepared from whole blood drawn from a volunteer. All chemicals were used without further purification.

Instruments

Peptide was purified using a preparative column on high-pressure liquid chromatography (HPLC) system (Shimadzu Corporation, Kyoto, Japan) equipped with gradient pumps and UV/VIS detector, a NaI(Tl) radioactivity monitor, and a rate meter. The reverse phase C18 microbound column (4.6 mm × 250 mm) served as the stationary phase, and two solvents: 0.1% Trifluoroacetic acid (TFA) in H₂O and 0.1% TFA in acetonitrile as mobile phase. The gradient was such that 10% CH₃CN in aqueous 0.1% CF₃COOH to 100% CH₃CN in 0.1% CF₃COOH at a flow rate of 1 mL/min over 28 min at 22°C. For thin layer chromatography (TLC), pre-coated aluminum sheets with silica gel were purchased from Merck (Kenilworth, NJ) and radioactivity was measured on Perkin Elmer 2480 wizard² automatic gamma counter (Waltham, MA). The pH of the solutions was measured using either a pH meter (Sartorius, Gottingen, Germany) or pH strips (Fisher scientific Inc.)

Synthesis of peptide conjugate

Briefly, the PACAP analogue with C-terminal NODAGA/DOTA chelators was synthesized on a Wang resin using an ABI 341A peptide synthesizer (Applied Biosystems, Foster City, CA). Fmoc-Lys (ivDde) was first introduced at the C terminus of the peptide, followed by 4-aminobutyric acid (γ -Aba). The 27-amino-acid-long PACAP sequence was then assembled by standard Fmoc coupling with the final histidyl residue, being a t-Boc-protected His(Trt) derivative. The capping t-Boc function was necessary to ensure that the N-terminal amino group remained protected during

subsequent deprotection and coupling cycles performed at the ϵ amino group of the C-terminal lysine. The ivDde group at the C-terminal lysine was then selectively removed with 2% hydrazine, followed by coupling with DOTA or NODAGA. The peptide was cleaved from the resin using TFA/water/phenol/thioanisole/ethanedithiol (82.5:5:5:5:2.5) and precipitated with diethyl ether. The crude peptide was purified to homogeneity by reversed phase HPLC using a preparative column. The peptide was characterized by matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry.

Radiolabeling

For radiolabeling, 300 μ L (14.5 – 18.5 MBq) of $^{68}\text{GaCl}_3$ was added in a clean glass test tube to which were added varying amounts of acetate buffer to study the influence of pH (1.0 – 6.0) and varying amount of a 4 mg/mL peptide-chelators water solution (1-20 μ g). The mixture was heated at predetermined temperatures (25, 50, 70 and 90 $^\circ\text{C}$) for pre chosen time periods (10, 15, 20 and 30 min) to study the kinetics of the radiolabeling reactions with respect to temperature and time. The radiolabeling was evaluated by Radio-HPLC.

Preparation of free $^{68}\text{GaCl}_3$, ^{68}Ga -hydrolyzed, ^{68}Ga -DTPA and ^{68}Ga -Tf as reference standard

To identify HPLC peaks of the ^{68}Ga species, HPLC peaks, that may have formed during the chelation experiments, free $^{68}\text{GaCl}_3$ (as eluted from generator with 0.1M HCl), ^{68}Ga -hydrolyzed products, ^{68}Ga -DTPA and ^{68}Ga -Transferrin (Tf) were prepared as described^{25-26,13} and their retention times were recorded on radio-HPLC.

Stability (in vitro), transchelation and transmetallation studies

For the stability, transchelation and transmetallation studies, ^{68}Ga -peptide-chelators were prepared by mixing 300 μ L (14.5 – 18.5 MBq) of $^{68}\text{GaCl}_3$ in 0.1 M HCl, 120 μ L of 1.0 M sodium acetate and 5 μ L (20 μ g) of peptide-chelators. The preparation was incubated at 90 $^\circ\text{C}$ for 30 min. Aliquots of ^{68}Ga -peptide-chelators solutions (0.3 mL, approx. 15 MBq) were alternatively mixed with (1) 0.3 mL of a 0.9% NaCl, (2) 0.3 mL of a 100 nM $\text{Fe}^{3+}/\text{Zn}^{2+}/\text{Ca}^{2+}$ (for transmetallation study), (3) 0.3 mL of a 200 μ M DTPA solution, (4) 0.3 mL of a 50 μ M transferrin solution (for transchelation study) and (5) 0.3 mL of human serum solution. The mixtures were incubated at 37 $^\circ\text{C}$ and analyzed after 0, 10, 30, 60 and 120 min of incubation, respectively. A 10 μ l (in 100 μ L of normal saline) aliquot preparations of (1), (2) and (3) were injected in the radio-HPLC while preparations (4, 5) were analyzed by ITLC-SG using 0.1M citric acid as a solvent.

Cell binding (in vitro) assay

Estrogen receptor-positive T47D human breast tumor cells from American Type Culture Collection were maintained in RPMI 1640 medium with 2 mM L-glutamine contained 10 mM HEPES, 1.0 mM sodium pyruvate, 1.5 g/L sodium bicarbonate, 4.5 g/L glucose and supplemented with 0.2 Units/ml 90 %bovine insulin, and 10% fetal bovine serum, at 37°C under 5% CO₂/95% air. The cells when confluent were detached using 0.25% trypsin-EDTA, washed and resuspended with RPMI-1640 medium at a concentration of ~ 5 × 10⁶ cells/mL. Each test tube contained 1 mL of cell suspension into which 10 μL of ⁶⁸Ga-peptide-chelator (approx. 0.74 MBq) was added and incubated for 2 hr at 37°C in a water bath. The incubation was terminated by adding 0.5 ml of cold normal saline. The mixture was centrifuged at 3000 rpm for 10 minutes. The supernatant from each wash was collected into marked test tubes. To the remaining pellet, 1.0 ml of normal saline was added and centrifuged again at 3000 rpm for 10 minutes. The supernatant was separated and collected in the marked test tube. The cells and supernatant were counted for radioactivity in a gamma counter and percentage of radioactivity bound on the cells was calculated.

Biodistribution studies

The biodistribution of ⁶⁸Ga labeled peptide was studied in normal (n = 10, athymic nude, female mice, 20-25 g). The resultant optimal preparation procedure was used which contained 300 μL of ⁶⁸GaCl₃ in 0.1 N HCl, 120 μL of 1.0 M sodium acetate solution and 20 μg of peptide conjugate, incubated for 10 min at RT (for NODAGA-peptide) and 90°C (for DOTA-peptide). Following determination of radiochemical purity (> 95%) by radio-HPLC analysis, ⁶⁸Ga-peptide (3.7 – 5.5 MBq) was administered intravenously through a lateral tail vein in 200 μL of the preparation (pH ≈ 5.5). Radioactivity in the syringe before and after administration was measured in an energy-calibrated dose calibrator (CRC-15; Capintec, Ramssey, NJ), and the exact quantity received by each animal was determined. A standard ⁶⁸GaCl₃ solution of a known quantity was prepared at the time of injection. The animals (n = 5, each study) were sacrificed at 1 h post injection by CO₂ inhalation. Various organs were removed, washed free of blood, blotted dry, and weighed. The associated radioactivity was counted in the γ-counter and results were calculated as percentage injected dose per gram (% ID/g) of tissue.

Results

Purity of peptide-chelators conjugates

NODAGA- and DOTA-peptide purity as determined by HPLC was 99%. The MALDI-TOF demonstrated that, for NODAGA-peptide observed molecular mass was 3716 (calculated molecular mass-3718) whereas for DOTA-peptide observed molecular weight was 3787 (calculated molecular mass-3785). The MALDI peaks for the peptide conjugates are given in Figure 1A and B.

Radiolabeling kinetics

Optimal radiolabeling (> 95.0%) of the DOTA-conjugated peptide required pH 2-5 and higher (50-90°C) temperature, while the NODAGA conjugated peptide needed incubation only at room temperature at same pH range. At higher temperature (90°C), the radiolabeling was not affected by pH for both peptide conjugates (Fig. 2A, B) and can be labeled at wide range of pH (1.0 – 6.0). NODAGA-peptide can bind ⁶⁸Ga at a minimal quantity of 1 µg but DOTA-peptide required minimum of 20 µg to yield optimal (≥ 95%) radiolabeling (Fig. 2C). The minimum incubation time needed for optimal (> 95%) radiolabeling was 10 minutes at 90°C (Fig. 2D). Radio-HPLC showed retention time for both radio-complexes was 9.9 ± 0.3 minutes as compared to 3.1 ± 0.2 minutes for free ⁶⁸Ga as shown in Fig. 3-b. The retention time was also determined for the reference standard like hydrolyzed ⁶⁸GaCl₃, ⁶⁸Ga-DTPA and ⁶⁸Ga-transferrin and found to be 3.5 ± 0.2 minutes.

Stability studies

Both ⁶⁸Ga labeled peptide conjugates were stable in 0.9% normal saline (Fig. 4A) and showed minimal (1-2%) degradation against (FeCl₃, CaCl₂ and ZnCl₂) transmetallation (Fig. 4B-D) up to 2 hours. The transchelation study with DTPA (Fig. 5A) and transferrin (Fig. 5B) solution showed that both radio-complexes were remained intact over 2 hours. Radio-TLC showed ratio factor (Rf) value of 0.0-0.1 for ⁶⁸Ga labeled peptide and 0.9-1.0 for free ⁶⁸GaCl₃.

Cell binding (in vitro) assay

Cell binding assay showed higher uptake for ⁶⁸Ga-NODAGA-peptide, $34.0\% \pm 0.8\%$ as compared with $24.5\% \pm 0.9\%$ for ⁶⁸Ga-DOTA-peptide at 15 minutes. However, at 2

hours of incubation at 37°C, the uptake was nearly same, 48.9% ± 3.9% for ⁶⁸Ga-NODAGA-peptide and 45.2% ± 2.1% for ⁶⁸Ga-DOTA-peptide (Table – I).

Biodistribution studies

The biodistribution pattern of the both ⁶⁸Ga labeled peptide conjugate is shown in Figure 6. In both cases, a higher uptake is seen in kidneys followed by liver, blood and lungs. It was evident that the ⁶⁸Ga-NODAGA-Peptide showed rapid blood clearance as compared to ⁶⁸Ga-DOTA-peptide. Liver uptake was 8.8% for ⁶⁸Ga-NODAGA-Peptide was lower (p = 0.01) as compared to 16.5% for ⁶⁸Ga-DOTA-peptide. Most of the activity was excreted through kidney.

Discussion

With its several advantages such as the availability in a generator form, cyclotron independency, multi elution ability in a single day, simplified radiolabeling chemistry with several biomolecules using bifunctional chelating agents, the use of ⁶⁸GaCl₃ is rapidly growing for diversified clinical applications.²⁷ The presence of even µg levels of metallic impurities (with ageing of generator) in ⁶⁸GaCl₃ or reagent used may hamper the radiolabeling complexation. The removal of radionuclide impurities however has helped expansion of its use for clinical application.²⁸

The eluate of ⁶⁸GaCl₃ from the generator may contain metal ion impurities such as Fe³⁺, Cu²⁺, Al³⁺, Zn²⁺ and Sn⁴⁺ which could hamper the radiolabeling of biomolecules. It is important therefore to identify the suitable chelating agent for the radiolabeling which is least affected by these impurities as well the stability of the labeled compound.²⁹⁻³⁰ DOTA is the most commonly used chelating agent and ⁶⁸Ga-DOTATOC is accepted in the European Pharmacopeia.³¹ ⁶⁸Ga labeled DOTA biomolecules conjugates (like peptides, antibody, and small molecules) have been successfully developed and used for many clinical applications. The disadvantage of DOTA conjugate is its slow reaction kinetics for complex formation with ⁶⁸Ga.³² These limitations prompted us to investigate the use of NODAGA for comparison with DOTA.

Chelator-peptide conjugates were synthesized by solid phase peptide synthesis method. Both peptide conjugates were synthesized with high purity (~ 99.0%). NODAGA-peptide conjugate showed faster kinetics as compare to DOTA- peptide conjugate and less influenced by the pH variation. NODAGA peptide conjugate yielded higher radiolabeling efficiency at minimal quantity of peptide (1 µg), at room temperature and within short incubation (5 min) time. For the preparations with short lived radionuclides like ⁶⁸Ga, these characteristics of NODAGA are advantageous. The faster radiolabeling

kinetics may be attributable to its ionic size compatibility with cavity size of the nine membered triazomacrocyclic of NODAGA and therefore, yielding a high radiolabeling in 5 min at room temperature.³³ A low radiolabeling was observed in acidic (pH-1.0) conditions for both peptide conjugates when incubated at temperature (25-70°C). It may attribute to strongly acidic environment which might have protonated the donor atoms of the ligand and destabilized the ⁶⁸Ga complex.³⁴ At high temperature (~ 90°C), the radiolabeling was found to be independent of pH variation (Fig. 2 a-b). However at pH 7.0 and above, labeling decreases (data not included). This may be due to a fact that at acidic pH, ⁶⁸Ga remains predominant in cation Ga³⁺ form (> 90%), though soluble complex ions Ga(OH)²⁺, Ga(OH)⁺² may be present in the solution. As the pH of the solution shifts towards basic (> 7.0), the hydrolyzed form of Gallium, Ga(OH)⁻⁴ increases and decreases the radiolabeling yield.³⁵ Once the peptide conjugate is radiolabeled, then adjusting the pH to physiological pH 7.0 does not adversely influence the chelated Ga-68.

The use of sodium acetate buffer maintained the pH and allowed strong formation BFC-⁶⁸Ga complex. The other advantage of using sodium acetate buffer is its biocompatibility for human use.³⁶ DOTA peptide conjugate showed labeling at high temperature. Therefore, DOTA may not be useful for those peptides which can denature at higher temperature and become unsuitable for molecular imaging. On the other hand, radiolabeling of NODAGA-peptide at room temperature ruled out major thermal denaturation. Radio-HPLC showed that the formation of the ⁶⁸Ga-NODAGA and DOTA peptide conjugate has retention time of around 9.9 min whereas free ⁶⁸Ga eluted at 3.6 min (Fig. 3-b) and complex formation between ⁶⁸Ga and NODAGA-peptide conjugate is shown in Fig. 3-a . HPLC UV detector also detected the UV peak for NODAGA and DOTA peptide conjugate at around 9.9 min. Therefore, it confirmed the complexation of the ⁶⁸Ga with peptide conjugate which leads to the retention of the ⁶⁸Ga labeled peptide approximately at the same time.

Both radio-complexes were found to be stable in normal saline (Fig. 4a) and human serum over 120 minute. The stability in human serum (at 37°C) indicated its suitability for *in vivo* applications and showed their inertness towards the metal ion/species that may be present in the serum. The transmetallation study was important not because these ions are present as impurities in the eluate but these ions are also present in serum in appreciable quantity.³⁷ ⁶⁸Ga-NODAGA and DOTA peptide conjugates showed minimal (1-2%) degradation in radiolabeling after 120 minute incubation with respective ion solution at 37°C (Fig. 4-b-d). This could also provide an indication about the stability of these radio-complexes in *in vivo* environment. Transferrin is the major protein in the plasma which is known to form a strong complex with ⁶⁷Ga-citrate.³⁸ The minimal transchelation (~1-2%) of ⁶⁸Ga-peptide conjugates with transferrin showed the stability of radio-complexation (Fig. 5-a-b).

The cell binding assay was performed on T47D breast cancer lines which express VPAC-1 receptors on their surface.¹⁹ ⁶⁸Ga-NODAGA-peptide showed higher uptake in 15 minute as compared to ⁶⁸Ga-DOTA-peptide but the uptake was almost same at 2hr (Table-1) which may be due to saturation of the receptor site. The biodistribution data (Fig. 6) showed that both radio-complexes were cleared through kidneys. Hence, renal system was the major route of clearance from the body. ⁶⁸Ga-NODAGA-peptide was cleared rapidly from the blood and had significantly less liver uptake ($p < 0.05$) as compare to ⁶⁸Ga-DOTA-peptide at 1 hr. The fast clearance from the blood may be advantageous in reducing the background radioactivity for PET imaging but it could also limit the accumulation of radioactivity at target (tumor) area. Hence, a detailed *in vivo* investigation will be required further for any conclusion.

In the present study, we have showed that ⁶⁸Ga-NODAGA-peptide exhibited most favorable features from radiochemistry point of view. Chelator NODAGA has flexibility over the influence of temperature and acidic environment.¹³ Labeling at room temperature provides an advantage and could prevent any deleterious effect on peptide conjugate due to heating. Both peptide conjugates were found to be stable in serum, against transchelation and transmetallation experimentations. A detailed *in vivo* study is in progress to investigate the influence of chelators on the biodistribution and tumor targeting.

Conclusions

A comparison between NODAGA and DOTA revealed that NODAGA-peptide conjugate could be efficiently radiolabeled with ⁶⁸Ga at room temperature with high yield. ⁶⁸Ga-NODAGA-peptide conjugate remained stable in serum, in presence of major metal ions impurities, against transchelation with DTPA or transferrin. The biodistribution data showed its clearance from kidneys, which minimize the radiation burden on the organs. The clearance of ⁶⁸Ga-NODAGA-peptide from blood was faster as compare to ⁶⁸Ga-DOTA-peptide and would be advantageous for PET imaging.

Acknowledgments

The research, in part, was supported by NIH/NCI RO1 CA 157372 (M.L.T.), NIH/NCI 1S10OD012406 (M.L.T.), and NIH/NCI S10RR23709 (M.L.T.).

Disclosure Statement

M.L.T. is consultant to NuView and Zevacor, Inc. No other potential conflicts of interest relevant to this article are reported.

References

1. Schrevens L, Lorent N, Doms C, et al. The role of PET scan in diagnosis, staging, and management of non-small cell lung cancer. *Oncologist* 2004; 9: 633.
2. Rahmim A and Zaidi H. PET versus SPECT: strengths, limitations and challenges. *Nuclear Medicine Communications* 2008; 29: 193.
3. Griffeth LK. Use of PET/CT scanning in cancer patients: technical and practical considerations. *BUMC proceedings* 2005; 18: 321.
4. Bading JR, Alauddin MM, Fissekis JD, et al. Blocking catabolism with eniluracil enhances PET studies of 5-18F-fluorouracil pharmacokinetics. *J Nucl Med* 2000; 41: 1714.
5. Sun H, Sloan A, Mangner TJ, et al. Imaging DNA synthesis in vivo with 18F-FMAU and PET. *J Nucl Med* 2005; 46: 292.
6. Ido T, Wan CN, Casella V, et al. Labeled 2-deoxy-D-glucose analogs. 18F-labeled 2-deoxy-2-fluoro-D-glucose, 2-deoxy-2-fluoro-D-mannose and C-14-2-deoxy-2-fluoro-D-glucose. *J Labelled Comp Radiopharm* 1978; 14: 175.
7. Choi SJ, Kim JS, Kim JH, et al. 18F-3'-Deoxy-3'-fluorothymidine PET for the diagnosis and grading of brain tumors. *Eur J Nucl Med Mol Imaging* 2005; 32: 653.
8. Asti M, De Pietri G, Fraternali A, et al. Validation of (68)Ge/(68)Ga generator processing by chemical purification for routine clinical application of (68)Ga-DOTATOC. *Nuclear medicine and biology* 2008; 35:721.
9. Decristoforo C, Knopp R, von Guggenberg E, et al. A fully automated synthesis for the preparation of 68Ga-labelled peptides. *Nuclear medicine communications* 2007; 28: 870.
10. Fani M, Andre JP, Maecke HR. 68Ga-PET: a powerful generator-based alternative to cyclotron- based PET radiopharmaceuticals. *Contrast media & molecular imaging* 2008; 3: 67.
11. Strand J, Honarvar H, Perols A, et al. Influence of macrocyclic chelators on the targeting properties of 68Ga-labeled synthetic affibody molecules: comparison with 111In-labeled counterparts. *PLoS One* 2013; 8(8): e70028.
12. Chakravarty R, Chakraborty S, Dash A, et al. Detailed evaluation on the effect of metal ion impurities on complexation of generator eluted 68Ga with different bifunctional chelators. *Nuclear Medicine and Biology* 2013; 40: 197.

13. Asti M, Iori M, Capponi PC, et al. Influence of different chelators on the radiochemical properties of a 68-Gallium labelled bombesin analogue. *Nucl Med Biol* 2014; 41(1): 24.
14. Reubi JC. Peptide receptors as molecular targets for cancer diagnosis and therapy. *Endocr Rev* 2003; 24(4): 389.
15. Weiner RE, Thakur ML. Radiolabeled peptides in oncology: role in diagnosis and treatment. *BioDrugs* 2005; 19(3): 145.
16. Thakur ML, Marcus CS, Saeed S, et al. Imaging tumors in humans with Tc-99m-VIP. *Ann N Y AcadSci* 2000; 921: 37.
17. Thakur ML, Zhang K, Berger A, et al. VPAC1 receptors for imaging breast cancer: a feasibility study. *J Nucl Med* 2013; 54(7): 1019.
18. Zhang K, Aruva MR, Shanthly N, et al. PET imaging of VPAC1 expression in experimental and spontaneous prostate cancer. *J Nucl Med* 2008; 49(1): 112.
19. Zhang K, Aruva MR, Shanthly N, et al. Vasoactive intestinal peptide (VIP) and pituitary adenylate cyclase activating peptide (PACAP) receptor specific peptide analogues for PET imaging of breast cancer: In vitro/in vivo evaluation. *RegulPept* 2007; 144(1-3): 91.
20. Thakur ML, Aruva MR, Garipey J, et al. PET imaging of oncogene overexpression using 64Cu-vasoactive intestinal peptide (VIP) analog: comparison with 99mTc-VIP analog. *J Nucl Med* 2004; 45(8): 1381.
21. Schulz S, Rocken C, Mawrin C, et al. Immunocytochemical identification of VPAC1, VPAC2 and PAC1 receptors in normal and neoplastic human tissues with subtype specific antibodies. *Clin Cancer Res* 2004; 10: 8234.
22. Reubi JC. Neuropeptide receptors in health and disease: The molecular basis for in vivo imaging. *J Nucl Med* 1995; 36: 1825.
23. Reubi JC. Regulatory peptide receptors as molecular targets for cancer diagnosis and therapy. *Q J Nucl Med* 1997; 41: 63.
24. Reubi JC, Läderach U, Waser B, et al. Vasoactive intestinal peptide/pituitary adenylate cyclase activating peptide receptor subtypes in human tumors and their tissues of origin. *Cancer Res* 2000; 60: 3105.
25. Asti M, Iori M, Erba PA, et al. Radiosynthesis of 68Ga-labelled DOTA-biocytin (68Ga-r-BHD) and assessment of its pharmaceutical quality for clinical use. *Nucl Med Commun* 2012; 33(11): 1179.

26. Kumar V, Boddeti DK, Evans SG, et al. Potential use of ^{68}Ga -apo-transferrin as a PET imaging agent for detecting *Staphylococcus aureus* infection. *Nucl Med Biol* 2011; 38: 393.
27. Schultz MK, Mueller D, Baum RP, et al. A new automated NaCl based robust method for routine production of gallium-68 labeled peptides. *Appl Radiat Isot* 2013; 76: 46.
28. Vis R, Lavalaye J, van de Garde EM. GMP-compliant (^{68}Ga) radiolabelling in a conventional small-scale radiopharmacy: a feasible approach for routine clinical use. *Eur J Nucl Med Mol Imaging* 2015; 5: 27.
29. Zhernosekov KP, Filosofov DV, Baum RP, et al. Processing of generator-produced ^{68}Ga for medical application. *J Nucl Med* 2007; 48: 1741.
30. Asti M, Pietri GD, Fraternali A, et al. Validation of $^{68}\text{Ge}/^{68}\text{Ga}$ generator processing by chemical purification for routine clinical application of ^{68}Ga -DOTATOC. *Nucl Med Biol* 2008; 35: 721.
31. European Pharmacopeia 7.7 (01/2013:2482 Gallium (^{68}Ga) Edotreotide injection) 2011; 23: 310.
32. Maecke HR, Hofmann M, Haberkorn U. ^{68}Ga -labeled peptides in tumor imaging. *J Nucl Med* 2005; 46: 172S.
33. Ferreira CL, Lamsa E, Woods M, et al. Evaluation of bifunctional chelates for the development of gallium-based radiopharmaceuticals. *Bioconjug Chem* 2010; 21: 531.
34. Velikyan I. Positron emitting ^{68}Ga -based imaging agents: chemistry and diversity. *Med Chem* 2011; 7: 338.
35. Ali M, Hsieh W, Tsopeles C. An improved assay for (^{68}Ga)-hydroxide in (^{68}Ga)-DOTATATE formulations intended for neuroendocrine tumour imaging. *J Labelled Comp Radiopharm* 2015; 58: 383.
36. Bauwens M, Chekol R, Vanbilloen H, et al. Optimal buffer choice of the radiosynthesis of ^{68}Ga -Dotatoc for clinical application. *Nuclear Medicine Communications* 2010; 31: 753.
37. Riss PJ, Burchardt C, Roesch F. A methodical ^{68}Ga -labelling study of DO2A-(butyl-L-tyrosine)₂ with cation exchange post processed ^{68}Ga : practical aspects of radiolabeling. *Contrast Media Mol Imaging* 2011; 6: 492.
38. Harris WR, Pecoraro VL. Thermodynamic binding constants for gallium transferrin. *Biochemistry* 1983; 22: 292.