

10-1-2018

A systematic review of treating recurrent head and neck cancer: a reintroduction of brachytherapy with or without surgery.


Julianna Rodin
Thomas Jefferson University

Voichita Bar-Ad
Thomas Jefferson University

David Cagnetti
Thomas Jefferson University

Joseph Curry
Thomas Jefferson University

Jennifer Johnson
Follow this and additional works at: <https://jdc.jefferson.edu/otofp>
Case Western Reserve University

 Part of the [Oncology Commons](#), and the [Otolaryngology Commons](#)

[Let us know how access to this document benefits you](#)

See next page for additional authors

Recommended Citation

Rodin, Julianna; Bar-Ad, Voichita; Cagnetti, David; Curry, Joseph; Johnson, Jennifer; Zender, Chad; Doyle, Laura; Kutler, David; Leiby, Benjamin E.; Keane, William M.; and Luginbuhl, Adam J., "A systematic review of treating recurrent head and neck cancer: a reintroduction of brachytherapy with or without surgery." (2018). *Department of Otolaryngology - Head and Neck Surgery Faculty Papers*. Paper 39.
<https://jdc.jefferson.edu/otofp/39>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Otolaryngology - Head and Neck Surgery Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Julianna Rodin, Voichita Bar-Ad, David Cognetti, Joseph Curry, Jennifer Johnson, Chad Zender, Laura Doyle, David Kutler, Benjamin E. Leiby, William M. Keane, and Adam J. Luginbuhl

A systematic review of treating recurrent head and neck cancer: a reintroduction of brachytherapy with or without surgery

Julianna Rodin, MD¹, Voichita Bar-Ad, MD², David Cagnetti, MD¹, Joseph Curry, MD¹, Jennifer Johnson, MD, PhD³, Chad Zender, MD⁴, Laura Doyle, MS², David Kutler, MD⁵, Benjamin Leiby, PhD⁶, William Keane, MD¹, Adam Luginbuhl, MD¹

¹Department of Otolaryngology, Thomas Jefferson University, Philadelphia, PA, ²Department of Radiation Oncology, Thomas Jefferson University, Philadelphia, PA, ³Department of Otolaryngology, Case Western Reserve University School of Medicine, Cleveland, OH,

⁴Department of Medical Oncology, Thomas Jefferson University, Philadelphia, PA, ⁵Department of Otolaryngology, Weill Cornell Medical Center, New York, NY, ⁶Department of Biostatistics, Thomas Jefferson University, Philadelphia, PA, USA

Abstract

Purpose: To review brachytherapy use in recurrent head and neck carcinoma (RHNC) with focus on its efficacy and complication rates.

Material and methods: A literature search of PubMed, Ovid, Google Scholar, and Scopus was conducted from 1990 to 2017. Publications describing treatment of RHNC with brachytherapy with or without surgery were included. The focus of this review is on oncologic outcomes and the safety of brachytherapy in the recurrent setting.

Results: Thirty studies involving RHNC treatment with brachytherapy were reviewed. Brachytherapy as adjunctive treatment to surgical resection appears to be associated with an improved local regional control and overall survival, when compared with the published rates for re-irradiation utilizing external beam radiotherapy (RT) or brachytherapy alone. Safety data remains variable with different isotopes and dose rates with implantable brachytherapy demonstrating a tolerable side effect profile.

Conclusions: Although surgery remains a mainstay treatment for RHNC, intraoperative interstitial brachytherapy delivery as adjunctive therapy may improve the treatment outcome and may be associated with fewer complication rates as compared to reirradiation using external beam radiotherapy. Further investigations are required to elucidate the role of brachytherapy for RHNC.

J Contemp Brachytherapy 2018; 10, 5: 454–462

DOI: <https://doi.org/10.5114/jcb.2018.79399>

Key words: cesium, head and neck cancer, recurrent.

Purpose

Head and neck carcinoma (HNC) continue to be a commonly diagnosed cancer with almost 50,000 new cases and almost 10,000 deaths each year [1]. More than 90% of HNC are classified as squamous cell carcinoma, and cases have increased by nearly 5,000 new patients over two years [2,3]. Patients with locally advanced disease generally are treated with an algorithm of surgical resection, radiotherapy (RT), and/or chemoradiotherapy with curative intent. Unfortunately, depending on subsite and stage, almost half of these advanced stage patients fail this approach in the first 5 years [4,5]. The predominant pattern of recurrence is in a locoregional manner [5]. Surgical resection for these patients has become the mainstay of treatment in resectable cases, and re-irradiation as the only locoregional treatment modality for unresectable cases.

Brachytherapy with or without salvage surgery has often been used for this group of cancer patients with RHNC. Compared to external beam radiation therapy (EBRT) delivery systems, including intensity modulated radiation therapy (IMRT) and proton beam therapy, brachytherapy offers focused dose delivery and fewer early toxic effects, while minimizing adjacent non-cancer tissue toxicity, specifically in previously irradiated fields [5]. This protection of adjacent normal tissue can result in preservation of laryngeal and tongue function. Although brachytherapy has been utilized for recurrent head and neck cancer (RHNC) for years, its use has been limited due to availability of technical expertise, variable isotope implantation technical issues, and introduction of IMRT and other alternative EBRT modalities. Brachytherapy can be delivered via permanent implants (using low-dose-rate brachytherapy or LDR or high-dose-rate brachyther-

Address for correspondence: Adam Luginbuhl, MD, Department of Otolaryngology, Thomas Jefferson University, 925 Chestnut Street, 6th floor, Philadelphia, PA 19107 United States, phone: +1 215-503-6828, e-mail: adam.luginbuhl@jefferson.edu

Received: 30.05.2017

Accepted: 21.10.2018

Published: 30.10.2018

apy or HDR) or via removable catheters through which typically high activity radioisotopes are inserted (using HDR or pulsed-dose-rate brachytherapy or PDR) [6]. Permanent implants may be advantageous in patients with recurrence in irregular surfaces that are not suitable for catheter-based isotope delivery [7]. This review describes brachytherapy efficacy in recurrent primary HNC and recurrence in neck disease, and potential complications of brachytherapy. We separated studies based on whether brachytherapy was the sole treatment or in conjunction with surgical tumor resection.

Material and methods

A literature search using PubMed, Ovid, Google Scholar, and Scopus was conducted. Databases were searched for the terms 'recurrent head and neck cancer' AND 'brachytherapy' between 1990 and 2017. Additional search terms included 'Cesium-131', 'Iridium-192', and 'Iodine-125'. The general criteria for the studies to be included were publications describing brachytherapy use in recurrent HNC of various sites, with particular focus on treatment outcomes and complications. Publications involving ^{131}Cs brachytherapy are also described. The search identified a total of 380 papers, with 350 papers being excluded after title and abstract screening. In these excluded papers, brachytherapy in recurrent HNC

of various sites was not the chief focus. The remaining 30 studies reported adequate information according to the criteria. Statistical analysis was also performed to compare recurrence rates in studies, which used brachytherapy with surgery versus brachytherapy alone.

Results

In the 30 reviewed studies, 95% (1,440 of 1,515 patients) had prior radiation therapy, making radiotherapy treatment options limited. In 23 of these studies, brachytherapy was administered using Iridium-192 (^{192}Ir) in 1,003 patients, Iodine-125 (^{125}I) in 131 patients, and Cesium-131 (^{131}Cs) in 35 patients. Seven studies did not explicitly state which type of isotope was used [8,9,10,11,12,13,14]. Brachytherapy was incorporated into treatment planning with or without salvage surgery in a combined modality approach. Occasionally, brachytherapy was also followed by further radiation therapy or chemotherapy. Reirradiation with brachytherapy was used for local, locoregional, or nodal recurrence or neck disease and not for distant metastatic spread. As with most forms of reirradiation, the toxicity profile of brachytherapy includes neurologic toxicity, dysphagia, carotid artery rupture, skin necrosis, fistulas, and osteoradionecrosis [15,16]. These aspects and other considerations within the studies are described in further detail below.

Table 1. Overview of studies treating recurrent head and neck squamous cell carcinoma using brachytherapy alone

Author, Year	N	Brachytherapy	Total dose	Additional therapy	Results	Complications
Grimard, 2006 [5]	82	Ir 192; LDR	≥ 55 Gy	Surgery (14), EBRT (3), Surgery + EBRT (6)	LC at 2 years: 37% OS at 2 years: 33%	8.5% acute toxicity 22% late toxicity
Strnad, 2014 [8]	104	PDR	55 Gy or 24 Gy (when +EBRT)	Surgery (51%), EBRT (32%), Chemo (56%)	LC at 5 years: 82% OS at 5 years: 21%	31% late toxicity
Strnad, 2014 [12]	51	PDR	60 Gy or 27 Gy (when +EBRT)	EBRT (22%), Chemo (69%)	LC at 5 years: 57%	10% acute grade 3 29% late toxicity
Puthawala, 2001 [17]	220	Ir 192; LDR	Median: 53 Gy	Chemo (88)	LC at 5 years: 51% OS at 5 years: 20%	60% acute toxicity 27% late toxicity (65% of which had RT > 65 Gy)
Hepel, 2005 [18]	30	Ir 192; HDR	Mean: 34 Gy	Chemo (43%), EBRT (6%)	LC at 2 years: 67% OS at 2 years: 37%	16% late grade 3-4 toxicity
Jiang, 2011 [19]	29	I 125; LDR, US-guided	Median: D ₉₀ 130 Gy	Not mentioned	LC at 2 years: 35% OS at 2 years: 28%	No carotid rupture or soft tissue/bone necrosis
Glatzel, 2002 [37]	90	Ir 192; HDR	Median: 17.5 Gy	Not mentioned	Median: OS 6 Mo	7% late grade 3-4
Zhu, 2013 [32]	19	I 125; LDR, US or CT-guided	Median: D ₉₀ 131 Gy	Not mentioned	LC at 2 years: 27.5% OS at 2 years: 18%	No grade 4 or 5 toxicity
Meng, 2012 [38]	17	I 125; LDR, US or CT-guided	Median: D ₉₀ 126 Gy	Not mentioned	LC at 2 years: 50% OS at 2 years: 24%	No grade 3 or 4 toxicity
Wiegand, 2013 [39]	12	Ir 192; HDR	Mean: 20-33 Gy	Not mentioned	Median: OS 8.5 Mo	No serious complications

Ir 192 – Iridium 192; I 125 – Iodine 125; Gy – Gray; LDR – low-dose-rate; HDR – high-dose-rate; PDR – pulsed-dose-rate; US – ultrasound; CT – computed tomography; EBRT – external beam radiation therapy; RT – radiotherapy; LC – local control; OS – overall survival

Brachytherapy for local disease: with and without salvage surgery

Re-irradiation with brachytherapy has been used as an option for resectable and unresectable RHNC cases. While salvage surgery in general is favored, its use may be limited by the extent of disease and the cancer’s intimate anatomic relationship to adjacent structures. Low-dose-rate (LDR), high-dose-rate (HDR), and pulsed-dose-rate (PDR) brachytherapy can be used as monotherapy alone or adjuvantly with salvage surgery. Tables 1 and 2 provide details of all studies using brachytherapy alone and with surgery, respectively. Some of these studies are further described below.

The use of brachytherapy alone was associated with lower local control (LC) and overall survival (OS) rates. In a study by Puthawala *et al.* [17], 220 patients with RHNC received salvage brachytherapy using LDR interstitial Iridium-192 (¹⁹²Ir) implants. They achieved 2-, 5-, and 10-year disease-free survival (DFS) rates of 60%, 33%, and 22%, respectively [17]. Hepel *et al.* [18] found similar results using HDR ¹⁹²Ir brachytherapy with DFS at 2 years

being 45% and OS at 1 and 2 years, 56% and 37%, respectively. Strnad *et al.* [8] found that their LC rates at 2, 5, and 10 years were 92.5%, 82.4%, and 58.9%, respectively, with a 5-year OS of 21% using PDR brachytherapy. Jiang *et al.* [19] used LDR ¹²⁵I ultrasound-guided permanent seeds in patients with RHNC ineligible for salvage surgery, and achieved LC rates at 1, 2, and 3 years of 53.1%, 34.8%, and 17.4%, and OS rates at 1, 2, and 3 years of 54.1%, 27.5%, and 27.5%, respectively.

In the recurrent setting, surgical resection remains the foundation of treatment for resectable disease, when looking at oncologic outcomes. In a study by Grimard *et al.* [5], brachytherapy was used as monotherapy in 22 of 45 patients and with surgery in 20 of 45 patients [5]. While not statistically significant, salvage surgery with adjuvant brachytherapy led to better locoregional control [5]. Narayana *et al.* [9] studied patients with recurrent HNC with 18/30 receiving surgery and HDR, 3/30 receiving EBRT and HDR, and 9/30 receiving HDR alone. There was improved local control in patients who received salvage surgery versus those without surgery: 88% vs. 40% [9]. In a study by Rudzianskas *et al.* [20], 43% of patients

Table 2. Overview of studies treating recurrent head and neck squamous cell carcinoma using surgery with brachytherapy

Author, year	N	Brachytherapy	Total dose	Surgical resection	Results	Complications
Henderson, 2016 [6]	23	Ir 192; HDR Cs-131; LDR	Mean: 21 Gy Mean: 81 Gy	Yes	Not mentioned	50% toxicity in HDR pts 30% toxicity in LDR pts
Narayana, 2007 [9]	30	HDR	34 Gy (+ surgery) 20 Gy (+ EBRT) 40 Gy (only BT)	60% Yes 40% No	LC at 2 years: 71% (88% + surgery, 40% – surgery) OS at 2 years: 63%	33% grade 2-3 toxicity
Martinez-Fernandez, 2017 [10]	63	HDR	Total: 32 or 40 Gy	Yes	LC at 15 years: 41% OS at 15 years: 12%	51% grade ≥ 3 toxicity
Schiefke, 2008 [11]	13	HDR	Median: 30 Gy	Yes	OS at 2 years: 65% LR failure in 38%	69% acute toxicity 31% late toxicity
Scala, 2013 [13]	76	HDR	Median rate: 1,200 cGy	Yes	LC at 2 years: 62% OS at 2 years: 42%	No grade 3-4 toxicity
Rudzianskas, 2012 [20]	30	Ir 192; HDR	Total: 30 Gy	43% Yes 57% No	LC at 2 years: 77% OS at 2 years: 62% <u>Non-surgical group:</u> LC at 2 years: 47% OS at 2 years: 35%	10% acute grade 2-3 10% late grade 2-4
Park, 1991 [21]	35	I 125; LDR	Mean rate: 8,280 cGy	Yes	OS at 2 years: 38% OS at 5 years: 29%	36% acute and/or late toxicity
Pellizzon, 2005 [22]	42	Ir 192; HDR	Median: 24 Gy	Yes	Crude LC 57% OS at 5 years: 52.5%	19% acute and/or late toxicity
Pham, 2015 [23]	18	Cs 131; LDR	Median: 80 Gy	Yes	OS at 18 mos: 45% LR failure in 33%	11% grade 3 toxicity
Teckie, 2013 [41]	57	Ir 192; HDR	Median: 15 Gy	Yes	OS at 3 years: 32% LR failure in 46%	37% grade 3 No grade 4-5 toxicity
Perry, 2010 [42]	34	Ir 192; HDR	Median: 15 Gy	Yes	OS at 2 years: 55%	29% grade 3-4 toxicity

Ir 192 – Iridium 192; I 125 – Iodine 125; Cs 131 – Cesium 131; Gy – Gray; LDR – low-dose-rate; HDR – high-dose-rate; EBRT – external beam radiation therapy; BT – brachytherapy; LC – local control; LR – locoregional; OS – overall survival

had surgery and HDR with ^{192}Ir , while 57% had HDR alone. Patients who received surgery and brachytherapy had improved 2-year local control rates (77%) and OS rates (62%) versus those with brachytherapy alone (47%, 35%, respectively) [20]. These differences in locoregional control underscore the overall benefit of surgical resection in the recurrent setting.

In a study by Park *et al.* [21], LDR ^{125}I seeds were implanted in patients with RHNC who all underwent salvage surgery. DFS at 1, 2, and 5 years was 64%, 47%, and 41%, respectively, with a 5-year no evidence of disease (NED) in patients (29%) [21]. Pellizzon *et al.* [22] studied patients with RHNC treated with surgery and HDR ^{192}Ir brachytherapy with or without further EBRT and found the OS at 5 and 8 years to be 52.5% and 48.1%. Martinez-Fernandez *et al.* [10] followed previously irradiated RHNC patients treated with salvage surgery and perioperative HDR brachytherapy for 15 years, finding a 15-year LRC rate of 41.2% and 15-year OS rate of 11.6%. Pham *et al.* [23] showed that in selected patients, ^{131}Cs intraoperative seeds with salvage surgery achieved 1-, 2-, and 3-year LC rates of 69%, 62%, and 52%, respectively.

Looking to historic data of re-irradiation using the standard or hyperfractionated IMRT combined with systemic chemotherapy has demonstrated a 2-year survival of 10-25% [24,25]. The previously discussed studies using surgery plus brachytherapy reported a 5-year overall survival of 33-67%, showing encouraging progress over EBRT plus chemotherapy.

We performed a statistical analysis of the studies to evaluate locoregional recurrence in RHNC patients receiving brachytherapy and surgery vs. brachytherapy alone. Logistic regression was used to compare recurrence rates, and generalized estimating equations (GEE) methods were used to account for clustering within the study. In studies where recurrence rates were not explicitly stated, recurrence rates were extrapolated based on the local and locoregional control rates. For the primary analysis, the rate of locoregional recurrence was 38.8% (143/369) in the surgery with brachytherapy group and 49.0% (329/671) in the brachytherapy alone group. The estimated odds ratio was 0.66 (95% CI: 0.38-1.12; $p = 0.13$).

Brachytherapy for nodal disease

Many studies also have demonstrated brachytherapy use in RHNC with regional failure in the cervical lymph nodes. Table 3 portrays the details of these studies. In the study by Bartochowska *et al.* [26], patients with unresectable isolated cervical lymph node recurrence had PDR or HDR with ^{192}Ir via catheters. Local control rates at 1 and 2 years were 31.7% and 19% [26]. Huang *et al.* [27] achieved better results in 31 patients with RHNC with neck metastases unmanageable with other techniques, achieving LC rates of 64.5% and 45.1% at 1 and 2 years. In 74 patients with inoperable recurrent lymphadenopathy, Tselis *et al.* [28] showed OS rates of 42%, 19%, and 6% at 1, 2, and 3 years, respectively.

In the study by Teudt *et al.* [14], patients with RHNC underwent revision surgery and perioperative HDR for their recurrent neck metastases. These patients were

found to have 78% and 67% OS at 2 and 5 years, respectively, with a median survival of 65 months [14]. Nutting *et al.* [29] also looked at patients with recurrent neck disease undergoing LDR brachytherapy alone, with resection, or with resection and flap surgery. Locoregional control at 2 years in those who had brachytherapy alone was 0%, while those who also had a surgery with or without a flap had improved LRC, 37% or 33%, respectively [29]. For the neck disease statistical analysis, the rate of locoregional recurrence was 30.9% (42/136) in the surgery with brachytherapy group and 49.0% (149/304) in the brachytherapy alone group. The estimated odds ratio was 0.46 (95% CI: 0.25-0.86; $p = 0.015$).

Complications

Due to the nature of the radioactive isotopes, brachytherapy is associated with various radiation exposure concerns. However, these radiation effects appear to be improved in terms of functional organ preservation and reduced irradiation to surrounding normal tissues compared to EBRT or IMRT (Quon). Toxicity is commonly monitored using the Radiation Toxicity Oncology Group (RTOG) common toxicity criteria [30]. Radiation safety procedures, including the use of lead shields and gloves, may be appropriate when handling specific radioisotope implants, which is based on a number of factors including isotope, total source activity, and implant location [31]. The recommended limit of radiation exposure to staff personnel based on federal regulation is 50 mSv/year for "stochastic effects", e.g., cancer or genetic effects, defined as "ones in which the probability of occurrence increases with increasing absorbed dose, but the severity does not depend on the magnitude of the absorbed dose" [31]. In some cases associated with specific radioisotopes, patients were isolated in order to reduce personnel and family radiation exposure, and were required to stay in the hospital longer.

Toxicity complications related to brachytherapy have been explored in various studies using ^{192}Ir and ^{125}I and usually involve necrosis, dysphagia, or fistulas. Tables 1-3 include toxicity rates for studies reviewed. In a review by Kasperts *et al.* [15], 17% (120/704) of patients developed soft tissue necrosis and 2% (15/704) developed orocutaneous fistulas. Another 2% of patients suffered from hemorrhage or carotid blowouts and another 3% developed osteoradionecrosis [15]. In the study by Grimard *et al.* [5], acute toxicity occurred in 15.6% of patients and late toxicity in 40%, with 4 patients developing grade 2 or 3 radionecrosis and 7 patients experiencing dysphagia. In Zhu *et al.*'s study [32], only one patient had a grade 1 skin reaction, with no patients suffering from soft tissue necrosis or other complications. Henderson *et al.* [6] looked at patients who underwent HDR or LDR brachytherapy and neck dissection with or without flap coverage. There were complications in 50% of HDR patients and 29.4% of LDR patients, with only 3 of 22 patients receiving flaps having complications (2 HDR, 1 LDR) [6]. Martinez-Fernandez *et al.* [10] found RTOG grade 3 or greater toxicities in 50.8% of patients, with 17 cases needing further operations due to fistula (6), wound dehiscence (4), bleeding (3),

Table 3. Overview of studies treating recurrent squamous cell carcinoma as neck disease

Author, year	N	Brachytherapy	Total dose	Surgical resection	Results	Complications
Bartochowska, 2015 [26]	60	Ir 192; PDR (49 pts) or HDR (11 pts)	Median: 20 Gy or 24 Gy	No	LC at 2 years: 27% OS at 2 years: 19%	33% acute toxicity 12% late toxicity
Huang, 2016 [27]	31	I 125; LDR, CT-guided	Mean: D ₉₀ 101 Gy	No	LC at 2 years: 45% OS at 2 years: 45%	No grade 3 or 4 toxicity
Tselis, 2011 [28]	74	Ir 192; HDR, CT-guided	Median: 30 Gy	No	LC at 2 years: 67% OS at 2 years: 19%	5% acute grade 3-4 toxicity 8% late grade 3-4 toxicity
Kolotas, 2007 [43]	49	Ir 192; HDR, CT-guided	Total: 30 Gy or 36 Gy	No	LC at 19 mos: 69% OS at 2 years: 31% OS at 3 years: 6%	16% acute grade 2-3 toxicity
Bollet, 2001 [44]	84	Ir 192; LDR	Mean: 56.5 Gy	No	LC at 2 years: 31% LC at 5 years: 0% OS at 2 years: 13% OS at 5 years: 1%	35% grade 3-4 toxicity 7% grade 5 (death)
Brachytherapy combined with surgical resection						
Teudt, 2016 [14]	9	HDR	Mean: 27 Gy	Yes	OS at 2 years: 78% OS at 5 years: 67%	No grade 3 or 4 toxicity
Kupferman, 2007 [24]	22	Ir 192; LDR	Median: 60 Gy	Yes	RC at 2 years: 67% OS at 5 years: 46%	14% acute grade 3-4 toxicity 18% late grade 3-4 toxicity
Nutting, 2006 [29]	72	Ir 192; LDR	Total: 60 Gy	Yes in 66 No in 6	LC at 5 years: 23% OS at 5 years: 23% <u>Non-surgical group:</u> LC at 2 years: 0% OS at 2 years: 0%	15% grade 3-4 toxicity
Cornes, 1996 [40]	39	Ir 192; LDR	Mean: 49.6 Gy	Yes	LC at 1 yr: 63% OS at 2 yrs: 38%	5% acute grade 3-4 toxicity 23% late grade 3-4 toxicity

Ir 192 – Iridium 192; I 125 – Iodine 125; Gy – Gray; LDR – low-dose-rate; HDR – high-dose-rate; PDR – pulsed-dose-rate; CT – computed tomography; LC – local control; RC – regional control; OS – overall survival

or other (4). Notably, many forms of brachytherapy did not use permanent implantable seeds but temporary catheters for the purpose of low-dose or high-dose-rate brachytherapy. Hemorrhage after catheter or tube removal was reported [15].

In contrast, Pham *et al.* [23] reported similar or lower rates of acute and chronic toxicity with ¹³¹Cs as compared to studies using ¹⁹²Ir and ¹²⁵I. Only 2 of 18 patients had grade 3 toxicity and none had grade 4-5 toxicity. When present, grade 1-2 complications were primarily dermatitis and hoarseness [23]. The risk to healthcare personnel and close family members has been minimized with isotopes like ¹³¹Cs compared to some previous brachytherapy isotopes, which required special precautions [31]. Many prior studies portrayed the effects of ¹²⁵I and ¹⁹²Ir brachytherapy on RHNC, but fewer studies have explored the use of permanent ¹³¹Cs. ¹³¹Cs offers a high-dose-rate, focused delivery, and less radiation exposure to adjacent tissues, personnel, and the patient's family [23,31]. Compared to ¹²⁵I, ¹³¹Cs has a dose-rate several times higher because of its similar energy but much lower half-life (9.69 days) [33]. This allows for ¹³¹Cs to deliver ~90% of its dose in a 33-day period, unlike 32% of the dose delivered by ¹²⁵I in that time frame [33]. This longer period of time exposes staff and family to potentially in-

creased radiation doses and may require special precautions. Suture strands containing ¹³¹Cs seeds can be fixed 0.5-1 cm apart within the tumor bed or come in a pre-embedded mesh (Figure 1) [31]. With brachytherapy done during salvage surgery, flaps also may help to prevent seed migration and dosimetry alterations. After implantation, seed placement can be monitored radiographically by generating a post-implant plan and toxicity monitored using RTOG common toxicity criteria [30]. In a study by Parasher *et al.* [31], 28 patients received a median number of twenty ¹³¹Cs seeds implanted at 0.5-1 cm distance for lung or HN cancer, with a median seed activity of 2.4 U. Median radiation exposure rate at the skin and at 1 meter was 0.43 mSv and 0.002 mSv, respectively. Minimum measurable radiation dose for OSL badge and TLD ring was 1 mrem and 30 mrem [31]. These minimal dose levels allow for patients to interact with family members and caregivers without special precautions like special lead protective equipment or isolation.

Discussion

Treatment options for RHNC remain limited. Many patients have undergone previous surgeries with or without radiation, chemotherapy, immunotherapies,




Fig. 1. A) Post-operative radiographic Cesium-131 seed dosimetry. B) Cesium-131 seed placement intra-operatively

and/or combination therapies, which limit retreatment options with radiation in general and EBRT in particular. The most common current treatment for curative intent is salvage surgery but may not be feasible for all cases [34,35,36]. Despite wide margins and G0 resection, locoregional recurrence still thwarts the curative efforts in the recurrent setting. Adding focused radiation directed at the resection margin during salvage surgery, while sparing toxicity to surrounding tissue, is in theory promising.

Brachytherapy has shown to be an effective treatment option with acceptable toxicity profiles for RHNC. When using brachytherapy alone, the reported LR control rates and OS rates at 2 years range from 27.5-92.5% and 18.2-43%, respectively [5,8,12,17,18,19,32,37,38,39]. Some studies have revealed brachytherapy to have enhanced and improved results when combined with salvage surgery [9,20,29,40]. Salvage surgery with brachytherapy as adjunctive therapy has 2-year LR control rates and OS rates ranging from 62-88% and 38-65.3%, respectively [6,9,10,11,13,20,21,22,23,41,42]. Studies by Narayana *et al.* and Rudzianskas *et al.* even compared LC and OS rates at 2 years between those who received brachytherapy with and without surgery and found improved rates in those also receiving surgery [9,20]. Although our analysis comparing locoregional recurrence rates of patients receiving surgery with brachytherapy vs. brachytherapy alone was not statistically significant, there was a trend of 38.8% vs. 49.0%. For brachytherapy use in patients with neck disease, LC

and OS rates at 2 years had likewise fair responses, with rates ranging from 27.3-67% and 13-78%, respectively [14,24,26,27,28,29,40,43,44]. Many of these studies had recurrent HNC with neck metastases that appeared inoperable or unmanageable with other therapies. Yet, analysis in these neck disease patients did show a statistically significant difference ($p = 0.015$) in locoregional recurrence rates (30.9% vs. 49.0%) between the combined treatment group and brachytherapy alone group, respectively.

In comparison, a review by Goodwin [34] described outcomes using salvage surgery alone, the most common salvage treatment. The weighted average of OS at 3 years was 37% (range: 28-80%) and at 5 years was 36.4% (range: 23-55%) [34]. Goodwin [34] also reported the weighted average of disease-free survival (DFS) rates at 2 years: 36.3% (range: 0-59%). Moreover, a literature review by Strojan *et al.* [35] examined local control and survival rates using conventional EBRT techniques and newer EBRT techniques like IMRT, with or without surgery, in recurrent HNC. Locoregional control rates and OS rates at 2 years using conventional radiation techniques with salvage surgery ranged from 13-74% (average: 38%) and 21-67% (average: 42.5%), respectively [35]. Newer RT techniques like IMRT have been showing better outcomes. For example, a study by Lee *et al.* [36] did show that IMRT had improved LR recurrence-free survival at 2 years over conventional RT techniques (52% vs. 20%) but did not significantly improve OS rates. In general, for patients receiving IMRT (with or without salvage

surgery), LR control and OS rates at 2 years ranged from 27-77% and 19.5-58%, respectively [35,45].

Similarly, to some brachytherapy studies, patients receiving reirradiation (either using conventional or newer RT techniques) with salvage surgery had improved outcomes. One phase III randomized trial by Janot *et al.* [46] divided patients into receiving salvage surgery alone or with adjuvant full-dose reirradiation with chemotherapy. Patients in the radiation arm (using conventional RT) had significantly improved LR control and DFS rates but no significant change in OS rates at 2 years [46]. Likewise, a study by Takiar *et al.* [47] showed improved disease control and survival outcomes using adjuvant IMRT after surgery (vs. surgery alone) in HNSCC patients with a 5-year OS rate of 57%.

Complications appear less frequent or severe compared to EBRT techniques including IMRT. In some brachytherapy studies, complication rates ranged as high as 50-60% for acute grade 1-2 toxicities including mucositis, dermatitis, hoarseness, infection, or hematoma [12,14,17,26,27]. Late grade 3-4 toxicities included complications like fistula, soft tissue necrosis, ORN, dysphagia, fibrosis, trismus, severe hemorrhage, and wound dehiscence, with rates ranging from 7-51% [9,10,12,17,18,20,21,28,29,37,40,42]. In the study by Takiar *et al.* [47] using IMRT for re-irradiation, there was a 5-year grade ≥ 3 toxicity rate of 48%, with three patients dying due to radiation toxicity. However, the majority of patients who now are receiving their first round of radiation with IMRT may likely have fewer overall complications, if or when they receive reirradiation. Literature is limited at this time to support this theory but will likely be shown in coming years.

We propose an algorithm to help guide treatment options for patients with RHNC (Figure 2). For example, for patients with resectable locoregional recurrence, consider surgical resection followed by brachytherapy or EBRT with or without chemotherapy or immunotherapy. Using brachytherapy versus external radiation techniques like IMRT may depend on the timing, type of previous RT, and location of recurrence. Despite the safety, ease of use, and promising efficacy advantages of ^{131}Cs , the use for specific RHNC populations requires further investigation. Unfortunately, the majority of published literature is retrospective, and the patient cohorts and treatment approaches used are very heterogeneous. In theory, combining a radiation boost to salvage surgery may provide dual therapy with less radiotherapy toxicity. ^{131}Cs with salvage surgery and other combinations with immunotherapy may offer a viable treatment pathway for RHNC for some patients and should be studied in future clinical trials. Our institution has recently been studying ^{131}Cs brachytherapy in RHNC with preliminary promising results. At this time, HDR and PDR appear to be the current techniques of choice due to optimization of implant dosimetry and geometric sparing with and reduced treatment time.

Conclusions

Treatment options for RHNC remain challenging. Although brachytherapy alone can be used to treat RHNC, locoregional control rates and overall survival rates are higher when brachytherapy is used as adjunctive therapy to salvage surgery. These rates remain comparable or improved when brachytherapy is utilized in




Fig. 2. Proposed algorithm for treating recurrent head and neck squamous cell carcinoma

neck disease with or without surgery. Regardless of the type of radiotherapy used, these patients continue to face significant risks for recurrence and complications including fistula formation, carotid rupture, dysphagia, osteoradionecrosis, and prolonged hospital stay [9,10,20,21,42,48,49,50]. With the recent introduction of isotopes with more favorable properties, complication rates have the potential to be lower while preserving local regional control rates.

Disclosure

Authors report no conflict of interest.

References

- Siegel R, Miller K, Jemal A. Cancer statistics. *CA Cancer J Clin* 2016; 66: 7-30.
- Siegel R, Miller K, Jemal A. Cancer statistics. *CA Cancer J Clin* 2017; 67: 7-30.
- Siegel R, Miller K, Jemal A. Cancer statistics. *CA Cancer J Clin* 2015; 65: 5-29.
- Vokes E, Weichselbaum R, Lippman S, Hong WK. Head and neck cancer. *N Engl J Med* 1993; 328: 184-194.
- Grimard L, Esche B, Lamothe A et al. Interstitial low-dose-rate brachytherapy in the treatment of recurrent head and neck malignancies. *Head Neck* 2006; 28: 888-895.
- Henderson P, Kutler D, Parashar B et al. Neither high-dose nor low-dose brachytherapy increases flap morbidity in salvage treatment of recurrent head and neck cancer. *J Contemp Brachytherapy* 2016; 8: 308-312.
- Quon H, Harrison LB. Brachytherapy in the treatment of head and neck cancer. *Oncology (Williston Park)* 2002; 16: 1379-1396.
- Strnad V, Lotter M, Kreppner S, Fietkau R. Reirradiation for recurrent head and neck cancer with salvage interstitial pulsed-dose-rate brachytherapy: Long-term results. *Strahlenther Onkol* 2015; 191: 495-500.
- Narayana A, Cohen G, Zaider M et al. High-dose-rate interstitial brachytherapy in recurrent and previously irradiated head and neck cancers-preliminary results. *Brachytherapy* 2007; 6: 157-163.
- Martinez-Fernandez M, Alcalde J, Cambeiro M et al. Perioperative high dose rate brachytherapy (PHDRB) in previously irradiated head and neck cancer: Results of a phase I/II re-irradiation study. *Radiother Oncol* 2017; 122: 255-259.
- Schieffe F, Hildebrandt G, Pohlmann S et al. Combination of surgical resection and HDR-brachytherapy in patients with recurrent or advanced head and neck carcinomas. *J Cranio-maxillofac Surg* 2008; 36: 285-292.
- Strnad V, Lotter M, Kreppner S, Fietkau R. Re-irradiation with interstitial pulsed-dose-rate brachytherapy for unresectable recurrent head and neck carcinoma. *Brachytherapy* 2014; 13: 187-195.
- Scala L, Hu K, Urken M et al. Intraoperative high-dose-rate radiotherapy in the management of locoregionally recurrent head and neck cancer. *Head Neck* 2013; 35: 485-492.
- Teudt I, Kovacs G, Ritter M et al. Intensity modulated perioperative HDR brachytherapy for recurrent and/or advanced head and neck metastases. *Eur Arch Otorhinolaryngol* 2016; 273: 2707-2715.
- Kasperts N, Slotman B, Leemans C et al. A review on re-irradiation for recurrent and second primary head and neck cancer. *Oral Oncol* 2005; 41: 225-243.
- Patel P, Salama J. Reirradiation for recurrent head and neck cancer. *Expert Rev Anticancer Ther* 2012; 12: 1177-1189.
- Puthawala A, Nisar Syed A, Gamie S et al. Interstitial low-dose-rate brachytherapy as a salvage treatment for recurrent head-and-neck cancers: long-term results. *Int J Radiat Oncol Biol Phys* 2001; 51: 354-362.
- Hepel J, Nisar Syed A, Puthawala A et al. Salvage high-dose-rate (HDR) brachytherapy for recurrent head-and-neck cancer. *Int J Radiat Oncol Biol Phys* 2005; 62: 1444-1450.
- Jiang P, Jiang Y, Wang J et al. Percutaneous ultrasonography-guided permanent iodine-125 implantation as salvage therapy for recurrent head and neck carcinomas. *Cancer Biother Radiopharm* 2011; 26: 753-757.
- Rudzianskas V, Inciura A, Juozaityte E et al. Reirradiation of recurrent head and neck cancer using high-dose-rate brachytherapy. *Acta Otorhinolaryngol Ital* 2012; 32: 297-303.
- Park R, Liberman F, Lee DJ et al. Iodine-125 seed implantation as an adjunct to surgery in advanced recurrent squamous cell cancer of the head and neck. *Laryngoscope* 1991; 101: 405-410.
- Pellizzon A, dos Santos Novaes P, Conte Maia M et al. Interstitial high-dose-rate brachytherapy combined with cervical dissection on head and neck cancer. *Head Neck* 2005; 27: 1035-1041.
- Pham A, Arora S, Wernicke G et al. Cesium-131 brachytherapy in high risk and recurrent head and neck cancers: first report of long-term outcomes. *J Contemp Brachytherapy* 2015; 7: 445-452.
- Kupferman M, Morrison W, Santillan A et al. The role of interstitial brachytherapy with salvage surgery for the management of recurrent head and neck cancers. *Cancer* 2007; 109: 2052-2057.
- Wong S, Machtay M, Li Y. Locally recurrent, previously irradiated head and neck cancer: concurrent re-irradiation and chemotherapy, or chemotherapy alone? *J Clin Oncol* 2006; 24: 2653-2658.
- Bartochowska A, Skowronek J, Wierzbicka M et al. Is there a place for brachytherapy in the salvage treatment of cervical lymph node metastases of head and neck cancers? *Brachytherapy* 2015; 14: 933-938.
- Huang H, Xu S, Li F et al. Clinical application of computed tomography-guided ¹²⁵I seed interstitial implantation for head and neck cancer patients with unmanageable cervical lymph node metastases. *Eur J Med Res* 2016; 21: 18.
- Tselis N, Ratka M, Vogt HG et al. Hypofractionated accelerated CT-guided interstitial ¹⁹²Ir-HDR-brachytherapy as re-irradiation in inoperable recurrent cervical lymphadenopathy from head and neck cancer. *Radiother Oncol* 2011; 98: 57-62.
- Nutting C, Horlock N, A'Hern R et al. Manually after-loaded ¹⁹²Ir low-dose rate brachytherapy after subtotal excision and flap reconstruction of recurrent cervical lymphadenopathy from head and neck cancer. *Radiother Oncol* 2006; 80: 39-42.
- Cox J, Stetz J, Pajak TF. Toxicity criteria of the Radiation Therapy Oncology Group (RTOG) and the European Organization for Research and Treatment of Cancer (EORTC). *Int J Radiat Oncol Biol Phys* 1995; 31: 1341-1346.
- Parasher B, Wernicke A, Pavese A et al. Cesium-131 permanent brachytherapy: dosimetric evaluation and radiation exposure to surgeons, radiation oncologists, and staff. *Brachytherapy* 2011; 10: 508-513.
- Zhu L, Jiang Y, Wang J et al. An investigation of ¹²⁵I seed permanent implantation for recurrent carcinoma in the head and neck after surgery and external beam radiotherapy. *World J Surg Oncol* 2013; 11: 60.
- Yondorf M, Schwartz T, Boockvar J et al. Radiation exposure and safety precautions following ¹³¹Cs brachytherapy in patients with brain tumors. *Health Phys* 2017; 112: 403-408.
- Goodwin WJ Jr. Salvage surgery for patients with recurrent squamous cell carcinoma of the upper aerodigestive tract: when do the ends justify the means? *Laryngoscope* 2000; 110: 1-18.

35. Strojan P, Corry J, Eisbruch A et al. Recurrent and second primary squamous cell carcinoma of the head and neck: when and how to irradiate. *Head Neck* 2015; 37: 134-150.
36. Lee N, Chan K, Bekelman J et al. Salvage re-irradiation for recurrent head and neck cancer. *Int J Radiat Oncol Biol Phys* 2007; 68: 731-740.
37. Glatzel M, Buntzel J, Schroder D et al. High-dose-rate brachytherapy in the treatment of recurrent and residual head and neck cancer. *Laryngoscope* 2002; 112: 1366-1371.
38. Meng N, Liang Jiang Y, Jie Wang J et al. Permanent implantation of iodine-125 seeds as a salvage therapy for recurrent head and neck carcinoma after radiotherapy. *Cancer Invest* 2012; 30: 236-242.
39. Wiegand S, Sesterhenn AM, Zimmermann AP et al. Interstitial HDR brachytherapy for advanced recurrent squamous cell carcinoma of the head and neck. *Anticancer Res* 2013; 33: 249-252.
40. Cornes P, Cox H, Rhys-Evans P et al. Salvage treatment for inoperable neck nodes in head and neck cancer using combined iridium-192 brachytherapy and surgical reconstruction. *Br J Surg* 1996; 83: 1620-1622.
41. Teckie S, Scala L, Ho F et al. High-dose-rate intraoperative brachytherapy and radical surgical resection in the management of recurrent head-and-neck cancer. *Brachytherapy* 2013; 12: 228-234.
42. Perry D, Chan K, Wolden S et al. High-dose-rate intraoperative radiation therapy for recurrent head-and-neck cancer. *Int J Radiat Oncol Biol Phys* 2010; 76: 1140-1146.
43. Kolotas C, Tselis N, Sommerlad M et al. Reirradiation for recurrent neck metastases of head-and-neck tumors using CT-guided interstitial ¹⁹²Ir HDR brachytherapy. *Strahlenther Onkol* 2007; 183: 69-75.
44. Bollet M, Lapeyre M, Marchal C et al. Cervical lymph node relapses of head-and-neck squamous cell carcinoma: is brachytherapy a therapeutic option? *Int J Radiat Oncol Biol Phys* 2001; 51: 1305-1312.
45. Sulman E, Schwartz D, Le T et al. IMRT reirradiation of head and neck cancer-disease control and morbidity outcomes. *Int J Radiat Oncol Biol Phys* 2009; 73: 399-409.
46. Janot F, de Raucourt D, Benhamou E et al. Randomized trial of postoperative reirradiation combined with chemotherapy after salvage surgery compared with salvage surgery alone in head and neck carcinoma. *J Clin Oncol* 2008; 26: 5518-5523.
47. Takiar V, Garden A, Ma D et al. Reirradiation of head and neck cancers with intensity modulated radiation therapy: outcomes and analyses. *Int J Radiat Oncol Biol Phys* 2016; 95: 1117-1131.
48. Vermorken JB, Specenier P. Optimal treatment for recurrent/metastatic head and neck cancer. *Ann Oncol* 2010; 21: 252-261.
49. Jeong S, Yoo EJ, Kim JY et al. Re-irradiation of unresectable recurrent head and neck cancer: Helical tomotherapy as image-guided intensity-modulated radiotherapy. *Rad Oncol J* 2013; 31: 206-215.
50. Vargo J, Kubicek G, Ferris R et al. Adjuvant stereotactic body radiotherapy + cetuximab following salvage surgery in previously irradiated head and neck cancer. *Laryngoscope* 2014; 124: 1579-1584.