

7-31-2023

Extracellular tRNA-Derived RNAs as Emerging Activators of Endosomal Toll-like Receptors: A Narrative Review

Justin Gumas
Thomas Jefferson University

Yohei Kirino
Thomas Jefferson University

Follow this and additional works at: <https://jdc.jefferson.edu/tjucompmedctrfp>

 Part of the [Genetic Processes Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Gumas, Justin and Kirino, Yohei, "Extracellular tRNA-Derived RNAs as Emerging Activators of Endosomal Toll-like Receptors: A Narrative Review" (2023). *Computational Medicine Center Faculty Papers*. Paper 52. <https://jdc.jefferson.edu/tjucompmedctrfp/52>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Computational Medicine Center Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.



Extracellular tRNA-derived RNAs as emerging activators of endosomal Toll-like receptors: a narrative review

Justin Gumas, Yohei Kirino

Computational Medicine Center, Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, PA, USA

Contributions: (I) Conception and design: Both authors; (II) Administrative support: None; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: Both authors; (V) Data analysis and interpretation: Both authors; (VI) Manuscript writing: Both authors; (VII) Final approval of manuscript: Both authors.

Correspondence to: Yohei Kirino, PhD. 1020 Locust Street, JAH Suite Rm222, Philadelphia, PA 19107, USA. Email: Yohei.Kirino@jefferson.edu.

Background and Objective: The innate immune system deploys various pattern-recognition receptors (PRRs), including Toll-like receptors (TLRs), to detect the invasion of pathogens and initiate protective responses. TLR7 and TLR8 are located within the endosome of immune cells and are activated by single-stranded RNAs (ssRNAs). In addition to foreign ssRNAs from bacteria and viruses, endogenous self-ssRNAs, such as microRNAs (miRNAs), have been shown to activate TLR7 and TLR8, but such endogenous ssRNA ligands have not yet been fully elucidated. This scientific knowledge gap is partly derived from the technical limitations of standard RNA-seq, particularly its inability to capture non-miRNA-short non-coding RNAs (sncRNAs) lacking 5'-phosphate and 3'-hydroxyl ends. However, recent advances in our understanding of “previously-hidden” sncRNAs, captured by RNA-seq using T4 polynucleotide kinase-treated RNA samples, have widened the pool of candidate ssRNA molecules that could act as endogenous ligands of ssRNA-sensing immune receptors. This has indeed been exemplified in the recent finding that, during immune response, transfer RNA (tRNA)-derived sncRNAs in macrophages are packaged into extracellular vesicles (EVs), and those tRNA-derived extracellular (ex-) sncRNAs can function as endogenous ligands of TLR7 when delivered to the endosome of recipient cells. In this review, we highlight advances in our understanding of the functional roles of tRNA-derived ex-sncRNAs as emerging activators of endosomal TLRs.

Methods: We searched PubMed for articles relevant to our topic and published prior to September 2022.

Key Content and Findings: tRNA-derived ex-sncRNAs could act as endogenous modulators of the immune system by activating endosomal TLRs in diverse pathological conditions, ranging from bacterial infection to cancers and neurological disorders.

Conclusions: tRNA-derived ex-sncRNAs may constitute a significant class of bioactive circulating RNAs, and they are promising candidates as biomarkers in various diseases. Further research is required to realize their full implication in human health and disease.

Keywords: tRNA-derived RNA; transfer RNA half (tRNA half); extracellular RNA; endosomal TLR; Toll-like receptor 7 (TLR7)

Received: 01 October 2022; Accepted: 10 March 2023. Published online: 31 July 2023.

doi: 10.21037/exrna-22-22

View this article at: <https://dx.doi.org/10.21037/exrna-22-22>

Introduction

One of the first lines of defense against invading pathogens begins when the pattern-recognition receptors (PRRs) of the innate immune system detect pathogen-associated molecular patterns (PAMPs), leading to initiation of

protective responses (1,2). Toll-like receptors (TLRs) are the most extensively studied PRRs, and are expressed in both innate immune cells (e.g., macrophages and dendritic cells) and non-immune cells (e.g., epithelial cells and fibroblast cells) (2-4). Among the 10 TLRs characterized in humans,

Table 1 Search information

Items	Specification
Date of search	August 31, 2022
Databases and other sources searched	PubMed
Search terms used	tRNA, tRNA half, tRF, tsRNA, exRNA, circulating RNA, immune response, TLR, TLR7, TLR8, neurodegeneration, cancer
Timeframe	1986–2022
Inclusion and exclusion criteria	Research and review manuscripts written in English were included, and articles written in a language other than English were excluded
Selection process	Both authors performed the search for suitable articles

TLR1, TLR2, TLR4, TLR5, TLR6, and TLR10 localize to the cell surface (surface TLRs), while TLR3, TLR7, TLR8, and TLR9 localize to endosomes and other intracellular compartments (endosomal TLRs). Each TLR specifically recognizes a different PAMP, ranging from polysaccharides and lipopeptides to nucleic acids (2-4). Endosomal TLRs are nucleic acid-sensing TLRs, and among those, TLR7 and TLR8 recognize single-stranded RNAs (ssRNAs) as their ligands (2-5). ssRNA recognition by these TLRs results in the recruitment of the adaptor protein MyD88, leading to NF- κ B-mediated transcription and downstream induction of interferon and cytokine production (2-4). While “foreign” ssRNAs from bacteria or viruses have been extensively studied as the ligands recognized by TLR7 and TLR8 (5-10), “endogenous” ssRNA ligands from host cells have not been fully characterized yet, and indeed the idea of endogenous ligands of PRRs remains controversial, as it complicates the traditional view of the role of PRRs in distinguishing self from non-self (11).

Once endogenous ssRNA molecules are packaged into extracellular vesicles (EVs), they can be delivered into the endosomes of recipient cells, where they can be sensed by TLR7 or TLR8. This has been observed with certain extracellular (ex-) microRNAs (miRNAs), which, upon delivery to recipient cell endosomes, become ligands for TLR7 and TLR8 and activate the downstream pathway (12,13). This miRNA-mediated TLR7/TLR8 activation is relevant not only to immune response (14,15) but has also been demonstrated to play roles in neurodegeneration (12), tumor growth and metastasis (16,17), autoimmunity (18), and pathobiology of various other diseases (19). Because miRNAs are the best-studied short non-coding RNAs (sncRNAs), it is natural that they have dominated current research on both ex-sncRNAs and endogenous ssRNA ligands of

TLRs. However, recent advances in our understanding of “previously-hidden” sncRNAs have widened the pool of candidate ssRNA molecules that could act as endogenous ligands of ssRNA-sensing immune receptors, exemplified in the recent finding that transfer RNA (tRNA)-derived sncRNAs can function as endogenous ligands of TLR7 (20). We present this article in accordance with the Narrative Review reporting checklist (available at <https://exrna.amegroups.com/article/view/10.21037/exrna-22-22/rc>).

Methods

To identify articles suitable for this review, we conducted a literature search via PubMed using search terms that included “tRNA”, “tRNA half”, “tRF”, “tsRNA”, “exRNA”, “circulating RNA”, “immune response”, “TLR”, “TLR7”, “TLR8”, “neurodegeneration”, and “cancer”. Articles written in English and published prior to August 31, 2022 were considered. These details are compiled in *Table 1*.

Most ex-sncRNAs are uncaptured by standard RNA-seq

Cellular sncRNA molecules generally possess either a hydroxyl group (OH), a monophosphate (P), or a 2',3'-cyclic phosphate (cP) at their termini (*Figure 1A*), and the terminal states of each sncRNA are determined by the catalytic machinery underlying the RNA cleavage that produces them (21,24). Although next-generation sequencing of RNA molecules (RNA-seq) has become a common tool to characterize RNA expression profiles, most sncRNA sequencing studies to date have relied on a standard small RNA-seq method in which 5'- and 3'-adaptors (AD) can be ligated only to 5'-P and 3'-OH ends of RNAs, respectively.




Figure 1 Infection-induced ex-5'-tRNA halves activate TLR7. (A) Chemical structures of general snRNA termini [modified from our previous review (21)]; (B) targeted snRNAs in standard RNA-seq, Phospho-RNA-seq, and cP-RNA-seq; (C) surface TLR stimulation culminates in activation of NF- κ B, leading to upregulation of ANG, which cleaves the anticodon-loops of tRNAs. The resultant 5'-tRNA halves are secreted into EVs, and then the ex-5'-tRNA halves are delivered into endosomes in recipient cells and activate TLR7. Following TLR7 activation, signal transduction proceeds through MyD88, leading to activation of a protein complex that includes members of the IRAK family and the TRAF family, along with IKK α (22). Activation of IRAKs, particularly IRAK1 and IRAK2, and of TRAF6 leads to activation of IRF7 and its subsequent translocation into the nucleus, where it promotes transcription of IFN (23). Formation of a protein complex including TRAF6 and the TAK1 leads to downstream activation of NF- κ B via degradation of its inhibitor. Liberated NF- κ B can then migrate to the nucleus, where it induces expression of inflammatory cytokines and promotes cellular immune response. TLR, Toll-like receptor; IRAK, IL-1R associated kinase; TRAF, tumor necrosis factor receptor-associated factor; IKK α , I κ B kinase α ; IRF7, interferon regulatory factor 7; IFN, type I interferon; TAK1, transforming growth factor- β -activated kinase 1.

While this standard method is suitable for efficient amplification and sequencing of 5'-P/3'-OH-containing snRNAs, such as miRNAs, it cannot capture RNAs with other terminal structures (i.e., 5'-OH, 3'-P, or cP) (Figure 1B). Due to this limitation, non-miRNA-sncRNAs lacking the 5'-P/3'-OH ends have been significantly underrepresented in many of the current snRNA analyses, and thus comprise a largely unexplored component in the

transcriptome.

This notion is especially important when it comes to the sequencing analyses of ex-sncRNAs. Human plasma samples contain many ex-sncRNAs which lack 5'-P or 3'-OH and are therefore uncaptured by standard RNA-seq (25). To sequence ex-sncRNAs with any given set of terminal phosphate states, Giraldez *et al.* developed Phospho-RNA-seq (25) (Figure 1B), which begins with the treatment of

RNA samples with T4 polynucleotide kinase (T4 PNK). Because T4 PNK possesses both 5'-phosphorylation (convert 5'-OH to 5'-P in the presence of ATP) and 3'-dephosphorylation (convert 3'-P/cP to 3'-OH) activities, the T4 PNK treatment leaves all RNA species with the 5'-P/3'-OH-ends, thus rendering them available for 5'- and 3'-AD ligation in subsequent small RNA-seq procedure. Phospho-RNA-seq has been successfully used to profile the expression of ex-sncRNAs in human plasma samples and their tissue specific signatures (25). Other human plasma sequencing studies also showed that the addition of T4 PNK treatment to the sequencing procedure significantly altered the proportion of the reads of the sncRNAs derived from tRNAs, messenger RNAs (mRNAs), ribosomal RNAs (rRNAs), and other non-coding RNAs (26,27), corroborating that standard RNA-seq without T4 PNK treatment cannot fully capture these ex-sncRNAs. Our study on EVs secreted from human monocyte-derived macrophages (HMDMs) further confirmed the necessity of T4 PNK treatment in ex-sncRNA sequencing (20). T4 PNK treatment was required to amplify the majority of the cDNAs derived from the EV ex-sncRNAs. Treatment with a mutant T4 PNK, which lacks 3'-dephosphorylation activity, resulted in dramatic reduction of cDNA yield, suggesting that most ex-sncRNAs in HMDM EVs are 3'-P- or cP-containing RNAs, and miRNAs and other 3'-OH containing RNAs comprise only a small part of those ex-sncRNA species packaged into the EVs (20). Based on these findings, recent ex-sncRNA sequencing studies have been utilizing T4 PNK treatment to make terminal formation of RNA molecules consistently 5'-P and 3'-OH, allowing for the capture of the full range of ex-sncRNAs.

tRNA-derived sncRNAs as a component of the expanding sncRNA world

It is well established that tRNAs, canonically understood as adaptor molecules of translational machinery, are also a source of abundant functional sncRNAs (28-31). The tRNA-derived sncRNAs are generated from mature tRNAs or their precursor transcripts, not as random degradation products but as specific functional molecules, and can be classified into two groups: tRNA halves and tRNA-derived fragments (tRFs) (28-31). The 30–35-nt tRNA halves are generally much more abundant than tRFs and are produced by endonucleolytic cleavage of the tRNA anticodon-loop. In mammalian cells, angiogenin (ANG), a member of the RNase A superfamily, is one of the major enzymes responsible for the anticodon

cleavage (32,33). Because ANG-mediated cleavage leaves a cP and 5'-P in 5'- and 3'-cleavage products (34), respectively, 5'-tRNA halves contain a 5'-P (from mature tRNAs) and a cP, while 3'-tRNA halves possess a 5'-OH and an amino acid (AA) at the 3'-end (from mature tRNAs), which has been experimentally validated (35). These terminal formations make tRNA half molecules the aforementioned “hidden class” of sncRNAs that are uncaptured by standard RNA-seq. Due to the presence of a cP in 5'-tRNA halves, a specific sequencing method for cP-containing RNAs, termed cP-RNA-seq (*Figure 1B*), has been utilized for the expressional characterization of 5'-tRNA halves (20,35-39). ANG-mediated tRNA cleavage can be induced by various biological factors/phenomena such as stress stimuli (32,33,38) and sex hormone signaling pathways (35). Although ANG is bound by its inhibitor ribonuclease/angiogenin inhibitor 1 (RNH1) in the cytoplasm, under stress conditions, degradation of RNH1 and/or translocation of ANG to stress granules, where it dissociates from RNH1, enable ANG-mediated tRNA cleavage (40-42). tRNA halves function in stress granule formation (43), regulation of translation (32,44), and promotion of cell proliferation (35); they are also associated with various disorders such as neurodegeneration (45,46). tRNA halves can further serve as direct precursors for shorter sncRNAs such as Piwi-interacting RNAs (piRNAs) (36,39).

tRFs are generally shorter than tRNA halves and can mainly be subclassified into 5'-tRFs, 3'-tRFs, and internal (i)-tRFs (28-31). While 5'- and 3'-tRFs are derived from 5'- and 3'-parts of mature tRNAs, respectively, i-tRFs are derived wholly from internal parts of mature tRNAs. Although Dicer and ANG are known to be involved in the biogenesis of some tRFs (47,48), detailed regulatory mechanisms and other biogenesis factors for tRFs remain elusive, leaving terminal phosphate states of tRFs undefined. This further emphasizes the importance of inclusive sequencing methods such as Phospho-RNA-seq for the comprehensive profiling of tRNA-derived sncRNAs. While many tRFs have been shown to function as miRNAs or piRNAs by binding to AGO or PIWI proteins (29,36,39), tRFs further have various functions, such as regulating mRNA stability or translation, preventing cell apoptosis, and promoting viral infection, and their dysregulation is involved in various diseases (28,30,31,49).

tRNA-derived sncRNAs as abundant ex-RNAs

tRNA-derived sncRNAs are increasingly recognized as an abundant class of ex-RNAs, released from cells under

diverse conditions and packaged into EVs (50-52) or bound to RNA-binding proteins (RNPs) (53) or lipoprotein particles (LPPs) (54). Functional roles of some of these tRNA-derived ex-sncRNAs have also been identified. For example, ex-5'-tRNA halves and ex-5'-tRFs contained in the vesicles deriving from the epididymis, termed epididymosomes, are transferred to maturing sperm (55) and regulate gene expressions in embryonic stem cells and embryos (55-57). tRNA derived ex-sncRNAs further appear to have potential as biomarkers in various disease contexts (58). For example, tRNA-derived ex-sncRNAs are differentially accumulated in breast cancer (59), prostate cancer (60), lung cancer (61), and chronic kidney disease (62), compared to healthy individuals. For further discussion of tRNA-derived ex-sncRNAs, we recommend a review from Tosar and Cayota (63). Experimental results on tRNA-derived ex-sncRNAs have continuously accumulated; more recent studies have provided further evidence that tRNA-derived ex-sncRNAs could be useful prognostic and diagnostic biomarkers in breast cancer (64-66), and have also shown their differential accumulation in metastatic hypopharyngeal cancer (67), systemic lupus erythematosus (68), and ischemic kidney injury (69).

Infection-induced ex-tRNA halves activate TLR7

We recently identified a novel role of ex-tRNA halves as activators of endosomal TLR (20). We demonstrated that infection of HMDMs with *Mycobacterium bovis* BCG and surface TLR activation by treatment with lipopolysaccharide (LPS) or peptidoglycan (PGN) upregulate the NF- κ B-mediated transcription of *ANG*, leading to accumulation of tRNA half molecules in HMDMs and their secreted EVs. In sequencing data of T4 PNK-treated RNAs from the EVs, 5'-tRNA halves comprised over 96% of tRNA-derived reads [in EV #1 library (20)], while 3'-tRNA halves (1.3%), 5'-tRFs (0.87%), 3'-tRFs (0.63%), and i-tRFs (0.99%) were minor species. The mechanisms underlying the specific and selective packaging of tRNA-derived sncRNAs into EVs are unknown, but specific RNA binding proteins, such as YBX1 that binds to 5'-tRNA halves (44) and functions in RNA sorting into EVs (70), could be involved in the mechanisms. Most ex-5'-tRNA halves were derived from a focused subset of cytoplasmic tRNAs, including tRNA^{ValCAC/AAC}, tRNA^{GlyGCC}, tRNA^{HisGUG}, and tRNA^{GluCUC}, which in aggregate were the sources of approximately 90% of the identified ex-5'-tRNA halves. One of the most remarkable characteristics of ex-tRNA halves is their

abundance. The ex-5'-tRNA^{HisGUG} half was >210 times more abundant than miR-150, one of the most abundant miRNAs in HMDM-derived EVs (20). While it has been demonstrated that ex-miRNAs can act as ligands for endosomal TLRs, the much greater abundance of ex-tRNA halves suggests that they might constitute a more significant, biologically relevant class of endogenous ligands of these immune receptors. Indeed, the EV-contained ex-tRNA halves were experimentally confirmed to be delivered into endosomes in recipient HMDMs where the 5'-tRNA^{HisGUG} half strongly activates TLR7 (20). The activity of the 5'-tRNA^{HisGUG} half is as high as that of HIV-derived ssRNA40, a widely used positive control ssRNA known as a strong activator of endosomal TLRs (5), suggesting that 5'-tRNA^{HisGUG} half could have the capacity to produce an immune response.

Expressional induction and secretion of tRNA halves are not limited to cell culture settings but have been further observed in actual pathological situations. By developing a sensitive multiplex tRNA half quantification method, we revealed an approximately 1,000-fold enrichment of plasma ex-5'-tRNA halves in patients infected with *Mycobacterium tuberculosis* (20,71). A dramatic increase in the levels of serum ex-5'-tRNA halves has also been observed in LPS-injected mice and monkeys and in patients experiencing active, but not quiescent, hepatitis B virus infection (72). Because upregulation of 5'-tRNA half expression has been reported upon infection with respiratory syncytial virus (73,74), *Rickettsia* (75), and hepatitis B and C viruses (76), it seems possible that induction of 5'-tRNA halves and their secretion as ex-5'-tRNA halves could be a universal phenomenon among infectious diseases. These studies suggest a novel role of ex-5'-tRNA half molecules as “immune activators,” but further studies are required to fully unveil the immune response pathways mediated by tRNA-derived ex-sncRNAs.

Future perspectives

The study of endogenous ex-sncRNA ligands of immune receptors is still at an initial stage and remains an area of some controversy. Likewise, we are just beginning to appreciate the previously hidden classes of sncRNAs not detected by standard RNA-seq methods. We expect that the 5'-tRNA^{HisGUG} half is just the first example of an immune receptor-simulating tRNA-derived ex-sncRNA, and other tRNA halves and tRFs could be found to be similarly immunostimulatory through the TLR7 or TLR8 axis. The 5'-tRNA^{HisGUG} half strongly activates TLR7, but not

TLR8 (20), probably because of differences in ligand specificity for TLR7 and TLR8 (6,77,78). Future exploration of immune stimulatory tRNA-derived sncRNAs should include review of studies looking into the characteristics of ssRNA ligands of TLR7 and TLR8.

Beyond tRNA-derived sncRNAs, our first genome-wide identification of cP-containing sncRNAs revealed abundant expression of rRNA- and mRNA-derived sncRNAs in various tissues (37), and, indeed, not only mature rRNAs (79) but also rRNA- and mRNA-derived sncRNAs have been shown to be abundantly secreted as ex-RNAs (26,80,81). These other unexplored classes of ex-sncRNAs should further be investigated as potential activators of TLR7 and TLR8.

Endosomal TLR3 also recognizes RNA molecules, but it binds to double-stranded RNAs (dsRNAs) (82). Although ssRNAs containing partial stem structures can also be TLR3 ligands, at least 40–50-base-pairs (bp) length of dsRNA molecule is required to ensure stable complex formation with TLR3 (83,84), and more than 90-bp length of dsRNA is required to activate TLR3 signal transduction in dendritic cell maturation (85). Therefore, ex-sncRNAs shorter than mature tRNAs are unlikely to participate in the TLR3 signaling. However, exRNAs more broadly should be considered as potentially major endogenous ligands of TLR3, as exemplified in a study showing that dsRNAs released from necrotic polymorphonuclear neutrophils induce inflammatory response in macrophages (86).

Although mature tRNAs have been reported to be incorporated into EVs (70,87), unlike the 5'-tRNA^{HisGUG} half, the full-length tRNA^{HisGUG} was incapable of stimulating endosomal TLR (20), possibly due to its rigid secondary and tertiary structures. Therefore, shortening mature tRNA^{HisGUG} into less-rigid 5'-half molecules by anticodon-cleavage is crucial to produce immunostimulatory sncRNAs. ANG is highly enriched in the EVs derived from aggressive brain tumor cells (52), while another RNase A superfamily member, RNase 1, has recently been shown to cleave non-vesicular tRNAs (88). Further research on immune responsive ribonucleases which cleave tRNAs, rRNAs, mRNAs, or other substrate RNAs to produce immune responsive ex-sncRNAs is necessary to fully understand the regulation and functional consequences of immune response mediated by these ex-sncRNAs.

TLR7 and TLR8 are involved in various biological processes and diseases (89). The symptom severity of COVID-19 is associated with TLR7 mutations (90–92). The genetic polymorphisms in TLR7 and TLR8 genes, which reduce the TLR activity, increase susceptibility to

Mtb infection (93), and upregulation and stimulation of TLR7 in macrophages suppress Mtb growth (94,95). TLR7 and TLR8 have been implicated in the progression of Parkinson's disease (96) and in Alzheimer's disease (12,97,98), possibly playing a role in promoting neuroinflammation and autoimmunity. A recent study also linked a specific gain-of-function mutation in TLR7 with severe systemic lupus erythematosus (99). Furthermore, missense mutations in TLR7 have been reported in the tumors of a subset of esophageal adenocarcinoma patients (100), and cellular immune response through activation of TLR7 and TLR8 can be directed against tumors (101). Despite these known linkages between TLR7/TLR8 and pathobiology of diseases, the endogenous ligands of these ssRNA-sensing TLRs remain poorly understood. The growing evidence of TLR7 and TLR8 involvement in the pathogenesis of non-infectious diseases highlights the need for a better understanding of the endogenous ligands of these receptors. Further research into tRNA-derived ex-sncRNAs, aided by upgraded sequencing methods, is necessary to identify these ligands and to gain a fuller understanding of the interplay between ex-RNAs, TLRs, and the immune system in the body's response to diverse pathologies.

Acknowledgments

We are grateful to Megumi Shigematsu and Takuya Kawamura (Thomas Jefferson University) for helpful discussions.

Funding: Work in the Kirino lab on this topic is currently supported by the National Institutes of Health Grants (Nos. GM106047, HL150560, AI151641, and AI168975 to YK) and the American Cancer Society Research Scholar Grant (No. RSG-17-059-01-RMC to YK).

Footnote

Provenance and Peer Review: This article was commissioned by the Guest Editors (Silvano Sozzani and Daniela Bosisio) for the series “Extracellular MiRNAs as Activators of Innate Immune Receptors” published in *ExRNA*. The article has undergone external peer review.

Reporting Checklist: The authors have completed the Narrative Review reporting checklist. Available at <https://exrna.amegroups.com/article/view/10.21037/exrna-22-22/rc>

Conflicts of Interest: Both authors have completed the

ICMJE uniform disclosure form (available at <https://exrna.amegrouops.com/article/view/10.21037/exrna-22-22/coif>). The series “Extracellular MiRNAs as Activators of Innate Immune Receptors” was commissioned by the editorial office without any funding or sponsorship. YK reports funding from National Institutes of Health Grants (Nos. GM106047, HL150560, AI151641, and AI168975) and the American Cancer Society Research Scholar Grant (No. RSG-17-059-01-RMC). The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Brubaker SW, Bonham KS, Zanoni I, et al. Innate immune pattern recognition: a cell biological perspective. *Annu Rev Immunol* 2015;33:257-90.
2. Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol* 2010;11:373-84.
3. Kawasaki T, Kawai T. Toll-like receptor signaling pathways. *Front Immunol* 2014;5:461.
4. Satoh T, Akira S. Toll-Like Receptor Signaling and Its Inducible Proteins. *Microbiol Spectr* 2016;4. doi: 10.1128/microbiolspec.MCHD-0040-2016.
5. Heil F, Hemmi H, Hochrein H, et al. Species-specific recognition of single-stranded RNA via toll-like receptor 7 and 8. *Science* 2004;303:1526-9.
6. Zhang Z, Ohto U, Shibata T, et al. Structural Analysis Reveals that Toll-like Receptor 7 Is a Dual Receptor for Guanosine and Single-Stranded RNA. *Immunity* 2016;45:737-48.
7. Lund JM, Alexopoulou L, Sato A, et al. Recognition of single-stranded RNA viruses by Toll-like receptor 7. *Proc Natl Acad Sci U S A* 2004;101:5598-603.
8. Melchjorsen J, Jensen SB, Malmgaard L, et al. Activation of innate defense against a paramyxovirus is mediated by RIG-I and TLR7 and TLR8 in a cell-type-specific manner. *J Virol* 2005;79:12944-51.
9. Triantafilou K, Orthopoulos G, Vakakis E, et al. Human cardiac inflammatory responses triggered by Coxsackie B viruses are mainly Toll-like receptor (TLR) 8-dependent. *Cell Microbiol* 2005;7:1117-26.
10. Gantier MP, Irving AT, Kaparakis-Liaskos M, et al. Genetic modulation of TLR8 response following bacterial phagocytosis. *Hum Mutat* 2010;31:1069-79.
11. Rifkin IR, Leadbetter EA, Busconi L, et al. Toll-like receptors, endogenous ligands, and systemic autoimmune disease. *Immunol Rev* 2005;204:27-42.
12. Lehmann SM, Krüger C, Park B, et al. An unconventional role for miRNA: let-7 activates Toll-like receptor 7 and causes neurodegeneration. *Nat Neurosci* 2012;15:827-35.
13. Fabbri M, Paone A, Calore F, et al. A new role for microRNAs, as ligands of Toll-like receptors. *RNA Biol* 2013;10:169-74.
14. Feng Y, Zou L, Yan D, et al. Extracellular MicroRNAs Induce Potent Innate Immune Responses via TLR7/MyD88-Dependent Mechanisms. *J Immunol* 2017;199:2106-17.
15. Zou L, He J, Gu L, et al. Brain innate immune response via miRNA-TLR7 sensing in polymicrobial sepsis. *Brain Behav Immun* 2022;100:10-24.
16. Fabbri M. TLRs as miRNA receptors. *Cancer Res* 2012;72:6333-7.
17. Casadei L, Calore F, Creighton CJ, et al. Exosome-Derived miR-25-3p and miR-92a-3p Stimulate Liposarcoma Progression. *Cancer Res* 2017;77:3846-56.
18. Wu WC, Song SJ, Zhang Y, et al. Role of Extracellular Vesicles in Autoimmune Pathogenesis. *Front Immunol* 2020;11:579043.
19. Groot M, Lee H. Sorting Mechanisms for MicroRNAs into Extracellular Vesicles and Their Associated Diseases. *Cells* 2020;9:1044.
20. Pawar K, Shigematsu M, Sharbati S, et al. Infection-induced 5'-half molecules of tRNA^{His}GUG activate Toll-like receptor 7. *PLoS Biol* 2020;18:e3000982.
21. Shigematsu M, Kirino Y. Making Invisible RNA Visible: Discriminative Sequencing Methods for RNA Molecules with Specific Terminal Formations. *Biomolecules* 2022;12:611.
22. Balka KR, De Nardo D. Understanding early TLR signaling through the Myddosome. *J Leukoc Biol*

- 2019;105:339-51.
23. Gilliet M, Cao W, Liu YJ. Plasmacytoid dendritic cells: sensing nucleic acids in viral infection and autoimmune diseases. *Nat Rev Immunol* 2008;8:594-606.
 24. Shigematsu M, Kawamura T, Kirino Y. Generation of 2',3'-Cyclic Phosphate-Containing RNAs as a Hidden Layer of the Transcriptome. *Front Genet* 2018;9:562.
 25. Giraldez MD, Spengler RM, Etheridge A, et al. Phospho-RNA-seq: a modified small RNA-seq method that reveals circulating mRNA and lncRNA fragments as potential biomarkers in human plasma. *EMBO J* 2019;38:e101695.
 26. Akat KM, Lee YA, Hurley A, et al. Detection of circulating extracellular mRNAs by modified small-RNA-sequencing analysis. *JCI Insight* 2019;5:e127317.
 27. Qin Y, Yao J, Wu DC, et al. High-throughput sequencing of human plasma RNA by using thermostable group II intron reverse transcriptases. *RNA* 2016;22:111-28.
 28. Kumar P, Kuscic C, Dutta A. Biogenesis and Function of Transfer RNA-Related Fragments (tRFs). *Trends Biochem Sci* 2016;41:679-89.
 29. Shigematsu M, Kirino Y. tRNA-Derived Short Non-coding RNA as Interacting Partners of Argonaute Proteins. *Gene Regul Syst Bio* 2015;9:27-33.
 30. Anderson P, Ivanov P. tRNA fragments in human health and disease. *FEBS Lett* 2014;588:4297-304.
 31. Magee R, Rigoutsos I. On the expanding roles of tRNA fragments in modulating cell behavior. *Nucleic Acids Res* 2020;48:9433-48.
 32. Yamasaki S, Ivanov P, Hu GF, et al. Angiogenin cleaves tRNA and promotes stress-induced translational repression. *J Cell Biol* 2009;185:35-42.
 33. Fu H, Feng J, Liu Q, et al. Stress induces tRNA cleavage by angiogenin in mammalian cells. *FEBS Lett* 2009;583:437-42.
 34. Shapiro R, Riordan JF, Vallee BL. Characteristic ribonucleolytic activity of human angiogenin. *Biochemistry* 1986;25:3527-32.
 35. Honda S, Loher P, Shigematsu M, et al. Sex hormone-dependent tRNA halves enhance cell proliferation in breast and prostate cancers. *Proc Natl Acad Sci U S A* 2015;112:E3816-25.
 36. Honda S, Kawamura T, Loher P, et al. The biogenesis pathway of tRNA-derived piRNAs in Bombyx germ cells. *Nucleic Acids Res* 2017;45:9108-20.
 37. Shigematsu M, Morichika K, Kawamura T, et al. Genome-wide identification of short 2',3'-cyclic phosphate-containing RNAs and their regulation in aging. *PLoS Genet* 2019;15:e1008469.
 38. Shigematsu M, Kirino Y. Oxidative stress enhances the expression of 2',3'-cyclic phosphate-containing RNAs. *RNA Biol* 2020;17:1060-9.
 39. Shigematsu M, Kawamura T, Morichika K, et al. RNase κ promotes robust piRNA production by generating 2',3'-cyclic phosphate-containing precursors. *Nat Commun* 2021;12:4498.
 40. Saikia M, Hatzoglou M. The Many Virtues of tRNA-derived Stress-induced RNAs (tiRNAs): Discovering Novel Mechanisms of Stress Response and Effect on Human Health. *J Biol Chem* 2015;290:29761-8.
 41. Saikia M, Krokowski D, Guan BJ, et al. Genome-wide identification and quantitative analysis of cleaved tRNA fragments induced by cellular stress. *J Biol Chem* 2012;287:42708-25.
 42. Pizzo E, Sarcinelli C, Sheng J, et al. Ribonuclease/angiogenin inhibitor 1 regulates stress-induced subcellular localization of angiogenin to control growth and survival. *J Cell Sci* 2013;126:4308-19.
 43. Emara MM, Ivanov P, Hickman T, et al. Angiogenin-induced tRNA-derived stress-induced RNAs promote stress-induced stress granule assembly. *J Biol Chem* 2010;285:10959-68.
 44. Ivanov P, Emara MM, Villen J, et al. Angiogenin-induced tRNA fragments inhibit translation initiation. *Mol Cell* 2011;43:613-23.
 45. Blanco S, Dietmann S, Flores JV, et al. Aberrant methylation of tRNAs links cellular stress to neurodevelopmental disorders. *EMBO J* 2014;33:2020-39.
 46. Hogg MC, Rayner M, Susdalzew S, et al. 5'ValCAC tRNA fragment generated as part of a protective angiogenin response provides prognostic value in amyotrophic lateral sclerosis. *Brain Commun* 2020;2:fcaa138.
 47. Haussecker D, Huang Y, Lau A, et al. Human tRNA-derived small RNAs in the global regulation of RNA silencing. *RNA* 2010;16:673-95.
 48. Li Z, Ender C, Meister G, et al. Extensive terminal and asymmetric processing of small RNAs from rRNAs, snoRNAs, snRNAs, and tRNAs. *Nucleic Acids Res* 2012;40:6787-99.
 49. Yu X, Xie Y, Zhang S, et al. tRNA-derived fragments: Mechanisms underlying their regulation of gene expression and potential applications as therapeutic targets in cancers and virus infections. *Theranostics* 2021;11:461-9.
 50. Nolte-t Hoen EN, Buermans HP, Waasdorp M, et al. Deep sequencing of RNA from immune cell-derived vesicles uncovers the selective incorporation of small non-coding RNA biotypes with potential regulatory functions.

- Nucleic Acids Res 2012;40:9272-85.
51. Baglio SR, Rooijers K, Koppers-Lalic D, et al. Human bone marrow- and adipose-mesenchymal stem cells secrete exosomes enriched in distinctive miRNA and tRNA species. *Stem Cell Res Ther* 2015;6:127.
 52. Wei Z, Batagov AO, Schinelli S, et al. Coding and noncoding landscape of extracellular RNA released by human glioma stem cells. *Nat Commun* 2017;8:1145.
 53. Tosar JP, Gámbaro F, Sanguinetti J, et al. Assessment of small RNA sorting into different extracellular fractions revealed by high-throughput sequencing of breast cell lines. *Nucleic Acids Res* 2015;43:5601-16.
 54. Allen RM, Zhao S, Ramirez Solano MA, et al. Bioinformatic analysis of endogenous and exogenous small RNAs on lipoproteins. *J Extracell Vesicles* 2018;7:1506198.
 55. Sharma U, Sun F, Conine CC, et al. Small RNAs Are Trafficked from the Epididymis to Developing Mammalian Sperm. *Dev Cell* 2018;46:481-494.e6.
 56. Sharma U, Conine CC, Shea JM, et al. Biogenesis and function of tRNA fragments during sperm maturation and fertilization in mammals. *Science* 2016;351:391-6.
 57. Chen Q, Yan M, Cao Z, et al. Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic disorder. *Science* 2016;351:397-400.
 58. Dhahbi JM, Spindler SR, Atamna H, et al. 5' tRNA halves are present as abundant complexes in serum, concentrated in blood cells, and modulated by aging and calorie restriction. *BMC Genomics* 2013;14:298.
 59. Dhahbi JM, Spindler SR, Atamna H, et al. Deep Sequencing of Serum Small RNAs Identifies Patterns of 5' tRNA Half and YRNA Fragment Expression Associated with Breast Cancer. *Biomark Cancer* 2014;6:37-47.
 60. Dhahbi JM, Atamna H, Selth LA. Data Mining of Small RNA-Seq Suggests an Association Between Prostate Cancer and Altered Abundance of 5' Transfer RNA Halves in Seminal Fluid and Prostatic Tissues. *Biomark Cancer* 2018;10:1179299X18759545.
 61. Shao Y, Sun Q, Liu X, et al. tRF-Leu-CAG promotes cell proliferation and cell cycle in non-small cell lung cancer. *Chem Biol Drug Des* 2017;90:730-8.
 62. Khurana R, Ranches G, Schafferer S, et al. Identification of urinary exosomal noncoding RNAs as novel biomarkers in chronic kidney disease. *RNA* 2017;23:142-52.
 63. Tosar JP, Cayota A. Extracellular tRNAs and tRNA-derived fragments. *RNA Biol* 2020;17:1149-67.
 64. Wang J, Ma G, Li M, et al. Plasma tRNA Fragments Derived from 5' Ends as Novel Diagnostic Biomarkers for Early-Stage Breast Cancer. *Mol Ther Nucleic Acids* 2020;21:954-64.
 65. Koi Y, Tsutani Y, Nishiyama Y, et al. Predicting the presence of breast cancer using circulating small RNAs, including those in the extracellular vesicles. *Cancer Sci* 2020;111:2104-15.
 66. Wang J, Ma G, Ge H, et al. Circulating tRNA-derived small RNAs (tsRNAs) signature for the diagnosis and prognosis of breast cancer. *NPJ Breast Cancer* 2021;7:4.
 67. Xi J, Zeng Z, Li X, et al. Expression and Diagnostic Value of tRNA-Derived Fragments Secreted by Extracellular Vesicles in Hypopharyngeal Carcinoma. *Onco Targets Ther* 2021;14:4189-99.
 68. Yang P, Zhang X, Chen S, et al. A Novel Serum tsRNA for Diagnosis and Prediction of Nephritis in SLE. *Front Immunol* 2021;12:735105.
 69. Lee HK, Lee BR, Lee TJ, et al. Differential release of extracellular vesicle tRNA from oxidative stressed renal cells and ischemic kidneys. *Sci Rep* 2022;12:1646.
 70. Shurtleff MJ, Yao J, Qin Y, et al. Broad role for YBX1 in defining the small noncoding RNA composition of exosomes. *Proc Natl Acad Sci U S A* 2017;114:E8987-95.
 71. Kawamura T, Shigematsu M, Kirino Y. In vitro production and multiplex quantification of 2',3'-cyclic phosphate-containing 5'-tRNA half molecules. *Methods* 2022;203:335-41.
 72. Zhang Y, Zhang Y, Shi J, et al. Identification and characterization of an ancient class of small RNAs enriched in serum associating with active infection. *J Mol Cell Biol* 2014;6:172-4.
 73. Zhou J, Liu S, Chen Y, et al. Identification of two novel functional tRNA-derived fragments induced in response to respiratory syncytial virus infection. *J Gen Virol* 2017;98:1600-10.
 74. Deng J, Ptashkin RN, Chen Y, et al. Respiratory Syncytial Virus Utilizes a tRNA Fragment to Suppress Antiviral Responses Through a Novel Targeting Mechanism. *Mol Ther* 2015;23:1622-9.
 75. Gong B, Lee YS, Lee I, et al. Compartmentalized, functional role of angiogenin during spotted fever group rickettsia-induced endothelial barrier dysfunction: evidence of possible mediation by host tRNA-derived small noncoding RNAs. *BMC Infect Dis* 2013;13:285.
 76. Selitsky SR, Baran-Gale J, Honda M, et al. Small tRNA-derived RNAs are increased and more abundant than microRNAs in chronic hepatitis B and C. *Sci Rep* 2015;5:7675.
 77. Tanji H, Ohto U, Shibata T, et al. Toll-like receptor 8 senses degradation products of single-stranded RNA. *Nat*

- Struct Mol Biol 2015;22:109-15.
78. Greulich W, Wagner M, Gaidt MM, et al. TLR8 Is a Sensor of RNase T2 Degradation Products. *Cell* 2019;179:1264-1275.e13.
 79. Preissner KT, Fischer S. Functions and cellular signaling by ribosomal extracellular RNA (rexRNA): Facts and hypotheses on a non-typical DAMP. *Biochim Biophys Acta Mol Cell Res* 2023;1870:119408.
 80. Li M, Zeringer E, Barta T, et al. Analysis of the RNA content of the exosomes derived from blood serum and urine and its potential as biomarkers. *Philos Trans R Soc Lond B Biol Sci* 2014;369:20130502.
 81. Jia J, Yang S, Huang J, et al. Distinct Extracellular RNA Profiles in Different Plasma Components. *Front Genet* 2021;12:564780.
 82. Alexopoulou L, Holt AC, Medzhitov R, et al. Recognition of double-stranded RNA and activation of NF-kappaB by Toll-like receptor 3. *Nature* 2001;413:732-8.
 83. Liu L, Botos I, Wang Y, et al. Structural basis of toll-like receptor 3 signaling with double-stranded RNA. *Science* 2008;320:379-81.
 84. Leonard JN, Ghirlando R, Askins J, et al. The TLR3 signaling complex forms by cooperative receptor dimerization. *Proc Natl Acad Sci U S A* 2008;105:258-63.
 85. Jelinek I, Leonard JN, Price GE, et al. TLR3-specific double-stranded RNA oligonucleotide adjuvants induce dendritic cell cross-presentation, CTL responses, and antiviral protection. *J Immunol* 2011;186:2422-9.
 86. Cavassani KA, Ishii M, Wen H, et al. TLR3 is an endogenous sensor of tissue necrosis during acute inflammatory events. *J Exp Med* 2008;205:2609-21.
 87. Tosar JP, Segovia M, Castellano M, et al. Fragmentation of extracellular ribosomes and tRNAs shapes the extracellular RNAome. *Nucleic Acids Res* 2020;48:12874-88.
 88. Nechooshtan G, Yunusov D, Chang K, et al. Processing by RNase 1 forms tRNA halves and distinct Y RNA fragments in the extracellular environment. *Nucleic Acids Res* 2020;48:8035-49.
 89. Lind NA, Rael VE, Pestal K, et al. Regulation of the nucleic acid-sensing Toll-like receptors. *Nat Rev Immunol* 2022;22:224-35.
 90. Asano T, Boisson B, Onodi F, et al. X-linked recessive TLR7 deficiency in ~1% of men under 60 years old with life-threatening COVID-19. *Sci Immunol* 2021;6:eabl4348.
 91. Solanich X, Vargas-Parra G, van der Made CI, et al. Genetic Screening for TLR7 Variants in Young and Previously Healthy Men With Severe COVID-19. *Front Immunol* 2021;12:719115.
 92. Fallerini C, Daga S, Mantovani S, et al. Association of Toll-like receptor 7 variants with life-threatening COVID-19 disease in males: findings from a nested case-control study. *Elife* 2021;10:e67569.
 93. Lai YF, Lin TM, Wang CH, et al. Functional polymorphisms of the TLR7 and TLR8 genes contribute to Mycobacterium tuberculosis infection. *Tuberculosis (Edinb)* 2016;98:125-31.
 94. Bao M, Yi Z, Fu Y. Activation of TLR7 Inhibition of Mycobacterium Tuberculosis Survival by Autophagy in RAW 264.7 Macrophages. *J Cell Biochem* 2017;118:4222-9.
 95. Lee HJ, Kang SJ, Woo Y, et al. TLR7 Stimulation With Imiquimod Induces Selective Autophagy and Controls Mycobacterium tuberculosis Growth in Mouse Macrophages. *Front Microbiol* 2020;11:1684.
 96. Campolo M, Filippone A, Biondo C, et al. TLR7/8 in the Pathogenesis of Parkinson's Disease. *Int J Mol Sci* 2020;21:9384.
 97. Frank S, Copanaki E, Burbach GJ, et al. Differential regulation of toll-like receptor mRNAs in amyloid plaque-associated brain tissue of aged APP23 transgenic mice. *Neurosci Lett* 2009;453:41-4.
 98. Dembny P, Newman AG, Singh M, et al. Human endogenous retrovirus HERV-K(HML-2) RNA causes neurodegeneration through Toll-like receptors. *JCI Insight* 2020;5:e131093.
 99. Brown GJ, Cañete PF, Wang H, et al. TLR7 gain-of-function genetic variation causes human lupus. *Nature* 2022;605:349-56.
 100. Fels Elliott DR, Perner J, Li X, et al. Impact of mutations in Toll-like receptor pathway genes on esophageal carcinogenesis. *PLoS Genet* 2017;13:e1006808.
 101. Schön MP, Schön M. TLR7 and TLR8 as targets in cancer therapy. *Oncogene* 2008;27:190-9.

doi: 10.21037/exrna-22-22

Cite this article as: Gumas J, Kirino Y. Extracellular tRNA-derived RNAs as emerging activators of endosomal Toll-like receptors: a narrative review. *ExRNA* 2023;5:2.