

5-2019

Letter to the Editor: Ketamine for pain management: Let's not neglect practical concerns

Steven P. Cohen

Johns Hopkins School of Medicine; Uniformed Services University of the Health Sciences

Eric S. Schwenk

Thomas Jefferson University

Eugene R. Viscusi

Thomas Jefferson University

Follow this and additional works at: <https://jdc.jefferson.edu/anfp>



Part of the [Anesthesiology Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Cohen, Steven P.; Schwenk, Eric S.; and Viscusi, Eugene R., "Letter to the Editor: Ketamine for pain management: Let's not neglect practical concerns" (2019). *Department of Anesthesiology Faculty Papers*. Paper 51.

<https://jdc.jefferson.edu/anfp/51>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Anesthesiology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.



Letter to the Editor: Ketamine for pain management: let's not neglect practical concerns

Steven P. Cohen^{a,b}, Eric S. Schwenk^c, Eugene R. Viscusi^c, on behalf of the American Society of Regional Anesthesia & Pain Medicine and American Academy of Pain Medicine Ketamine Guidelines Committees

We read with interest the recent article on ketamine by Bell and Kalso.¹ In recent years, the use of ketamine to treat chronic pain and depression has exploded, and this article represents an easy-to-follow overview of this often misunderstood medication.

In their article, the authors focus on systematic reviews to draw broad conclusions on acute and chronic pain, without delving into the specifics of studies, including selection criteria and limitations. Although this methodology is sufficient for a topical review that summarizes basic information to include mainstream generalizations about efficacy and side effects, it is not capable of providing clinicians with practical recommendations that can be used to guide care such as indications and contraindications, dose regimens, benefit as an adjunct to opioids for acute pain, evidence for follow-up therapy with oral ketamine and other *N*-methyl-D-aspartate antagonists, how often and in whom to repeat procedures, how to prevent and treat adverse effects, and practical concerns such as how and by whom these patients should be monitored.

For both acute and chronic pain, this information and more is provided in recent guidelines along with a narrative review developed in a joint effort by the American Society of Regional Anesthesia & Pain Medicine, the American Academy of Pain Medicine, and the American Society of Anesthesiologists.^{2,3} Considering that anesthesiologists (not to mention other clinicians) comprise a plurality of International Association for the Study of Pain membership, these pragmatic concerns have clinical relevance for the Journal's readership. We assume the time frame of this publication, which was revised and accepted in June 2018, prevented the inclusion of these guidelines, but would encourage readers interested in the practical elements of intravenous ketamine use for pain treatment to explore these resources.

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

^a Departments of Anesthesiology and Critical Care Medicine, Neurology and Physical Medicine & Rehabilitation, Johns Hopkins School of Medicine, Baltimore, MD, ^b Departments of Anesthesiology and Physical Medicine & Rehabilitation, Uniformed Services University of the Health Sciences, Bethesda, MD, ^c Department of Anesthesiology, Jefferson Medical College, Philadelphia, PA

Copyright © 2019 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of The International Association for the Study of Pain. This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

PR9 4 (2019) e728

<http://dx.doi.org/10.1097/PR9.0000000000000728>

Disclosures

The authors have no conflict of interest to declare.

The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense.

Acknowledgements

All authors served on the ASRA, AAPM, and ASA acute and chronic pain management ketamine guidelines committee, and Dr Schwenk is preparing an NIH grant evaluating ketamine for submission, but they report no conflicts related to this letter.

Article history:

Received 15 January 2019

Accepted 30 January 2019

References

- [1] Bell RF, Kalso EA. Ketamine for pain management. *PAIN Rep* 2018;3:e674.
- [2] Cohen SP, Bhatia A, Buvanendran A, Schwenk ES, Wasan AD, Hurley RH, Viscusi ER, Narouze S, Davis FN, Ritchie EC, Lubenow TR, Hooten WM. Consensus guidelines on the use of intravenous ketamine infusions for chronic pain from the American Society of Regional Anesthesia and Pain Medicine (ASRA), American Academy of Pain Medicine (AAPM) and the American Society of Anesthesiologists (ASA). *Reg Anesth Pain Med* 2018; 43:521–46.
- [3] Schwenk E, Viscusi ER, Buvanendran A, Hurley RH, Wasan AD, Narouze S, Bhatia A, Davis FN, Hooten WM, Cohen SP. Consensus guidelines on the use of intravenous ketamine infusions for acute pain management from the American Society of Regional Anesthesia and Pain Medicine (ASRA), American Academy of Pain Medicine (AAPM) and the American Society of Anesthesiologists (ASA). *Reg Anesth Pain Med* 2018;43:456–66.