

12-4-2014

Toward an improved understanding of the ionizing radiation induced DNA damage/response networks in human malignancies.

Daphne A Haas-Kogan

Departments of Radiation Oncology and Neurological Surgery, Helen Diller Family Comprehensive Cancer Center, UCSF Benioff Children's Hospital, University of California San Francisco

David R Raleigh

Department of Radiation Oncology, University of California San Francisco

Adam Paul Dicker

Department of Radiation Oncology, Kimmel Cancer Center, Jefferson Medical College, Thomas Jefferson University

Follow this and additional works at: <https://jdc.jefferson.edu/radoncfp>



Part of the [Radiology Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Haas-Kogan, Daphne A; Raleigh, David R; and Dicker, Adam Paul, "Toward an improved understanding of the ionizing radiation induced DNA damage/response networks in human malignancies." (2014). *Department of Radiation Oncology Faculty Papers*. Paper 50.
<https://jdc.jefferson.edu/radoncfp/50>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Radiation Oncology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.



Toward an improved understanding of the ionizing radiation induced DNA damage/response networks in human malignancies

Daphne A. Haas-Kogan^{1*†}, David R. Raleigh^{2†} and Adam Paul Dicker^{3*†}

¹ Departments of Radiation Oncology and Neurological Surgery, Helen Diller Family Comprehensive Cancer Center, UCSF Benioff Children's Hospital, University of California San Francisco, San Francisco, CA, USA

² Department of Radiation Oncology, University of California San Francisco, San Francisco, CA, USA

³ Department of Radiation Oncology, Kimmel Cancer Center, Jefferson Medical College, Thomas Jefferson University, Philadelphia, PA, USA

*Correspondence: dhaaskogan@radonc.ucsf.edu; Adam.Dicker@jefferson.edu

†All authors contributed equally to this Editorial.

Edited and reviewed by:

Timothy James Kinsella, Warren Alpert Medical School of Brown University, USA

Keywords: ionizing radiation, ionizing radiation induced DNA damage, DNA damage, radiation, editorial

GLIOBLASTOMA EGFR MUTATION PREDICTS THE RESPONSE TO CYTOTOXIC AND TARGETED THERAPIES

Several molecular aberrations in glioblastoma have been shown to have prognostic significance, and investigations of targeted therapies for glioma patients are ongoing. Here, Wachsberger and colleagues demonstrate that overexpression of a constitutively active form of the epidermal growth factor receptor (EGFR) found in some glioblastomas sensitizes cells to both multimodal cytotoxic therapy and cetuximab (1). The data further indicate that EGFR mutation may predispose gliomas to defective homologous recombination repair (HRR). These findings are relevant to future clinical trials, and suggest that targeted agents may achieve the greatest benefit in glioma patients who have been stratified according to molecular markers.

MicroRNA MEDIATES THE CELLULAR RESPONSE TO DNA DAMAGE

DNA damage triggers a signaling cascade that affects expression, subcellular localization, and activity of mediators; the strength and nature of molecular interactions; and chromatin structure. Ongoing discovery of novel effectors has amplified understanding of the complex network that mediates the DNA damage response (DDR). Recent evidence demonstrates that genotoxic stress regulates expression of distinct microRNAs (miRNA), some of which have been shown to directly influence expression of DDR effectors including ATM, DNA-PKcs, and BRCA1. The links between miRNAs and the DDR are summarized here by Wright and colleagues, as are the implications of future investigation into this emerging field (2).

MODULATION OF DOUBLE-STRAND BREAK REPAIR PATHWAYS TO RADIOSENSITIZE CANCER CELLS

Disparate sensitivity to the cytotoxic effects of radiation mediates selective tumor cell killing with relative preservation of normal tissues. Significant effort has been dedicated to understanding the mechanisms underlying differential radiosensitivity with the ultimate goal of enhancing the therapeutic ratio of ionizing radiation. Homologous recombinational repair is among the primary

pathways utilized by cells to process otherwise lethal DNA damage, and is therefore a promising target to improve the efficacy of radiation therapy. The major pathways of double-strand break repair are summarized here by Mladenov and colleagues, as are potential targets for radiosensitizing cancer cells (3).

RADIATION-INDUCED SECOND NEOPLASMS

Every year, more than 700 childhood cancer survivors will join the 300,000 individuals in the United States who have survived their cancer diagnoses, but will struggle with the consequences of the treatments that cured them. Twenty-five years after the first cancer diagnosis, the death rate due to subsequent malignancies exceeds that due to all other causes. Technological advances have transformed the delivery of radiation, resulting in more conformal radiation plans. We must investigate whether the benefit of decreasing acute side effects comes at a cost of lower-dose radiation distributed throughout the child's body that may increase the risk of fatal radiation-induced cancers. Braunstein and colleagues highlight such investigations into the current state of epidemiologic modeling and radiotherapy delivery data and their impacts on patient care (4).

THE COMBINATION OF NOVEL TARGETED MOLECULAR AGENTS AND RADIATION IN THE TREATMENT OF PEDIATRIC GLIOMAS

A comprehensive, cutting edge approach to gliomas requires first a critical understanding of the molecular underpinnings of glioma pathogenesis and response to therapy. Advances in molecular, genetic, and biochemical technologies have provided unprecedented insight into glioma biology and these are reviewed herein by Dasgupta and colleagues (5). Targeted therapeutics are available for many implicated pathways; however, follow up biological studies defining the appropriate, most effective agents for each alteration are lacking. Clinical studies must be designed that collect and harness molecular data within prospective trials, in order to generate pre-clinical and clinical data that will determine the best combination of agents for each molecular subtype.

IONIZING RADIATION IN GLIOBLASTOMA INITIATING CELLS

Rivera and colleagues explore the mechanisms underlying radiation resistance in GBM initiating cells (GICs) that are critical to enhanced therapeutic efficacy against neural progenitor cells (NPCs) (6). The importance of this approach is highlighted by several recent reports of improved glioma control and clinical outcome when aggressive therapy and higher doses of radiation are administered to the adult human subventricular zone, which constitutes the largest area of neurogenesis and houses the greatest concentration of NPCs. Such approaches are now possible with our modern arsenal of sophisticated dose delivery technology, with which we can modulate the dose to substructures of the brain.

HYPOFRACTIONATED RADIATION AND MOUSE MODELS OF NON-SMALL CELL LUNG CANCER

Recent clinical trials have demonstrated high rates of long-term local control and survival from early stage non-small cell lung cancer (NSCLC) treated with stereotactic body radiotherapy (SBRT). Compared to conventionally fractionated radiation regimens, the success of SBRT for NSCLC suggests that hypofractionated irradiation may induce distinct molecular pathways to enhance cell killing. Using two genetically engineered mouse models of NSCLC, Perez and colleagues demonstrate superior tumor control with hypofractionated radiation. The cellular response to SBRT is incompletely understood, but the genetic reagents described here will be useful to elucidating the molecular mechanisms that underlie the SBRT DDR (7).

REFERENCES

1. Wachsberger PR, Lawrence RY, Liu Y, Rice B, Daskalakis C, Dicker AP. Epidermal growth factor receptor mutation status and rad51 determine the response of glioblastoma to multimodality therapy with cetuximab, temozolomide, and radiation. *Front Oncol* (2013) 3:13. doi:10.3389/fonc.2013.00013
2. Wright CM, Dan T, Dicker AP, Simone NL. microRNAs: the short link between cancer and RT-induced DNA damage response. *Front Oncol* (2014) 4:133. doi:10.3389/fonc.2014.00133
3. Mladenov E, Magin S, Soni A, Iliakis G. DNA double-strand break repair as determinant of cellular radiosensitivity to killing and target in radiation therapy. *Front Oncol* (2013) 3:113. doi:10.3389/fonc.2013.00113
4. Braunstein S, Nakamura JL. Radiotherapy-induced malignancies: review of clinical features, pathobiology, and evolving approaches for mitigating risk. *Front Oncol* (2013) 3:73. doi:10.3389/fonc.2013.00073
5. Dasgupta T, Haas-Kogan DA. The combination of novel targeted molecular agents and radiation in the treatment of pediatric gliomas. *Front Oncol* (2013) 3:110. doi:10.3389/fonc.2013.00110
6. Rivera M, Sukhdeo K, Yu J. Ionizing radiation in glioblastoma initiating cells. *Front Oncol* (2013) 3:74. doi:10.3389/fonc.2013.00074
7. Perez BA, Ghafoori AP, Lee CL, Johnston SM, Li Y, Moroshek JG, et al. Assessing the radiation response of lung cancer with different gene mutations using genetically engineered mice. *Front Oncol* (2013) 3:72. doi:10.3389/fonc.2013.00072

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 31 October 2014; accepted: 05 November 2014; published online: 04 December 2014.

Citation: Haas-Kogan DA, Raleigh DR and Dicker AP (2014) Toward an improved understanding of the ionizing radiation induced DNA damage/response networks in human malignancies. *Front. Oncol.* 4:335. doi: 10.3389/fonc.2014.00335

This article was submitted to *Radiation Oncology*, a section of the journal *Frontiers in Oncology*.

Copyright © 2014 Haas-Kogan, Raleigh and Dicker. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.