9-16-2019

Continuation of angiotensin converting enzyme inhibitors on the day of surgery is not associated with increased risk of hypotension upon induction of general anesthesia in elective non-cardiac surgeries.

U. Yoon

Thomas Jefferson University

A. J. Setren

Thomas Jefferson University

A. Chen

Thomas Jefferson University

Follow this and additional works at: <https://jdc.jefferson.edu/anfp>

T. Nguyen

[Thomas Jefferson University Commons](#)

[Let us know how access to this document benefits you](#)

M. C. Torjman

Thomas Jefferson University

Recommended Citation

Yoon, U.; Setren, A. J.; Chen, A.; Nguyen, T.; Torjman, M. C.; and Kennedy, T. L., "Continuation of angiotensin converting enzyme inhibitors on the day of surgery is not associated with increased risk of hypotension upon induction of general anesthesia in elective non-cardiac surgeries." (2019). *Department of Anesthesiology Faculty Papers*. Paper 53.
<https://jdc.jefferson.edu/anfp/53>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Anesthesiology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

U. Yoon, A. J. Setren, A. Chen, T. Nguyen, M. C. Torjman, and T. L. Kennedy

Continuation of Angiotensin Converting Enzyme Inhibitors on the Day of Surgery Is Not Associated with Increased Risk of Hypotension Upon Induction of General Anesthesia in elective Non-Cardiac Surgeries

Yoon U, MD¹, Setren AJ, MD¹, Chen A, BS¹, Nguyen T, BA¹, Torjman MC, PhD¹, Kennedy TL, MD¹,

¹ Department of Anesthesiology, Thomas Jefferson University Hospital, Philadelphia, PA

Corresponding Author:

Uzung Yoon, MD, MPH

Department of Anesthesiology

Thomas Jefferson University Hospital,

Suite 8290 Gibbon, 111 South 11th Street, Philadelphia, PA 19107

Phone: (215) 955-2370

Fax: (215) 955-0677

Email: uzyoon@gmail.com

Declarations of interest: none

Financial disclosure: none

Brief title: angiotensin converting enzyme inhibitors and hypotension

Keywords: angiotensin converting enzyme inhibitors, angiotensin II receptor blockers, induction, intraoperative hypotension

Dear Editor

Angiotensin-converting-enzyme inhibitors (ACE-I) and angiotensin-II receptor blockers (ARB) have become progressively popular to treat hypertension due to their beneficial effect in reducing morbidity and mortality. At the same time the rate of major noncardiac surgical procedures is increasing and the use of these agents may pose challenges in the perioperative setting. Intraoperative hypotension in the setting of ACE-I and ARBs use has been reported in the literature[1]. However, the effect on ACE-I and ARB on induction related hypotension has not been studied. Induction related hypotension almost always occurs, is immediate and plateaus soon after due to increased volume loading, vasopressor use and cardiovascular adaptation[2]. The question was raised whether ACE-I or ARB continuation increases the risk of induction related intraoperative hypotension within 15 minutes after general anesthesia in elective non-cardiac surgeries.

WE conducted a retrospective-cohort-study among elective non-cardiac surgery patients. The study was approved by the institutional review board and informed consent was waived. 400 electronic medical records with patients who had an ACE-I or ARB on their list of preoperative home medications were screened. Patients were divided into two groups: continuation of ACE/ARB and discontinuation of ACE/ARB. Patient who received regional anesthesia were excluded. Hypotension was defined as either: systolic-BP <90 mm Hg or systolic-BP decrease of >20%.

349 patients were included for final analysis. The mean admission ASA-status was 2.7 ± 0.5 , age 65 ± 11 years, and BMI 31 ± 6.9 kg/m². There were no statistically significant changes between the ACE-I/ARB continuation and discontinuation group in terms of SBP ($p=0.853$), DBP ($p=0.357$), MAP ($p=0.782$) and HR ($p=0.220$), figure 1. There were also no significant differences in induction medication dose (propofol, fentanyl and rocuronium) and pressor use ($p=0.137$). Hypotension ($p<0.001$) occurred equally in both groups over 15 minutes.

Hollman et al[3] meta-analysis showed that continuation of ACE-I/ARBs was associated with increased risk of intraoperative hypotension up to 60 minutes after induction. However, most included studies had a low sample size, pertain to a specific type of surgery such as vascular, cardiac, or bariatric, and reported only small incidences of hypotension. The heterogeneity was very high ($I^2=71\%$) and the results may not be generalizable or reproducible in a larger patient population. Furthermore, no study looked specifically for the effect on induction related hypotension. Despite intraoperative hypotension was reported, the meta-analysis was unable to demonstrate an association between perioperative ACE-I/ARB administration and mortality, MACE, CHF, AKI, or CVA.

The decision whether to continue or withhold ACE-I/ARB preoperatively in elective non-cardiac surgeries is still controversial. Currently, the 2014 American College of Cardiology/American Heart Association guidelines[4] state that it is reasonable to continue therapy preoperatively, and if withheld, therapy may be reinstituted as soon as clinically feasible.

The European Society of Cardiology/European Society of Anaesthesiology [5] bases its recommendations on the indication for treatment with an ACE-I/ARB, recommending discontinuation for 24-hours before surgery if prescribed for hypertension, and continuation if prescribed for heart failure and left ventricular systolic dysfunction.

In conclusion the continuation of ACE-I/ARB on the day of surgery was not associated with increased risk of intraoperative hypotension upon induction and within 15 minutes of general anesthesia in elective non-cardiac surgeries. However, the risk of intraoperative hypotension even after the induction phase with continuation of ACE/ARB still remains controversial, and the current evidence has failed to correlate the impact of intraoperative hypotension with adverse patient outcome. Our study adds another piece of data into the discussion and we suggest that individual perioperative risk factors should be considered before making the decision whether to continue or withhold ACE-I/ARB. Discontinuation may be considered for patients at high risk for hypotension-related complications, including patients with critical cardiovascular or cerebrovascular disease.

Reference

1. Roshanov PS, Rochwerg B, Patel A, et al. Withholding versus Continuing Angiotensin-converting Enzyme Inhibitors or Angiotensin II Receptor Blockers before Noncardiac Surgery: An Analysis of the Vascular events In noncardiac Surgery patients cOhort evaluationN Prospective Cohort. *Anesthesiology*. 2017;126:16-27.
2. Pigott DW, Nagle C, Allman K, Westaby S, Evans RD. Effect of omitting regular ACE inhibitor medication before cardiac surgery on haemodynamic variables and vasoactive drug requirements. *Br J Anaesth* 1999;83:715–20
3. Hollmann C, Fernandes NL, Biccard BM. A systematic review of outcomes associated with withholding or continuing angiotensin-converting enzyme inhibitors and angiotensin receptor blockers before noncardiac surgery. *Anesth Analg*. 2018;127:678–687.
4. Fleisher LA, Fleischmann KE, Auerbach AD, et al.; American College of Cardiology; American Heart Association. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *J Am Coll Cardiol*. 2014;64:e77–e137
5. Kristensen SD, Knuuti J, Saraste A, et al.; Authors/Task Force Members. 2014 ESC/ESA guidelines on non-cardiac surgery: cardiovascular assessment and management: the joint task force on non-cardiac surgery: cardiovascular assessment and management of the European Society of Cardiology (ESC) and the European Society of Anaesthesiology (ESA). *Eur Heart J*. 2014;35:2383–2431.

Figure 1. Mean $\pm$ SEM Mean Arterial Pressure (MAP) measured pre-induction and at six additional time points for 15 minutes post induction