6-1-2010

Prevalence of obesity, type II diabetes mellitus, hyperlipidemia, and hypertension in the United States: findings from the GE Centricity Electronic Medical Record database.

Albert G Crawford

Thomas Jefferson University School of Population Health

Christine Cote

Johnson and Johnson, Corporate Office of Science and Technology

Joseph Couto

Thomas Jefferson University School of Population Health

Follow this and additional works at: <https://jdc.jefferson.edu/healthpolicyfaculty>

Mehmet Daskiran
Part of the [Community Health and Preventive Medicine Commons](#), and the [Health Services Research Commons](#)
Johnson and Johnson, Corporate Office of Science and Technology

Candace Gunnarsson

S2Statistical Solutions, Inc.

Let us know how access to this document benefits you

Recommended Citation

~~See next page for additional authors.~~

Crawford, Albert G; Cote, Christine; Couto, Joseph; Daskiran, Mehmet; Gunnarsson, Candace; Haas, Kara; Haas, Sara; Nigam, Somesh C; and Schuette, Rob, "Prevalence of obesity, type II diabetes mellitus, hyperlipidemia, and hypertension in the United States: findings from the GE Centricity Electronic Medical Record database." (2010). *College of Population Health Faculty Papers*. Paper 54.

<https://jdc.jefferson.edu/healthpolicyfaculty/54>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in College of Population Health Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Albert G Crawford, Christine Cote, Joseph Couto, Mehmet Daskiran, Candace Gunnarsson, Kara Haas, Sara Haas, Somesh C Nigam, and Rob Schuette

Prevalence of Obesity, Type II Diabetes Mellitus, Hyperlipidemia, and Hypertension in the United States: Findings from the GE Centricity Electronic Medical Record Database

Albert G. Crawford, PhD,¹ Christine Cote, MD,² Joseph Couto, PharmD,¹ Mehmet Daskiran, PhD,²
Candace Gunnarsson, PhD,³ Kara Haas, MD,⁴ Sara Haas, MS,³
Somesh C. Nigam, PhD,⁵ and Rob Schuette, PhD³

Abstract

This study analyzed GE Centricity Electronic Medical Record (EMR) data to examine the effects of body mass index (BMI) and obesity, key risk factor components of metabolic syndrome, on the prevalence of 3 chronic diseases: type II diabetes mellitus, hyperlipidemia, and hypertension. These chronic diseases occur with high prevalence and impose high disease burdens. The rationale for using Centricity EMR data is 2-fold. First, EMRs may be a good source of BMI/obesity data, which are often underreported in surveys and administrative databases. Second, EMRs provide an ideal means to track variables over time and, thus, allow longitudinal analyses of relationships between risk factors and disease prevalence and progression. Analysis of Centricity EMR data showed associations of age, sex, race/ethnicity, and BMI with diagnosed prevalence of the 3 conditions. Results include uniform direct correlations between age and BMI and prevalence of each disease; uniformly greater disease prevalence for males than females; varying differences by race/ethnicity (ie, African Americans have the highest prevalence of diagnosed type II diabetes and hypertension, while whites have the highest prevalence of diagnosed hypertension); and adverse effects of comorbidities. The direct associations between BMI and disease prevalence are consistent for males and females and across all racial/ethnic groups. The results reported herein contribute to the growing literature about the adverse effects of obesity on chronic disease prevalence and about the potential value of EMR data to elucidate trends in disease prevalence and facilitate longitudinal analyses. (*Population Health Management* 2010;13:151–161)

Introduction

THERE IS A VAST LITERATURE ON THE ETIOLOGY of type II diabetes mellitus, hyperlipidemia, and hypertension. A key concept in understanding the origin and course of these chronic conditions is metabolic syndrome, which refers to a cluster of risk factors for cardiovascular disease (CVD). These risk factors were originally described by Reaven as a composite of obesity, dyslipidemia, hypertension, and hyperglycemia. The exact measures and thresholds for these components have undergone several iterations in the past 2 decades.¹ Currently, the most frequently used definitions are those developed by (1) the World Health Organization in

1999,² with important later revisions by the European Group for the Study of Insulin Resistance in 1999³; (2) the National Cholesterol Education Program Adult Treatment Panel III in 2001⁴; and (3) the International Diabetes Federation (IDF) in 2006.⁵ It remains arguable whether the composite metabolic syndrome categorization is better than the sum of its parts at predicting CVD, as well as whether the latest definitional iteration (IDF) lends additional predictive power to older definitions.^{6,7} One obvious advantage of the IDF definition, however, may be the recent recognition that one measure of obesity may not fit all ethnicities; hence, the IDF definition of metabolic syndrome may be a more globally useful definition.

¹Thomas Jefferson University School of Population Health, Philadelphia, Pennsylvania.
Johnson & Johnson, ²Corporate Office of Science and Technology, and ³Office of Evidence Based Medicine, Medical Device and Diagnostics New Brunswick, New Jersey.

³Statistical Solutions, Inc., Cincinnati, Ohio.

⁴Ethicon Endo-Surgery, Inc., New Brunswick, New Jersey.

The various definitions and indicators noted have been used to assess the prevalence of metabolic syndrome and the clinical outcomes of patients in several large observational and interventional cohorts.^{8–13} Among the more prominent data sets are those of the Framingham Study, including the Framingham Offspring Study, in the United States, and the General Practice Research Database in the United Kingdom, the largest non-US database of “anonymized” primary care medical records.^{14,15}

Turning to the effects of body mass index (BMI) and obesity, Bays and colleagues¹⁶ explored the relationships between BMI and prevalence of diabetes, hypertension, and dyslipidemia, examined BMI distributions among patients with these conditions, and compared results from 2 national surveys: (1) the screening questionnaire-based 2004 Study to Help Improve Early evaluation and management of risk factors Leading to Diabetes (SHIELD), and (2) the interview, clinical, and laboratory data of the 1999–2002 National Health and Nutrition Examination Surveys (NHANES). Bays et al found that mean BMI was virtually identical in the 2 data sets: 27.8 kg/m² for SHIELD and 27.9 kg/m² for NHANES. Increased BMI was associated with increased prevalence of diabetes mellitus, hypertension, and dyslipidemia in both studies ($P < 0.001$). For each condition, more than 75% of patients had a BMI > 25 kg/m². Estimated prevalence of diabetes mellitus and hypertension was similar in both studies, while prevalence of dyslipidemia was substantially higher in NHANES than SHIELD.

McTigue et al¹⁷ studied 90,185 women who participated in the Women’s Health Initiative to determine how cardiovascular and mortality risks differ across clinical weight categories in women, especially the extreme obesity category. The duration of follow-up averaged 7.0 years (1993–2004). The researchers found that, among women, extreme obesity prevalence differed with race/ethnicity, from 1% among Asians and Pacific Islanders to 10% among black women. All-cause mortality rates per 10,000 person-years were 68.49 (95% confidence interval [CI], 65.26–71.68) for normal BMI (< 25), 71.16 (95% CI 67.68–74.82) for overweight ($25 \leq \text{BMI} < 30$), 84.47 (95% CI 78.90–90.42) for obesity 1 ($30 \leq \text{BMI} < 35$), 102.85 (95% CI 92.90–113.86) for obesity 2 ($35 \leq \text{BMI} < 40$), and 116.85 (95% CI 103.36–132.11) for extreme obesity ($\text{BMI} \geq 40$). Analyses adjusting for the effects of age, smoking, educational achievement, region of the United States, and physical activity level showed that weight-related all-cause mortality risk, coronary heart disease (CHD) mortality, and CHD incidence did not differ by race/ethnicity. Such adjusted analyses among both white and black participants showed positive trends in all-cause mortality and CHD incidence with increasing weight. Much of the obesity-related mortality and CHD risk was mediated by diabetes, hypertension, and hyperlipidemia. For white women, weight-related all-cause mortality risk varied by age, with obesity producing less risk among older women. McTigue and colleagues advocate using more weight categories and conclude that “considering obesity as a body mass index of 30 or higher may lead to misinterpretation of individual and population risks.”

Sturm¹⁸ asserts that “the most serious health problems are not associated with overweight or moderate obesity ... but

with clinically severe or morbid obesity (eg, more than 100 pounds (45 kg) overweight).” He uses data from the Behavioral Risk Factor Surveillance System to estimate trends for extreme weight categories ($\text{BMI} > 40$ and $\text{BMI} > 50$) for the period from 1986–2005 in the United States and assesses whether trends have changed since 2000. Sturm finds that from 2000–2005, while the prevalence of obesity (self-reported $\text{BMI} > 30$) increased by 24%, the prevalence of $\text{BMI} > 40$ increased by 50% and the prevalence of $\text{BMI} > 50$ increased by 75%. Sturm concludes that “the heaviest BMI groups have been increasing at the fastest rates for 20 years” and that “because comorbidities and resulting service use are much higher among severely obese individuals, the widely published trends for overweight/obesity underestimate the consequences for population health.”

The analyses reported herein employ GE Centricity^a electronic medical record (EMR) data to assess the prevalence of the 3 chronic conditions and the role of obesity in their etiology and progression. The GE database comprises the “anonymized” longitudinal medical records of over 6 million patients, is considered one of the few high-quality patient databases available commercially, and is generally representative of the population of the United States.

A growing proportion of ambulatory care physician practices in the United States are adopting EMR systems to support various clinical processes including documentation of patient encounters and secure exchange of data with other providers. In 2007, the GE Centricity EMR was used by more than 20,000 clinicians in 49 states for medical record documentation for approximately 30 million patients. These electronic records contain a wide range of demographic and clinical variables. The structured user interface supports consistent and accurate documentation by physicians and thus promotes internal data validity for each patient and for the aggregate. As with other EMRs, a significant attribute of Centricity is its capacity to track patients’ clinical conditions over time and, thus, to support longitudinal multivariate analytic designs, including retrospective cohort studies. Cumulatively, these attributes of Centricity and other EMRs make them rich resources for population health and outcomes research.

Research interest in the GE Centricity database is evident in the growing peer-reviewed literature employing this resource. Our review of recently published peer-reviewed literature employing GE Centricity EMR data revealed 2 especially useful and relevant studies. First, Brixner et al¹⁹ used clinical, diagnostic, and treatment/prescription information in the Centricity database to examine the prevalence of cardiometabolic risk (CMR) factors that contribute to metabolic syndrome in the primary care setting. The authors concluded that “... the distribution of CMR factors in a primary care database is similar to that established by prospective national health surveys such as NAMCS. A key method for identifying risk factors is using clinical outcomes, including BMI and lab values. Future studies on metabolic syndrome need to link clinically based information with more readily available treatment and diagnosis information.” The second noteworthy study employing Centricity data is Gill and Chen’s 2008 evaluation of lipid management (ie,

^aCentricity Physician Office is a registered trademark of GE Medical Systems Information Technologies.

adequacy of lipid testing, achievement of lipid goals, appropriate use of lipid-lowering medication).²⁰ The authors note that "National EHR networks are excellent vehicles for large outpatient quality of care studies, particularly for measuring clinical outcomes such as lipid levels." These studies demonstrate that national EMR databases such as Centricity are valuable tools for epidemiological, outcomes, and health services research.

Methods

GE Centricity Database

The GE Centricity database captures patient-level clinical data elements obtained from the Centricity Physician Office EMR (formerly Logician) for Clinical Data Services (CDS) reporting. The Centricity EMR and its predecessors have been in use for over 20 years, are certified by the Certification Commission for Healthcare Information Technology, and are currently used by over 30,000 clinicians throughout the United States. The CDS database contains data from 133 provider groups with 7259 clinicians (including approximately 60% primary care providers and 40% specialists) at 98 installations. The CDS database includes de-identified, standardized data on more than 8.9 million patients, with the median duration of documentation being 985 days, or approximately 2.7 years. Among the strengths of the Centricity database are its incorporation of documentation of a wide range of diagnostic and therapeutic services, specifically laboratory test results (with exact amounts and units of measurement) and medications ordered, and, perhaps most important, inclusion of the above-mentioned data at multiple points in time, allowing longitudinal analyses not possible with most other data sets.

While the CDS database includes a large and diverse group of providers, participation is voluntary, raising concerns regarding self-selection by providers. For example, Gill and Chen suggest that "practices that are more interested in measuring (and improving) quality of care are more likely to join MQIC [Medical Quality Improvement Consortium]."²⁰ Moreover, a recent study by Oderda et al²¹ that compared the patient population in MQIC to the general US population found that the MQIC population is older (with approximately 50% age 45 years or older, compared to 37% of the US population), more likely to be from the northeast or midwest regions (62% vs. 41%), and more likely to have commercial insurance (73% vs. 59%) or Medicare (20% vs. 11%). However, the MQIC population is very similar in its racial distribution (79% vs. 81% white, 15% vs. 13% African American). Although the proportion of patients with hypertension in the MQIC database is virtually identical to that of the US population (25.9% vs. 25.8%), according to data from the National Health Interview Survey, the proportion with diabetes is somewhat higher (9.8% vs. 7.4%). To summarize, as Gill and Chen note, "while the MQIC population differs somewhat from the general US population, it is unclear whether this reflects differences in persons who seek care in outpatient settings, differences in providers who use an EHR [electronic health record], or differences in providers who use an EHR and join MQIC."²⁰

Among the specific potential limitations of the Centricity database are (1) invalid retrospective patient data, (2) unconfirmed diagnoses, and (3) duplicate patient data. The present research design addresses these 3 potential limita-

tions by applying inclusion/exclusion criteria to maximize the validity of the data.

Inclusion/exclusion criteria

Exclusion of invalid retrospective patient data. When a new provider's data are entered into the Clinical Data Services database, the visit dates may be back-filled, and thus may be inaccurate (ie, the database entry date is assigned as the visit date for activities that occurred long before that date). As a result of the potential inaccuracy of visit dates, the duration and prevalence of chronic conditions may be underestimated. Therefore, the research design used herein excludes patient data for the first year after the patient data were initially entered into the database.

Exclusion of unconfirmed diagnoses. Patients merely evaluated for a condition and patients with confirmed diagnoses have identical *International Classification of Diseases, Ninth Revision* (ICD-9) codes in the database. Exclusion criteria were developed to exclude instances when patients were merely evaluated for a diagnosis (with no confirmation of the evaluated diagnosis) through the following algorithm: (a) include only records where the Diagnosis Qualifier contains any of the phrases "diagnosis of," "hospitalized for," "history of" (but not "family history of"), "minor diagnosis of," "take note of," "recurrence of," or "status post"; and (b) exclude records where the Problem Concept description contains any of the phrases "rule out," "family history of," "risk of," or "screening."

Further investigation may reveal that disease counts should be supplemented with data derived from sources in the database other than problems and explicit ICD-9 diagnosis codes. For instance, for diabetes, there may be a medication record indicating regular use of insulin or oral hypoglycemics for a patient with no ICD-9 diagnosis code for diabetes in the database. Likewise, for hypertension, there may be recorded observations of blood pressure for a patient who may not have an ICD-9 diagnosis code for hypertension in the database.

Exclusion of duplicate patient data. An exclusion algorithm was applied to exclude cases for which there was potential duplication of patient data as a result of a patient visiting more than 1 Centricity client practice. This is critical because the data are "anonymized" (ie, all unique patient identifiers are removed in the de-identification process). To avoid duplication of patients, the research design specified exclusion of data originating from 3-digit zip codes containing 3 or more providers who employ the Centricity EMR system. The rationale for this exclusion criterion is that a patient residing in a 3-digit zip code area that contains more than 1 practice employing Centricity may appear in the database as more than 1 patient.

In addition to applying the aforementioned inclusion/exclusion algorithms, additional data (ie, diagnosis and procedure codes, patient age) were obtained from the Centricity EMR database. Current Procedural Terminology (CPT) code numbers indicating specific procedures can be found in either the Order table or the Problem table, requiring that both be mined to identify procedures. To identify each procedure, our design scanned all records in both of

FIG. 1. Attrition flow diagram.

these tables to identify relevant CPT procedure codes. Likewise, age was derived from either the "Activity Fact" table or the "Patient Dimension" table, with preference given to the former table. A summary of the results of employing these inclusion/exclusion algorithms is shown in Figure 1.

Analytic methods

Patient age groups used in the bivariate analyses are <18 years, 18–24, 25–34, 35–44, 45–54, 55–59, 60–64, 65–69, and ≥70. For simplicity, the age groups used in the multivariate analyses are <20, 20–39, 40–59, and ≥60 years. The narrower age ranges were selected so as to discern subtle patterns as well as general ones. BMI categories are BMI < 18.5, $18.5 \leq \text{BMI} < 25$, $25 \leq \text{BMI} < 30$, $30 \leq \text{BMI} < 35$, $35 \leq \text{BMI} < 40$, $40 \leq \text{BMI} < 45$, $45 \leq \text{BMI} < 50$, and BMI ≥ 50; these categories were likewise selected to be sufficiently narrow to detect subtle patterns. Racial/ethnic categories include white, African American, Hispanic, Oriental/Asian, Native American, Other, and Unknown.

In the initial analyses, contingency tables were used to assess the bivariate relationships of BMI category, age group, sex, and race/ethnicity with the diagnosed prevalence of

type II diabetes mellitus, hyperlipidemia, and hypertension. Sets of 3-way tables stratified the relationships between BMI category and diagnosed prevalence of the 3 diseases as a function of age group, sex, and race/ethnicity. Next, a contingency table was used to assess the relationship between BMI category and the diagnosis of obesity per se. Finally, logistic regressions were performed to summarize the effects of demographic factors (age, sex, and race/ethnicity), obesity, and comorbidities on the prevalence of each of the 3 conditions. These analyses consisted of an initial step including only the demographic factors and obesity, and then a second step adding the relevant comorbidities.

Results

Figures 2–5 and Table 1 present GE Centricity EMR results for the number and percent distribution of patients with diagnosed type II diabetes, hyperlipidemia, and hypertension, as a function of patient BMI, age group, race/ethnicity, and sex, for the United States in 2005. The GE Centricity data show that the prevalence of each condition is positively associated with both BMI and age (see Figs. 2 and 3). Figure 4 shows the associations between race/ethnicity and disease prevalence, and indicates that African Americans have the highest proportions of patients with diagnosed diabetes (12.1%) and hypertension (28.9%), while whites have the highest proportion with diagnosed hyperlipidemia (25.7%). The associations between sex and disease prevalence are shown in Figure 5, which portrays how males are more likely than females to have each of the 3 conditions: 9.3% vs. 7.2% have diabetes, 25.1% vs. 19.2% have hyperlipidemia, and 22.8% vs. 20.1% have hypertension.

As shown in Figures 2 and 3 and Table 2, the proportions of patients with type II diabetes mellitus, hyperlipidemia, and hypertension increase as both BMI and age increase. Figures 6A, 6B, and 6C show the combined effects of BMI and age on the prevalence of diabetes mellitus, hyperlipidemia,

FIG. 2. Diabetes, hyperlipidemia, and hypertension prevalence by BMI category.

FIG. 3. Diabetes, hyperlipidemia, and hypertension prevalence by age group.

and hypertension, respectively. There are strong linear associations between both age and BMI and disease prevalence for all 3 conditions, and these associations are consistently additive rather than interactive, that is, age and BMI combine in a straightforward fashion without any marked unpredictable interactions. Among the oldest patients (≥ 60) with the highest BMIs (≥ 50), the highest disease prevalences are found: 50.0% for diabetes, 52.3% for hyperlipidemia, and 72.6% for hypertension. There are only a few small inconsistencies in the linear relationships noted between BMI and disease prevalence after stratifying by age group; almost all of these inconsistencies reflect slightly higher disease prevalence for the underweight (< 18.5 BMI) group compared to the next lowest BMI group ($18.5 \leq \text{BMI} < 25$); these may reflect the effect of disease on underweight status rather than the effect of underweight status on disease.

Table 3 shows the combined effects of BMI and sex on the prevalence of diabetes, hyperlipidemia, and hypertension. As noted above, males have a higher prevalence of all 3 conditions, and the direct association between BMI and disease prevalence is virtually identical for males and females.

Table 4 shows the combined effects of BMI and race/ethnicity on the prevalence of diabetes, hyperlipidemia, and hypertension. As was true for the combined effects of BMI

and sex, the positive association between BMI and disease prevalence is highly consistent across all of the racial/ethnic groups.

Table 5 presents results of the association between BMI and the diagnosis of obesity. While BMI and reports of obesity as a medical problem are positively associated, obesity is substantially underreported. The medical definition of obesity includes having a BMI of 30 or above. However, only a little more than 10% of patients with $30 \leq \text{BMI} < 35$ are reported as obese, and only 57% were reported as obese at the highly serious threshold of $\text{BMI} \geq 50$.

The final analyses reported herein (Table 6) are logistic regressions that summarize the effects of demographic factors, obesity, and comorbidities on the prevalence of each of the 3 chronic conditions. These analyses begin with inclusion of the demographic factors and obesity and then add the relevant comorbidities. The analyses of diabetes prevalence reinforce the findings presented above: Whites have a higher risk of having diabetes than the reference category (odds ratio [OR] = 1.377), but African Americans and Hispanics have even higher odds (ORs = 2.201 and 2.754, respectively); males have higher odds than females (OR = 1.517); increasing age increases one's odds of having diabetes (for each year, OR = 1.060); and obesity leads to higher odds of having diabetes (OR = 1.113).

FIG. 4. Diabetes, hyperlipidemia, and hypertension prevalence by race/ethnicity.

FIG. 5. Diabetes, hyperlipidemia, and hypertension prevalence by sex.

Similarly, the logistic regression analyses of hypertension prevalence reinforce the findings presented earlier: Whites have a higher risk of having hypertension than the reference category ($OR=1.257$), and African Americans have even higher odds ($OR=2.572$), but the odds for Hispanics are

lower than those for whites (1.198); males have higher odds than females ($OR=1.269$); age increases one's odds of having hypertension (for each year, $OR=1.077$); and obesity also leads to higher odds ($OR=1.094$).

The logistic regression analyses of hyperlipidemia prevalence also confirm the earlier findings: while whites have a higher risk of having hyperlipidemia than the reference category ($OR=1.275$), African Americans and Hispanics have lower odds ($ORs=0.911$ and 0.937 , respectively); males have higher odds than females ($OR=1.560$); age increases one's odds of having hyperlipidemia (for each year, $OR=1.062$); and obesity also leads to higher odds ($OR=1.059$).

Discussion

The key role of obesity in the etiology of chronic disease in the United States is a focus of health researchers, health policy experts, and policy makers. For example, the American Heart Association and National Heart, Lung, and Blood Institute Scientific Statement includes, as one of its 7 conclusions, that "in the United States, the [metabolic] syndrome is strongly associated with the presence of abdominal obesity." The Statement also "recognizes several issues related to the metabolic syndrome that require additional research for

TABLE 1. PROPORTION OF PATIENTS WITH TYPE II DIABETES MELLITUS, HYPERLIPIDEMIA, AND HYPERTENSION BY BODY MASS INDEX (BMI) BAND, US, 2007

Patient Characteristics	Type II Diabetes		Hyperlipidemia		Hypertension		Total	% of Population
	n	%	n	%	n	%		
BMI Band								
BMI<18.5	4,449	1.0%	11,187	2.4%	13,873	3.0%	466,836	12.9%
18.5≤BMI<25	32,286	2.9%	140,251	12.7%	128,133	11.6%	1,103,105	30.5%
25≤BMI<30	74,113	7.6%	271,118	27.9%	236,754	24.4%	971,308	26.9%
30≤BMI<35	77,841	13.3%	197,816	33.7%	193,810	33.0%	586,469	16.2%
35≤BMI<40	51,130	18.5%	93,623	33.9%	104,940	38.0%	276,517	7.7%
40≤BMI<45	27,382	22.2%	40,148	32.6%	50,201	40.8%	123,078	3.4%
45≤BMI<50	13,396	25.9%	16,274	31.4%	22,676	43.8%	51,774	1.4%
BMI≥50	10,137	29.1%	10,502	30.1%	16,456	47.2%	34,861	1.0%
Total	290,734	8.0%	780,919	21.6%	766,843	21.2%	3,613,948	100.0%
Age Band								
Under 20	1,756	0.2%	4,269	0.5%	2,940	0.3%	883,151	24.4%
20–39 years	20,983	2.3%	66,254	7.1%	57,401	6.2%	931,816	25.8%
40–59 years	100,281	10.2%	298,927	30.3%	267,662	27.2%	985,418	27.3%
≥60 years	159,566	20.8%	389,476	50.8%	418,671	54.6%	766,812	21.2%
Missing	8,148	17.4%	21,993	47.0%	20,169	43.1%	46,751	1.3%
Total	290,734	8.0%	780,919	21.6%	766,843	21.2%	3,613,948	100.0%
Race								
White	83,224	9.3%	229,191	25.7%	214,944	24.1%	891,188	24.7%
African American	21,316	12.1%	30,183	17.2%	50,872	28.9%	175,952	4.9%
Hispanic	3,660	7.5%	4,852	9.9%	5,287	10.8%	48,812	1.4%
Oriental/Asian	782	7.2%	1,594	14.7%	1,395	12.9%	10,841	0.3%
Native American	567	8.5%	930	13.9%	938	14.0%	6,692	0.2%
Other Race	1,997	6.4%	4,792	15.4%	5,043	16.2%	31,142	0.9%
Unknown	179,188	7.3%	509,377	20.8%	488,364	19.9%	2,449,321	67.8%
Total	290,734	8.0%	780,919	21.6%	766,843	21.2%	3,613,948	100.0%
Sex								
Female	153,577	7.2%	410,588	19.2%	430,017	20.1%	2,135,205	59.1%
Male	137,132	9.3%	370,239	25.1%	336,762	22.8%	1,477,546	40.9%
Unknown	25	2.1%	92	7.7%	64	5.3%	1,197	0.0%
Total	290,734	8.0%	780,919	21.6%	766,843	21.2%	3,613,948	100.0%

TABLE 2. PROPORTION OF PATIENTS WITH TYPE II DIABETES MELLITUS, HYPERLIPIDEMIA, AND HYPERTENSION, BY BODY MASS INDEX (BMI) BAND AND AGE GROUP, US, 2007

	Under 20		20–39 years		40–59 years		≥60 years			
	Yes	%	Yes	%	Yes	%	Yes	%	Total	% of Population
Type II Diabetes										
BMI<18.5	231	0.1%	325	1.4%	1,161	7.6%	2,653	12.9%	466,836	12.9%
18.5≤BMI<25	491	0.2%	2,038	0.6%	6,749	2.9%	22,489	11.2%	1,103,105	30.5%
25≤BMI<30	299	0.3%	3,462	1.3%	19,485	6.1%	49,025	17.7%	971,308	26.9%
30≤BMI<35	244	0.7%	4,499	2.9%	26,635	12.1%	44,152	27.0%	586,469	16.2%
35≤BMI<40	217	1.4%	4,212	5.3%	21,175	19.2%	23,792	36.1%	276,517	7.7%
40≤BMI<45	137	2.3%	2,946	7.6%	12,724	25.1%	10,642	42.7%	123,078	3.4%
45≤BMI<50	68	2.9%	1,804	10.5%	6,786	30.8%	4,328	47.8%	51,774	1.4%
BMI≥50	69	4.1%	1,697	14.3%	5,566	35.5%	2,485	50.0%	34,861	1.0%
Total	1,756	0.2%	20,983	2.3%	100,281	10.2%	159,566	20.8%	3,613,948	100.0%
Hyperlipidemia										
BMI<18.5	472	0.1%	770	3.3%	2,911	18.9%	6,812	33.1%	466,836	12.9%
18.5≤BMI<25	1,171	0.4%	10,384	3.1%	40,763	17.5%	84,761	42.2%	1,103,105	30.5%
25≤BMI<30	933	1.1%	19,403	7.2%	97,513	30.6%	145,756	52.6%	971,308	26.9%
30≤BMI<35	777	2.2%	16,983	10.9%	81,514	37.1%	92,585	56.6%	586,469	16.2%
35≤BMI<40	494	3.3%	9,643	12.2%	42,604	38.6%	37,930	57.6%	276,517	7.7%
40≤BMI<45	251	4.1%	5,041	13.0%	19,494	38.4%	14,047	56.4%	123,078	3.4%
45≤BMI<50	102	4.3%	2,301	13.4%	8,357	37.9%	4,987	55.1%	51,774	1.4%
BMI≥50	69	4.1%	1,729	14.6%	5,771	36.8%	2,598	52.3%	34,861	1.0%
Total	4,269	0.5%	66,254	7.1%	298,927	30.3%	389,476	50.8%	3,613,948	100.0%
Hypertension										
BMI<18.5	244	0.1%	768	3.3%	3,027	19.7%	9,604	46.7%	466,836	12.9%
18.5≤BMI<25	588	0.2%	5,971	1.8%	28,402	12.2%	90,876	45.3%	1,103,105	30.5%
25≤BMI<30	518	0.6%	12,604	4.6%	71,020	22.3%	146,657	52.9%	971,308	26.9%
30≤BMI<35	521	1.4%	13,955	9.0%	73,779	33.6%	99,886	61.1%	586,469	16.2%
35≤BMI<40	475	3.2%	10,569	13.3%	46,548	42.1%	44,067	66.9%	276,517	7.7%
40≤BMI<45	285	4.7%	6,555	16.9%	24,308	47.9%	17,455	70.0%	123,078	3.4%
45≤BMI<50	174	7.3%	3,650	21.2%	11,657	52.9%	6,517	72.0%	51,774	1.4%
BMI>50	135	8.0%	3,329	28.1%	8,921	56.8%	3,609	72.6%	34,861	1.0%
Total	2,940	0.3%	57,401	6.2%	267,662	27.2%	418,671	54.6%	3,613,948	100.0%

clarification. Foremost is the need for improved strategies to achieve and sustain long-term weight reduction and increased physical activity.²² Similarly, the Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure includes obesity along with hypertension, dyslipidemia, and diabetes mellitus in its list of 9 major cardiovascular risk factors.²³

Utilizing the longitudinal data available in the GE Centricity EMR database, this study has assessed the associations between obesity and 3 key chronic disease states: type II diabetes, hyperlipidemia, and hypertension. The positive association between obesity and the 3 comorbid conditions in the GE Centricity database is similar to that found for the US population using other data sets and methods. Although obesity is underdiagnosed as an explicit problem in the Centricity EMR, as it is in medical records generally, the inclusion of BMI data in the Centricity database allows identification of obesity and allows estimation of the degree to which it is underreported.

The EMR data show both the high rates of prevalence of the 3 chronic diseases and the significant role of obesity in their etiology. Both bivariate and multivariate analyses document the role of obesity in predicting and explaining the prevalence of the 3 conditions and the value of the BMI data found in the Centricity EMR. While the demonstrated impact

of obesity is strong, this impact is most likely underestimated by using only 5 years of data, as was done in these analyses. Further analyses should be performed utilizing a longer time frame, an opportunity which will be facilitated by the expanding use of EMRs in the future.

Additional implications of the current analyses relate to optimal methods of analyzing EMR data. Specifically, exclusion algorithms were used to exclude retrospectively entered patient data, unconfirmed diagnoses, and duplicate patient data. Further research is needed to assess the feasibility and effectiveness of the existing techniques to maximize validity, as well as the extension of these techniques and development of additional ones.

Limitations

The limitations of this study originate primarily from 2 sources: (1) limitations of medical records in general and EMRs in particular, and (2) limitations that have been identified in the GE Centricity EMR database in particular. All medical records, manual or electronic, are subject to threats to validity, although EMRs generally reduce data errors and increase data validity compared to paper charts. For example, EMRs eliminate the problem of illegibility; yet EMR users can still make data entry errors. Similarly, regardless of whether records are manual or electronic, there will be gaps

FIG. 6. Prevalence of (A) diabetes, (B) hyperlipidemia, and (C) hypertension determined by body mass index (BMI) category and age group.

TABLE 3. PROPORTION OF PATIENTS WITH TYPE II DIABETES MELLITUS, HYPERLIPIDEMIA, AND HYPERTENSION, BY BODY MASS INDEX (BMI) BAND AND SEX, US, 2007

	<i>Female</i>		<i>Male</i>		<i>Total</i>	
	<i>n</i>	<i>%</i>	<i>n</i>	<i>%</i>	<i>n</i>	<i>% of Population</i>
Type II Diabetes						
BMI<18.5	2,833	1.2%	1,615	0.7%	466,836	12.9%
18.5≤BMI<25	17,622	2.4%	14,660	3.9%	1,103,105	30.5%
25≤BMI<30	33,889	6.5%	40,221	8.9%	971,308	26.9%
30≤BMI<35	37,333	11.6%	40,500	15.3%	586,469	16.2%
35≤BMI<40	28,331	16.3%	22,794	22.2%	276,517	7.7%
40≤BMI<45	17,240	20.1%	10,139	27.1%	123,078	3.4%
45≤BMI<50	9,017	23.8%	4,379	31.7%	51,774	1.4%
BMI≥50	7,312	27.9%	2,824	32.8%	34,861	1.0%
Total	153,577	7.2%	137,132	9.3%	3,613,948	100.0%
Hyperlipidemia						
BMI<18.5	7,714	3.2%	3,471	1.6%	466,836	12.9%
18.5≤BMI<25	89,920	12.4%	50,314	13.3%	1,103,105	30.5%
25≤BMI<30	125,288	24.0%	145,799	32.4%	971,308	26.9%
30≤BMI<35	92,833	28.9%	104,953	39.7%	586,469	16.2%
35≤BMI<40	51,387	29.6%	42,229	41.1%	276,517	7.7%
40≤BMI<45	25,019	29.2%	15,126	40.4%	123,078	3.4%
45≤BMI<50	10,888	28.7%	5,386	39.0%	51,774	1.4%
BMI≥50	7,539	28.7%	2,961	34.4%	34,861	1.0%
Total	410,588	19.2%	370,239	25.1%	3,613,948	100.0%
Hypertension						
BMI<18.5	9,909	4.1%	3,961	1.8%	466,836	12.9%
18.5≤BMI<25	83,761	11.6%	44,363	11.7%	1,103,105	30.5%
25≤BMI<30	118,459	22.7%	118,274	26.3%	971,308	26.9%
30≤BMI<35	97,457	30.3%	96,335	36.4%	586,469	16.2%
35≤BMI<40	60,297	34.7%	44,637	43.5%	276,517	7.7%
40≤BMI<45	32,558	38.0%	17,639	47.2%	123,078	3.4%
45≤BMI<50	15,584	41.1%	7,092	51.3%	51,774	1.4%
BMI≥50	11,992	45.7%	4,461	51.8%	34,861	1.0%
Total	430,017	20.1%	336,762	22.8%	3,613,948	100.0%

if information is not forwarded from one provider to another, even though EMRs should eventually improve such information exchange. Additionally, there may also be free text data in an EMR that are not available for analysis because of the complexities and costs of text mining. Still, in general, EMRs have great potential to mitigate many of the above-mentioned problems.

The second class of methodological limitations, those identified as potential problems in using GE Centricity EMR data, include invalid retrospective patient data, unconfirmed diagnoses, and duplicate patient data. As described earlier, inclusion/exclusion algorithms were developed and applied to address each of these issues. However, available data do not allow comprehensive assessment of the success of this strategy to eliminate all possible duplication and underestimation/overestimation. A specific potential limitation of the study stems from the fact that physicians' diagnoses of hypertension may not have taken into account that, for diabetics, hypertension is defined as systolic blood pressure ≥130 mmHg and diastolic blood pressure ≥80 mmHg, while for other populations, systolic blood pressure is defined as ≥140 mmHg and diastolic blood pressure is defined as ≥90 mmHg. However, it is unlikely that physicians failed to take this distinction into account when making diagnoses and recording problems. Still, to the degree that this distinction was ignored in physicians' diagnoses, the present

analyses underestimate diabetes prevalence and probably underestimate associations between prevalence of diabetes, hypertension, and hyperlipidemia.

Conclusions

The study presented herein analyzes GE Centricity EMR data in order to determine the effects of BMI and obesity, key risk factors, and components of metabolic syndrome, on the prevalence of 3 chronic diseases that have both high prevalence and high disease burdens: type II diabetes mellitus, hyperlipidemia, and hypertension. While BMI and obesity are often underreported in surveys and administrative databases, EMR databases such as the GE Centricity EMR database may be good sources of relevant data and may allow tracking such variables over time, and facilitate longitudinal analyses of increasing disease prevalence and progression.

This study demonstrates strong positive associations of age, sex, race/ethnicity, BMI, and comorbidities with diagnosed prevalence of the 3 conditions. Results include uniform positive associations between both age and BMI and the prevalence of each condition; uniformly greater prevalence for males than for females; and associations with race/ethnicity, with African Americans having the highest prevalence of diagnosed type II diabetes and hypertension and whites having the highest prevalence of diagnosed

TABLE 4. PROPORTION OF PATIENTS WITH TYPE II DIABETES MELLITUS, HYPERLIPIDEMIA, AND HYPERTENSION, BY BODY MASS INDEX (BMI) BAND AND RACE/ETHNICITY, US, 2007

	White		African American		Hispanic		Oriental/Asian		Native American			
	n	%	n	%	n	%	n	%	n	%	n	% of Population
Diabetes Type II												
BMI<18.5	1,343	1.3%	423	1.4%	43	0.3%	19	0.9%	9	0.7%	466,836	12.9%
18.5≤BMI<25	9,157	3.4%	2,113	5.2%	405	3.1%	277	5.6%	63	3.4%	1,103,105	30.5%
25≤BMI<30	21,065	8.7%	4,750	12.5%	1,039	10.2%	257	10.4%	139	8.2%	971,308	26.9%
30≤BMI<35	22,247	14.9%	5,387	17.8%	1,030	16.5%	151	17.5%	149	14.5%	586,469	16.2%
35≤BMI<40	14,922	21.0%	3,960	21.9%	584	20.3%	47	20.1%	103	22.7%	276,517	7.7%
40≤BMI<45	7,847	24.8%	2,332	24.0%	328	27.9%	17	20.0%	58	31.2%	123,078	3.4%
45≤BMI<50	3,788	28.6%	1,241	26.5%	121	25.8%	10	38.5%	24	30.0%	51,774	1.4%
BMI≥50	2,855	33.3%	1,110	28.6%	110	32.1%	4	30.8%	22	30.6%	34,861	1.0%
Total	83,224	9.3%	21,316	12.1%	3,660	7.5%	782	7.2%	567	8.5%	3,613,948	100.0%
Hyperlipidemia												
BMI<18.5	3,564	3.4%	469	1.5%	61	0.4%	31	1.4%	15	1.1%	466,836	12.9%
18.5≤BMI<25	41,590	15.4%	3,617	8.9%	616	4.7%	618	12.5%	145	7.7%	1,103,105	30.5%
25≤BMI<30	78,187	32.2%	8,134	21.3%	1,689	16.5%	601	24.3%	309	18.3%	971,308	26.9%
30≤BMI<35	57,825	38.6%	8,037	26.6%	1,356	21.8%	240	27.9%	240	23.4%	586,469	16.2%
35≤BMI<40	27,835	39.2%	5,061	27.9%	658	22.9%	66	28.2%	134	29.6%	276,517	7.7%
40≤BMI<45	12,077	38.2%	2,655	27.3%	280	23.8%	26	30.6%	50	26.9%	123,078	3.4%
45≤BMI<50	4,875	36.8%	1,221	26.1%	116	24.7%	10	38.5%	20	25.0%	51,774	1.4%
BMI≥50	3,238	37.7%	989	25.5%	76	22.2%	2	15.4%	17	23.6%	34,861	1.0%
Total	229,191	25.7%	30,183	17.2%	4,852	9.9%	1,594	14.7%	930	13.9%	3,613,948	100.0%
Hypertension												
BMI<18.5	4,096	3.9%	1,146	3.8%	72	0.5%	45	2.0%	23	1.8%	466,836	12.9%
18.5≤BMI<25	37,021	13.7%	6,683	16.4%	663	5.1%	490	9.9%	115	6.1%	1,103,105	30.5%
25≤BMI<30	66,549	27.4%	12,721	33.4%	1,661	16.2%	496	20.0%	271	16.1%	971,308	26.9%
30≤BMI<35	53,647	35.8%	12,645	41.9%	1,426	22.9%	247	28.7%	251	24.5%	586,469	16.2%
35≤BMI<40	29,172	41.1%	8,386	46.3%	816	28.4%	70	29.9%	160	35.3%	276,517	7.7%
40≤BMI<45	13,915	44.0%	4,728	48.6%	360	30.6%	28	32.9%	58	31.2%	123,078	3.4%
45≤BMI<50	6,171	46.5%	2,376	50.8%	157	33.5%	14	53.8%	30	37.5%	51,774	1.4%
BMI≥50	4,373	51.0%	2,187	56.3%	132	38.5%	5	38.5%	30	41.7%	34,861	1.0%
Total	214,944	24.1%	50,872	28.9%	5,287	10.8%	1,395	12.9%	938	14.0%	3,613,948	100.0%

hypertension. Despite the differences in disease prevalence by sex and by racial/ethnic group, the positive associations between BMI and disease prevalence are consistent for males and females and across all racial/ethnic groups. Finally, the logistic regression analyses control for possible confounding and demonstrate the unique effects of each of the factors—age, sex, racial/ethnic group, obesity, and comorbidities—in explaining and predicting the prevalence of the 3 chronic conditions. All of these results contribute to the growing literature regarding the adverse effects of obesity and

TABLE 5. PROPORTION OF ACTUAL DIAGNOSIS OF OBESITY BY BODY MASS INDEX (BMI) BAND*

BMI Band	Total	Obesity Diagnosis	Obesity Diagnosis %
BMI<18.5	311,513	1,921	0.6%
18.5≤BMI<25	760,079	3,381	0.4%
25≤BMI<30	683,330	15,932	2.3%
30≤BMI<35	412,522	42,187	10.2%
35≤BMI<40	193,518	43,458	22.5%
40≤BMI<45	86,010	30,209	35.1%
45≤BMI<50	36,068	16,489	45.7%
BMI≥50	24,158	13,728	56.8%
Total	2,507,198	167,305	6.7%

*ICD-9-CM diagnosis codes 278.00 and 278.01

TABLE 6. LOGISTIC REGRESSION ANALYSES: EACH DIAGNOSIS BY DEMOGRAPHICS AND OBESITY, WITHOUT AND WITH COMORBID CONDITIONS

	Significance	Odds ratio
Type II Diabetes Diagnosis		
White	0.000	1.377
African American	0.000	2.201
Hispanic	0.000	2.754
Age	0.000	1.060
Obesity	0.000	1.113
Sex	0.000	1.517
Hypertension		
White	0.000	1.257
African American	0.000	2.572
Hispanic	0.000	1.198
Age	0.000	1.077
Obesity	0.000	1.094
Sex	0.000	1.269
Hyperlipidemia		
White	0.000	1.275
African American	0.000	0.911
Hispanic	0.001	0.937
Age	0.000	1.062
Obesity	0.000	1.059
Sex	0.000	1.560

Variable(s) entered: White, African American, Hispanic, age, obesity, sex.

demonstrate the value of EMR databases for studying and documenting such effects.

Author Disclosure Statement

Dr. Haas is employed by Ethicon Endo-Surgery, Inc. Ethicon Endo-Surgery, Inc., funded this study, and it contracted with all other authors listed on this article. Drs Crawford, Cote, Couto, Daskiran, Gunnarsson, Haas, Nigam, and Schuette, and Ms. Haas disclosed no other financial ties or conflicts of interest.

References

1. Reaven GM. Role of insulin resistance in human disease. The Banting lecture. *Diabetes* 1988;37:1595–1607.
2. Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: Diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 1998;15:539–553.
3. Balkau B, Charles MA. Comment on the provisional report from the WHO consultation. European Group for the Study of Insulin Resistance (EGIR). *Diabet Med* 1999;16:442–443.
4. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* 2001;285:2486–2497.
5. Alberti K, Zimmet P, Shaw J. Metabolic syndrome—A new world-wide definition. A consensus statement from the International Diabetes Federation. *Diabet Med* 2006;23:469–480.
6. Lawlor D, Davey Smith GD, Ebrahim S. Does the new International Diabetes Federation definition of the metabolic syndrome predict CHD any more strongly than older definitions? Findings from the British Women's Heart and Health Study. *Diabetologia* 2006;49:41–48.
7. Ford ES. Risks for all-cause mortality, cardiovascular disease, and diabetes associated with the metabolic syndrome: A summary of the evidence. *Diabet Care* 2005;28:1769–1778.
8. Malik S, Nathan D, Wong S, et al. Impact of the metabolic syndrome on mortality from coronary heart disease, cardiovascular disease, and all causes in United States adult. *Circulation* 2004;110:1245–1250.
9. Meigs JB, Wilson PWF, Nathan DM, et al. Prevalence and characteristics of the metabolic syndrome in the San Antonio Heart and Framingham Offspring Studies. *Diabetes* 2003;52:2160–2167.
10. Hanley AJG, Karter AJ, Williams K, et al. Prediction of type 2 diabetes mellitus with alternative definitions of the metabolic syndrome: The insulin resistance atherosclerosis study. *Circulation* 2005;112:3713–3721.
11. Hong Y, Jin X, Mo J, et al. Metabolic syndrome, its preeminent clusters, incident coronary heart disease and all-cause mortality. Results of prospective analysis for the atherosclerosis risk in communities study. *J Intern Med* 2007;262:113–123.
12. Meigs JB, Rutter MK, Sullivan LM, et al. Impact of insulin resistance on risk of type 2 diabetes and cardiovascular disease in people with metabolic syndrome. *Diabetes Care* 2007;30:1219–1225.
13. Park Y, Shu S, Palaniappan L, et al. The metabolic syndrome. *Arch Int Med* 2003;163:427–436.
14. National Heart, Lung, and Blood Institute and Boston University. Framingham heart study. Available at: <http://www.nhlbi.nih.gov/about/framingham/index.html>. Accessed December 12, 2007.
15. Kannel WB, Feinleib M, McNamara PM, Garrison RJ, Castelli SP. An investigation of coronary heart disease in families: The Framingham Offspring Study. *Am J Epidemiol* 1979;110:281–290.
16. Bays HE, Chapman RH, Grandy S, for the SHIELD Investigators' Group. The relationship of body mass index to diabetes mellitus, hypertension and dyslipidaemia: comparison of data from two national surveys. *Int J Clin Pract* 2007;61:737–747.
17. McTigue K, Larson JC, Valoski A, et al. Mortality and cardiac and vascular outcomes in extremely obese women. *JAMA* 2006;296:79–86.
18. Sturm R. Increases in morbid obesity in the USA: 2000–2005. *Public Health* 2007;121:492–496.
19. Brixner D, Said Q, Kirkness C, Oberg B, Ben-Joseph R, Oderda G. Assessment of cardiometabolic risk factors in a national primary care electronic health record database. *Value in Health* 2007;10:S29–S36.
20. Gill J, Chen Y. Quality of lipid management in outpatient care: A national study using electronic health records. *Am J Med Qual* 2008;23:375–381.
21. Oderda G, Brixner D, Lieberman M. Comparison of EMR data to US national data. Paper presented at: International Society for Pharmacoeconomics and Outcomes Research (ISPOR) 10th Annual European Congress; October 20–23, 2007; Dublin, Ireland.
22. Grundy SM, Cleeman JI, Daniels SR, et al. Management of the metabolic syndrome: An American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 2005;112:2735–2752.
23. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, and the National High Blood Pressure Education Program Coordinating Committee. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: The JNC 7 report. *JAMA* 2003;289:2560–2572.

Address correspondence to:

Albert G. Crawford, Ph.D.

Thomas Jefferson University School of Population Health

1015 Walnut Street, Suite 115

Philadelphia PA 19107

E-mail: Albert.Crawford@jefferson.edu

This article has been cited by:

1. Ahmad Ghiyath Shayeb, Kirsten Harrild, Eileen Mathers, Siladitya Bhattacharya. 2012. Response: Re: An exploration of the association between male body mass index and semen quality. *Reproductive BioMedicine Online* . [[CrossRef](#)]
2. Fasil Tekola Ayele, Ayo Doumatey, Hanxia Huang, Jie Zhou, Bashira Charles, Michael Erdos, Jokotade Adeleye, Williams Balogun, Olufemi Fasanmade, Thomas Johnson, Johnnie Oli, Godfrey Okafor, Albert Amoah, Benjamin A. Eghan, Kofi Agyenim-Boateng, Joseph Acheampong, Clement A. Adebamowo, Alan Herbert, Norman Gerry, Michael Christman, Guanjie Chen, Daniel Shriner, Adebowale Adeyemo, Charles N. Rotimi. 2011. Genome-wide associated loci influencing interleukin (IL)-10, IL-1Ra, and IL-6 levels in African Americans. *Immunogenetics* . [[CrossRef](#)]
3. Li Cai, Aiping Liu, Lei Zhang, Shuping Li, Peiyu Wang. 2011. Prevalence, Awareness, Treatment, and Control of Hypertension among Adults in Beijing, China. *Clinical and Experimental Hypertension* 1-8. [[CrossRef](#)]
4. A. Janet Tomiyama, Mary F. Dallman, Elissa S. Epel. 2011. Comfort food is comforting to those most stressed: Evidence of the chronic stress response network in high stress women. *Psychoneuroendocrinology* . [[CrossRef](#)]
5. Fahad Javed, Emad F. Aziz, Manpreet S. Sabharwal, Girish N. Nadkarni, Shahzeb A. Khan, Juan P. Cordova, Alexandre M. Benjo, Dymphna Gallagher, Eyal Herzog, Franz H. Messerli, F. Xavier Pi-Sunyer. 2011. Association of BMI and Cardiovascular Risk Stratification in the Elderly African-American Females. *Obesity* **19**:6, 1182-1186. [[CrossRef](#)]
6. Frank K. Friedenberg, Derek M. Tang, Vishwas Vanar, Thais Mendonca. 2011. Predictive Value of Body Mass Index at Age 18 on Adulthood Obesity: Results of a Prospective Survey of an Urban Population. *The American Journal of the Medical Sciences* 1. [[CrossRef](#)]
7. Ehud Grossman, Franz H. Messerli. 2011. Management of Blood Pressure in Patients With Diabetes. *American Journal of Hypertension* . [[CrossRef](#)]
8. Ashley Sinclair Felix, Faina Linkov, George Larry Maxwell, Camille Ragin, Emanuela Taioli. 2011. Racial Disparities in Risk of Second Primary Cancers in Patients With Endometrial Cancer. *International Journal of Gynecological Cancer* 1. [[CrossRef](#)]