

2013

Infections in Anterior Cruciate Ligament Reconstruction

C. Stucken

Rothman Institute, Thomas Jefferson University, Philadelphia, Pennsylvania.

D.N. Garras

Rothman Institute, Thomas Jefferson University, Philadelphia, Pennsylvania.

J.L. Shaner

Rothman Institute, Thomas Jefferson University, Philadelphia, Pennsylvania.

Steven B. Cohen

Thomas Jefferson University, Steven.Cohen@rothmaninstitute.com

Let us know how access to this document benefits you

Follow this and additional works at: https://jdc.jefferson.edu/rothman_institutePart of the [Orthopedics Commons](#)

Recommended Citation

Stucken, C.; Garras, D.N.; Shaner, J.L.; and Cohen, Steven B., "Infections in Anterior Cruciate Ligament Reconstruction" (2013). *Rothman Institute*. Paper 59.https://jdc.jefferson.edu/rothman_institute/59

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Rothman Institute by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

As submitted to:

Sports Health

And later published as:

Infections in Anterior Cruciate Ligament Reconstruction

Stucken, C., Garras, D. N., Shaner, J. L., & Cohen, S. B.
Volume 5, Issue, 6, pages: 553-557.
2013

DOI: 10.1177/1941738113489099

Abstract:

Context: Anterior cruciate ligament (ACL) reconstruction is a safe, common, and effective method of restoring stability to the knee after injury, but evolving techniques of reconstruction carry inherent risk. Infection after ACL reconstruction, while rare, carries a high morbidity potentially resulting in poor clinical outcome.

Evidence Acquisition: Data were obtained from previously published peer-reviewed literature through a search of the entire PubMed database (up to December 2012) as well as from textbook chapters.

Results: Treatment with culture-specific antibiotics and debridement with graft retention is generally more effective than graft removal, but with persistent infection consideration should be given to graft removal. Graft type likely has no effect on infection rates.

Conclusions: The early diagnosis of infection and appropriate treatment are necessary to avoid the complications of articular cartilage damage and arthrofibrosis.

Keywords: Anterior cruciate ligament, infection, graft type

Introduction

Anterior cruciate ligament (ACL) rupture is the most frequent ligamentous injury of the knee (42). It is frequently injured in the young athletes performing cutting and pivoting sports, and predisposes the knee to subsequent injuries, as well as the potential for earlier onset of osteoarthritis (15, 35, 51). Arthroscopic ACL reconstruction is a common and effective method of restoring stability to the knee after injury, with more than 400,000 ACL reconstructions performed annually in the U.S. (18, 40, 39, 26, 38, 43, 45, 46, 65, 79, 27). Like any surgical procedure, a number of potential complications have been recognized that may affect functional outcome (41, 57, 12, 3, 69). Specifically infection after ACL reconstruction can be a devastating complication. Overall, infection rates are low after ACL reconstruction (58). Despite theoretical risk of disease transmission and higher graft failure in irradiated grafts, the use of allograft tissue continues to gain popularity for a number of reasons.

Allograft vs. Autograft: Clinical Outcomes

Graft material has received a significant amount of attention when focusing on surgical variables (14). The ideal graft would have low infection rates, low cost, low donor-site morbidity, fast incorporation, and wide availability. Autograft reconstruction remains the gold standard for younger patients, and despite the frequency of reconstruction surgery there is considerable controversy and variability in graft selection. No perfect graft exists. The use of

allografts in ACL reconstruction has increased significantly as both surgeons and patients desire limited donor-site morbidity, ease of use, decreased operative time, lower incidence of postoperative arthrofibrosis, low residual muscle weakness, availability and variety of graft sizes, decreased postoperative pain, improved cosmetic appearance, and commencement of earlier rehabilitation with faster return to work (27, 80, 70, 11, 67). Autograft harvest is not a completely benign event and has potential short and long-term morbidity, limiting options during procedural complications of graft harvest and revision reconstruction (5). Desire to avoid the sacrifice of autologous tissue, as well as to minimize surgical trauma and postoperative donor-site morbidity, has promoted consideration and increased use of allograft tissue. Tissue autografts amenable for use include central third bone-patellar tendon-bone (BTB), semitendinosus-gracilis graft from the hamstring (HS), or quadriceps tendon graft. Allograft tissues include the tibialis anterior or posterior, BTB, Achilles tendon with bone plug, fascia lata, peroneus longus, HS, and quadriceps tendons. Ultimately, graft selection is influenced by preoperative physical exam, patient age and activity level, as well as by the surgeon's preference, experience, and bias (28). A number of studies have demonstrated allograft efficacy in improving stability and function, and clinical outcomes comparable to reconstruction with autologous tissue, making them an acceptable alternative to autografts (32, 75, 52, 47, 81, 13, 21). Yet, other studies have suggested higher failure rates with allograft tissue (8, 55, 60, 16, 7, 53, 71). While no consensus exists, allografts have been recommended for patients over 45 because they permit quicker recovery when morbidity associated with graft procurement is eliminated (80).

Allograft vs. Autograft: Incidence of Sepsis

Allograft tissue has become an acceptable graft choice for ACL reconstruction, raising considerable questions regarding the risk of viral and bacterial transmission from contaminated tissue. Although the risk is low, these confirmed cases represent a major medical and surgical challenge. To date there have been three reported incidences of viral disease transmission from bone-patellar tendon-bone allografts used to reconstruct the ACL. One case of HIV was reported in 1995, and two cases of hepatitis C have been reported (82, 29, 76, 77, 25). The American Association of Tissue Banks (AATB) recommends routine serologic screening for HIV, human T-cell leukemia virus, hepatitis B, hepatitis C, aerobic and anaerobic bacteria, and syphilis. Overall, the risk of HIV transmission with connective tissue allografts is estimated to be one in 1.6 million (19, 17, 27).

The Center for Disease Control (CDC) and Prevention reported 26 cases of bacterial infections associated with musculoskeletal tissue allografts after the reported death of a recipient of an allograft (femoral condyle) contaminated with *Clostridium*. Thirteen of the cases were infected with *Clostridium*, and fourteen were associated with a single tissue processor. All were processed aseptically, but none underwent terminal sterilization. The CDC also described two cases of septic arthritis following allograft ACL reconstructions from a common donor at a Texas-based tissue bank, and two from a common donor at a Florida-based tissue bank (23, 24). Completed sterilization techniques were confirmed from the Texas-based tissue bank, but sterilization procedures were mistakenly not performed on the tissue from the Florida-based tissue bank. Tissue bank regulation has increased dramatically since the occurrence of these documented transmissions, and as of May 2005, all tissue banks in the United States were required to conform to the FDA's "Good Tissue Practice" guidelines, which permit inspection of

tissue bank facilities and specify minimum standards for tissue recovery, testing, and processing (48).

Septic arthritis remains a rare but devastating complication following ACL reconstruction. Reports in the literature show an incidence of 0.14% to 1.70% (20, 33, 44, 59, 86, 88). Reports evaluating the difference in the rates of postoperative infection with allografts versus autografts are limited, and the current literature provides conflicting data. Using the PubMed database we sought to study the largest available trials that specifically investigated the incidence of infection in allograft and autograft ACL reconstruction (Table 1). Some papers have suggested no difference in infection rates between allo- and autograft. Barker et al. reviewed 3,126 ACL reconstructions with 1,777 autografts and 1,349 allografts, finding infection rates of 0.44% in allografts and 0.68% in autografts (6). Although their infection rates were lower, Indelli et al also found no significant difference between allograft and autograft (44). They reviewed 3,500 arthroscopic ACL reconstructions and found infection rates of 0.29% (4/1,400) for BTB autograft and 0.1% (2/2,100) for allograft. Greenberg et al examined 861 patients and found no infections in either group (36). Our research has also suggested no difference as well (34). We looked at 788 ACL reconstructions, 535 allograft and 253 autograft, with infection rates of 0.74% (four patients) and 0.79% (two patients), respectively.

Conversely, several studies have reported differences in the rates of infection associated with graft type. Katz et al showed an increased rate of infection among autografts (1.2%) versus allografts (0.63%), but it was not statistically significant (50). Explanation for this trend toward a higher infection rate in the autograft population included longer surgical time, more invasive tissue dissection, and longer preparation of graft. Furthermore, the same type of “tube-within-a-tube” tissue harvester suspected by Tuman et al. as a source of contamination when not

disassembled during sterilization was used during their study (83). Meanwhile, data reported by both Wang et al and Crawford et al suggested a higher rate of infection in allograft tissue than autograft tissue (Wang 1.11% versus 0.5%; Crawford 3.8% versus 0%, respectively) (87, 30). Both of these studies have their flaws. In the study by Wang et al, the one infection in 90 allograft reconstructions skewed their results when compared to 20 infections in 3,978 autografts. The Crawford et al study reported 11 infections in 290 allografts (3.8%) in which none of the allografts had undergone sterilization procedures, and no infections in 41 autografts. These seven studies represent the largest series comparing infection rates in allografts with autografts. As shown in Table 1, pooling these data together to include over 13,000 reconstructions shows infection rates for autograft and allografts to be 0.51% and 0.49%, respectively.

Although it has been hypothesized that allograft contamination has the potential for disease transmission, the link between contaminated grafts and clinical infections has been called into question. Guelich et al evaluated the utility of culturing allografts, demonstrating a positive bacterial culture rate of 9.7% (24 of 247 allografts) (37). These patients did not receive antibiotics in addition to the routine use of preoperative prophylactic antibiotics, and none went on to develop septic arthritis or wound complications. Likewise, Diaz-de-Rada et al, had 24 positive cultures from 181 allograft implantations analyzed; no patients showed clinical infection during follow-up (31). Hence, culture positive evidence of allograft contamination did not correlate with infectious complications (22).

Autograft: Bone-patella tendon-bone vs. hamstring incidence of sepsis

Some reports suggest an increased rate of infection with hamstring tendon autograft as compared with BTB autograft. Barker et al had 18 cases of septic arthritis after ACL reconstruction, there was a statistically significant increased risk of infection with hamstring tendon autograft (1.44%) compared with BTB autograft (0.49%) (6). Other studies have demonstrated a trend toward increased risk with hamstrings, but have not achieved statistical significance. In one of the largest reports, Wang et al reviewed 21 cases of infection from a population of 4,068 patients with ACL reconstruction (87). The majority of reconstructions were performed using autografts, and 20 (0.57%) infections occurred among 3536 patients reconstructed with hamstrings; no infections occurred among 442 post BTB autografts. A possible explanation included the frequent use of flash sterilization that was correlated with the occurrence of infections. Judd et al found similar results in a review of 217 BTB autografts and 192 hamstring autografts where all 11 intra-articular infections occurred in the hamstring group (49). The high rate of infection among hamstring autografts was attributed to the choice of graft fixation or additional soft-tissue dissection rather than the graft itself. Katz et al also had two of 118 hamstring autograft infections and 0 of 52 BTB (50). Burks et al reported eight postoperative ACL infections from a population of 1,918 arthroscopic ACL reconstructions, including seven hamstring tendon autografts and only one BTB autograft (20). One explanation for this trend was the relatively short length of hamstring graft places suture material inside the joint, which acts as a foreign body predisposing the joint to infections (10).

Diagnosis of infection

Prompt diagnosis and treatment of a septic joint is necessary for infection control and to achieve best long-term clinical outcomes. Postoperative infections are classified as either

acute (<two weeks), subacute (two weeks to two months) or late (>two months). Most patients have acute or subacute presentation of symptoms (20, 33, 44, 59, 86, 88, 6, 50, 87, 49, 10, 85, 74, 72). The most consistent findings include increased pain, inflammation and moderate effusion, whereas fevers, chills, erythema and drainage are not consistently present (44, 88, 49, 72). While Armstrong et al state that typical postoperative pain usually lasts for only 1-2 days and pain lasting longer than this should be suggestive of septic arthritis, there is a high degree of patient variability with infected patients generally having more pain than expected (2). The diagnosis of septic arthritis can be difficult to make in the early postoperative period as knee swelling, inflammation, and stiffness may be interpreted as normal, making laboratory data crucial to establishing the diagnosis. Laboratory values including erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP) are recommended to confirm diagnosis as they have high negative predictive values. Wang et al used ESR, CRP, and fibrinogen (FIB) levels more than 50 mm/h, 6 mg/mL, and 800 mg/mL, respectively, for septic arthritis (87). Margheritini et al believe CRP to be a more accurate predictor of postoperative complications than ESR, as both can be elevated postoperatively, but CRP rises and falls more rapidly than ESR levels. A sustained elevation of CRP beyond two weeks or new rise should prompt investigation to rule out infection (56). Complete CRP levels normalization is generally seen after 2-12 weeks. Synovial fluid aspirate is still the best diagnostic test for infection (20). Progressively higher synovial WBC counts are seen with septic arthritis; polymorphonuclear cells greater than 90% are highly predictive (44). Magnetic Resonance Imaging (MRI) can substantiate the diagnosis of infection in uncertain scenarios, and it can further determine the extent of the infection and detect any specific fluid collection. Findings on imaging can include joint effusion, synovitis, bone

erosions, edema of adjacent soft tissues and bone marrow, sinus tracts, and soft tissue abscesses (68).

Risk factors for infection

Several risk factors common for infection following ACL reconstruction include intra-articular corticosteroid injection, systemic corticosteroids, immunocompromised state, prior or concomitant procedures on the same knee, and history of previous knee infection (88, 49, 62, 1). More recent literature suggests other possible risk factors including graft type, operative time, tourniquet time, foreign body load, and usage of drains (3, 69, 20, 59, 86, 88, 74, 72, 58, 4). Individual reports have implicated methods of sterilization, instrumentation, and specific hardware used for graft fixation. Wang et al noted that flash sterilization of instrumentation was correlated with high rates of infections (87). Tuman et al suggested that failure to disassemble a “tube-within-a-tube” hamstring harvester may lead to unsatisfactory sterilization, providing a potential source for contamination (83). Judd et al found a higher incidence of infection associated with post/washer/braided suture construct in hamstring autograft fixation. This may be related to soft tissue injury during hamstring harvesting combined with relatively subcutaneous position of the metallic construct (49). Eight of 11 patients had concomitant extra-articular wound infection at this site with cultures positive for the same causative organism.

Many different microorganisms have been cultured from synovial fluid of with septic arthritis, but the most common pathogen is *Staphylococcus aureus*. Causative bacteria also include coagulase-negative *Staphylococcus*, *Propionibacter acnes*, *Peptostreptococcus*, *Enterobacter*, *Enterococcus*, *Pseudomonas aeruginosa*, *Escherichia coli*, and *Klebsiella* (20, 33, 44, 59, 86, 88, 6, 50, 87, 49, 10, 85, 74, 72). Methicillin-resistant *Staphylococcus aureus*

(MRSA) were also cultured among patients. (Au: Reference needed.) More unusual organisms have also identified in case reports, including *Mycobacterium fortium*, *Mycobacterium tuberculosis*, *Staphylococcus lugdenesis*, *Erysipelothrix rhusio pathiae*, and fungal species *Rhizopus*, *Microsporus* and *Candida albicans* (61, 84, 63, 66, 64). Establishing the causative organism in an intra-articular infection is crucial to predicting prognosis and commencing appropriate treatment. MRSA joint fluid is a virulent organism which typically presents acutely and entails greater need for lavage and for graft removal (54). Given that the majority of infections are with skin flora and are associated with concomitant extra-articular sites for infection, inoculation may occur at the time of surgery or shortly thereafter.(59, 49) including concomitant inside-out meniscus repair (59, 88). In subacute or late septic knee arthritis, the tibial or femoral sites may be infected and spread to the knee joint from hematoma collection in the subcutaneous tissue (49, 10, 33, 44).

Treatment of infection

Suspected joint infection is an emergency with the two main treatment goals: protecting the articular cartilage and the graft. In an animal study cartilage test, more than half its glycosaminoglycan and collagen within seven days from the onset of infection (78). Prompt intravenous antibiotic therapy to cover the most common organisms (*Staphylococcus aureus* and coagulase-negative *Staphylococcus*) should be given as soon as laboratory studies and joint fluid have been obtained. A third-generation cephalosporin or vancomycin is recommended (44, 59, 86, 88, 49, 72). When there is a strong clinical suspicion, antibiotic therapy should be continued even if synovial fluid cultures are negative (44, 86, 72).

There are several options for treatment of post-reconstruction septic arthritis of the knee (45). Barrett and Field recommended joint debridement with graft and hardware removal (9). Burks et al recommend arthroscopic graft removal with six weeks of antibiotics, followed by early re-implantation within six weeks of completing antibiotic treatment (20). Open arthrotomy with removal of hardware and curettage of tunnels has been recommended (89). Staging treatment with multiple initial debridements followed by placement of antibiotic-impregnated polymethylmethacrylate (PMMA) beads, and final re-implantation at 6-8 months may be best (74). A full course of antibiotic treatment may proceed with arthroscopic debridement in case of persistent clinical or abnormal laboratory findings (86). There is an algorithm for the treatment of infections based on a series of seven infections in 2,500 ACL reconstructions (88).

A survey of Sports Medicine Fellowship Program Directors regarding a standard of care for infections found 85% use culture-specific antibiotics and surgical irrigation of the joint with graft retention as the initial treatment for infected patellar tendon autografts and 64% for infected allografts (58). For cases resistant to initial treatment, the most common treatment (39%) favored continuing IV antibiotics with repeated surgical irrigation and graft retention. Thirty-one percent of respondents did recommend a combination of IV antibiotics, hardware removal and graft removal in resistant cases. Overall, graft removal was not considered the standard of care for initial treatment; it was chosen for 6% of autografts and 33% of allografts. After graft removal, the earliest time interval for revision procedure is was 6 to 9 months. Since there was no consensus in the literature, Matava et al concluded that primary infection can be treated with culture-specific intravenous antibiotics and surgical joint irrigation with graft retention as the initial treatment.

Outcomes after infection

The outcomes after deep infection after ACL reconstruction are mixed. Complications include pain, stiffness, arthrofibrosis, articular cartilage degeneration, and graft weakening or failure. While patients can generally perform pain-free activities of daily living, knee function following infection is impaired and results are much less satisfactory than for patients without postoperative infections (49). A full return to athletic activities was not certain, and pain followed by arthrofibrosis was the most common causes of unsatisfactory results (49). In 13 knees, of which only two had positive cultures, all treated with antibiotics initially, six patients failed to improve underwent arthroscopic irrigation and debridement (86). Of these 13, four had pain with stair climbing, three had slight impairment with squatting exercises, and three anterior knee pain with activities of daily living. In contrast, clinical outcomes inferior to control subjects without infection appeared to be secondary to damage to the articular cartilage from the infection (59). Four infections with the most significant functional limitations had severe bicondylar focal articular surface irregularities at MRI final follow-up. These patients averaged 12.25 days of inpatient hospital stay for these infections.

Recently, a long-term follow-up after infection on four of 831 patients that developed septic arthritis postoperatively required an average of 2.75 additional procedures for eradication (73). At 17.9 year average follow-up each patient had a decline in SF-36, Lysholm, and IKDC scores, and increase in KT-1000 displacement. Radiographic and MRI studies showed progression of arthritis in all patients, as compared to their 36 month follow up.

Conclusion

Patients should be educated about the signs of infection, and surgeons should always err on the side of caution to provide early treatment. Management with antibiotics and debridement with graft retention is usually the recommended treatment. While the diagnosis of an infection after reconstruction has significant morbidity, eradication of infection can usually be attained and patients may still have a functional knee with a much higher likelihood of osteoarthritis.

References:

1. Armstrong RW, Bolding F. Septic arthritis after arthroscopy: the contributing roles of intra-articular steroids and environmental factors. *Am J Infect Control* 1994;22:16-18.
2. Armstrong RW, Bolding F, Joseph R. Septic arthritis following arthroscopy: Clinical syndromes and analysis of risk factors. *Arthroscopy* 1992; 8:213-223.
3. Azar FM, Arthur ST. Complications of anterior cruciate ligament reconstruction. *Tech Knee Surg* 2004; 3:238-250.
4. Babcock HM, Carroll C, Matava M, L'ecuyer P, Praser V. Surgical site infections after arthroscopy: Outbreak investigation and case control study. *Arthroscopy* 2003;19:172-181.
5. Barber AF, McGuire DA, Johnson DH. Should allografts be used for routine anterior cruciate ligament reconstructions? *J Arthrosc Rel Surg* 2003;19(4):421-425.

6. Barker JU, Drakos MC, Maak TG, Warren RF, Williams III RJ, Allen AA. Effect of graft selection on the incidence of postoperative infection in anterior cruciate ligament reconstruction. *Am J Sports Med* 2010;38 (2):281-286.
7. Barrett GR, Lubert K, Replogle WH, Manley JL. Allograft anterior cruciate ligament reconstruction in the young, active patient; Tegner activity level and failure rate. *Arthroscopy* 2010;26:1593-1601.
8. Barrett AM, Craft JA, Replogle WH, Hydrick JM, Barrett GR. Anterior cruciate ligament graft failure: a comparison of graft type based on age and Tegner activity level. *Am J Sports Med* 2011;39:2194-2198.
9. Barrett GR, Field LD. Comparison of patella tendon versus patella tendon/Kennedy ligament augmentation device for anterior cruciate ligament reconstruction: Study of results, morbidity and complications. *Arthroscopy* 1993;9:624-632.
10. Binnet MS, Basahir K. Risk and outcome of infection after different arthroscopic anterior cruciate ligament reconstruction techniques. *Arthroscopy* 2007;23:862-868
11. Bear GS, Harner CD. Clinical outcomes of allograft versus autograft in anterior cruciate ligament reconstruction. *Clin Sports Med* 2007;26:661-681.
12. Berg EE. Tibial bone plug nonunion: a cause of anterior cruciate ligament reconstructive failure. *Arthroscopy* 1992;8:380-4.
13. Beynon BD, Uh BS, Johnson RJ et al. Rehabilitation after anterior cruciate ligament reconstruction: a prospective, randomized double-blind comparison of programs administered over 2 different time intervals. *Am J Sports Med* 2005;33:347-359.
14. Beynon BD, Johnson RJ, Abate JA, Fleming BC, Nichols CE. Treatment of anterior cruciate ligament injuries, part 1. *Am J Sports Med* 2005;33:1579-1602

15. Beynon BD, Johnson RJ, Abate JA, Fleming BC, Nichols CE. Treatment of anterior cruciate ligament injuries, part 2. *Am J Sports Med* 2005;33:1751-1767.
16. Borchers JR, Pedroza A, Kaeding C. Activity level and graft type as risk factors for anterior cruciate ligament graft failure: a case-control study. *Am J Sports Med* 2009;37:2362-2367.
17. Boyce T, Edwards J, Scarborough N. Allograft bone. The influence of processing on safety and performance. *Orthop Clin North Am.* 1999;30:571-581.
18. Brown CH, Carson EW. Revision anterior cruciate ligament surgery. *Clin Sports Med* 1999;18:109-171
19. Buck BE, Resnick L, Shah SM, Malinin TI. Human immunodeficiency virus cultured from bone. Implications for transplantation. *Clin Orthop Relat Res.* 1990;251:249-53.
20. Burks R, Freiderichs M, Fink B, Luker M, West H, Greis P. Treatment of postoperative anterior cruciate ligament infections with graft removal and early reimplantation. *Am J Sports Med* 2003;31:414-418.
21. Carey JL, Dunn WR, Dahm DL, Zeger SL, Spindler KP. A systemic review of anterior cruciate ligament reconstruction with autograft compared with allograft. *JBJS* 2009;91:2242-2250.
22. Centeno, JM, Woolf S, Reid JB, Lubowitz JH. Do anterior cruciate ligament allograft culture results correlate with clinical infections? *Arthroscopy* 2007;23(10):1100-1103
23. Centers for Disease Control and prevention. Septic arthritis following anterior cruciate ligament reconstruction using tendon allografts- Florida and Louisiana, 2000. *MMWR Morb Mortal Wkly Rep* 2001;50:1081-1083.

24. Centers for Disease Control and prevention. Update: Allograft associated bacterial infection- United States 2002. *MMWR Morb Mortal Wkly Rep* 2002;51(10):207-210.
25. Centers for Disease Control and Prevention (CDC). Hepatitis C virus transmission from an antibody-negative organ and tissue donor—United States, 2000-2002. *MMWR Morb Mortal Wkly Rep*. 2003;52:273-4, 276.
26. Clancy WG, Nelson DA, Reider B, et al. Anterior cruciate ligament reconstruction using one-third of the patellar ligament augmented by extra-articular tendon transfers. *J Bone Joint Surg Am* 1982;64:352-359.
27. Cohen SB, Sekiya JK. Allograft safety in anterior cruciate ligament reconstruction. *Clin Sports Med*. 2007 Oct;26(4):597-605.
28. Cohen SB, Yucha DT, Ciccotti MC, Goldstein DT, Ciccotti MA, Ciccotti MG. Factors affecting patient selection of graft type in anterior cruciate ligament reconstruction. *Arthroscopy* 2009;25(9):1006-1010.
29. Conrad EU, Gretch DR, Obermeyer KR, et al. Transmission of the hepatitis-C virus by tissue transplantation. *J Bone Joint Surg* 1995;77A:214-24.
30. Crawford C, Kainer M, Jernigan D, et al. Investigation of postoperative allograft-associated infection in patients who underwent musculoskeletal allograft implantation. *Clin Infect Dis* 2005;41:195-200.
31. Diaz-de-Rada P, Barriga A, Barrosos JL, Garcia-Barrecheguren E, Alfanso M, Valenti JR. Positive cultures in allograft ACL-reconstruction: what to do? *Knee Surg Sports Traumatol Arthrosc* 2003;11:219-22

32. Edgar CM, Zimmer S, Kakar S, Jones H, Schepsis AA. Prospective comparison of auto and allograft hamstring tendon constructs for ACL reconstruction. *Clin Orthop Relat Res.* 2008;466(9):2238-2246.
33. Fong SY, Tan JL. Septic arthritis after arthroscopic anterior cruciate ligament reconstruction. *Ann Acad Med Singapore.* 2004;33:228-234.
34. Garras DN, Shaner J, Lawson K, Cohen SB. Comparison of infection rates in allograft versus autograft ACL reconstructions. Arthroscopy Association North America Annual Meeting, E-poster presentation, May 2012. Orlando, FL.
35. Gillquist J, Messner K. Anterior cruciate ligament reconstruction and the long-term incidence of gonarthrosis. *Sports Med* 1999;27:143-156.
36. Greenberg DD, Robertson M, Vallurupalli S, White RA, Allen WC. Allograft compared with autograft infection rates in primary anterior cruciate ligament reconstruction. *J Bone Joint Surg Am* 2010;92(14):2402-8.
37. Guelich DR, Lowe WR, Wilson B. The Routine culture of allograft tissue in anterior cruciate ligament reconstruction. *Am J Sports Med* 2007;35(9):1495-99.
38. Harner CD, Olson E, Irrgang JJ, et al. Allograft versus autograft anterior cruciate ligament reconstruction: 3-to 5-year outcome. *ClinOrthop*1996;(324):134-44
39. Harner CD, Poehling GG. Double bundle or double trouble? *Arthroscopy* 2004;20(10)1013-4.
40. Harner CD, Giffin JR, Dunteman RC, et al. Evaluation and treatment of recurrent instability after anterior cruciate ligament reconstruction. *Instr Course Lect* 2001;50:463-74.

41. Harner CD, Irrgang JJ, Paul J, Dearwater S, Ru FH. Loss of motion after anterior cruciate ligament reconstruction. *Am J Sports Med* 1992;20:499-506.
42. Hirshman HP, Daniel DM, Miysaka K. The fate of unoperated knee ligament injuries. In: Daniel DM, Akeson W, O'Connor J, editors. *Knee ligaments: structure, function, injury and repair*. New York: Raven Press, 1990: 481-503.
43. Howe JG, Johnson RJ, Kaplan MJ, et al. Anterior cruciate ligament reconstruction using quadriceps patellar tendon graft. Part I. Long-term follow-up. *Am J Sports Med* 1991;19:447-457.
44. Indelli P, Dillingham M, Fanton G, Schurman D. Septic arthritis in postoperative anterior cruciate ligament reconstruction. *Clin Orthop* 2002;398:182-188.
45. Johnson RJ, Eriksson E, Haggmark T, et al. Five –to ten-year follow-up evaluation after reconstruction of the anterior cruciate ligament. *Clin Orthop* 1984;183:122-140.
46. Jones KG. Reconstruction of the anterior cruciate ligament using the central one-third of the patellar ligament. A follow-up report. *J Bone and Joint Surg Am* 1970;52: 1302-1308
47. Jost PW, Dy CJ, Robertson CM, Kelly AM. Allograft use in anterior cruciate ligament reconstruction. *HSSJ* 2011;7:251-256.
48. Joyce MJ. Safety and FDA regulations for musculoskeletal allografts: perspectives of an orthopaedic surgeon. *Clin Orthop* 2005;435:22-30.
49. Judd D, Bottoni C, Kim D, Burke M, Hooker S. Infections following anterior cruciate ligament reconstruction. *Arthroscopy* 2006;22:375–384
50. Katz LM, Battaglia TC, Patino P, Reichmann W, Hunter DJ, Richmond JC. A retrospective comparison of the incidence of bacterial infection following anterior

cruciate ligament reconstruction with autograft versus allograft. *Arthroscopy* 2008;24:1330-1335.

51. Keays SL, Newcombe PA, Bullock-Saxton JE, Bullock MI, Keays AC. Factors involved in the development of osteoarthritis following anterior cruciate ligament surgery. *Am J Sports Med* 2010;38:455-463.
52. Kleipool AE, Zijl JA, Willems WJ. Arthroscopic anterior cruciate ligament reconstruction with bone-patellar tendon-bone allograft or autograft. A prospective study with an average follow up of 4 years. *Knee Surg Sports Traumatol Arthrosc*. 1998;6(4):224-230.
53. Krych AJ, Jackson JD, Hoskin TL, Dahm DL. A meta-analysis of patellar tendon autograft versus patellar tendon allograft in anterior cruciate ligament reconstruction. *Arthroscopy* 2008;24:292-298.
54. Kurokouchi K, Takahashi S, Yamada T, Yamamoto H. Methicillin-resistant staphylococcus aureus-induced septic arthritis after anterior cruciate ligament reconstruction. *Arthroscopy* 2008;24 (5):615-617
55. Leal-Blanquet J, Alentorn-Geli E, Tuneu J, Valenti JR, Maestro A. Anterior cruciate ligament reconstruction: a multicenter prospective cohort study evaluating 3 different grafts using same bone drilling method. *Clin J Sport Med* 2011;21:294-300.
56. Margheritini F, Camillieri G, Mancini L, Mariani PP. C-reactive protein and erythrocyte sedimentation rate changes following arthroscopically assisted anterior cruciate ligament reconstruction. *Knee Surg, Sports Traumatol, Arthrosc* 2001;9:343-345.

57. Marzo JM, Bowen MK, Warren RF, Wickiewicz TL, Altchek DW. Intra-articular fibrous nodule as a cause of loss of extension following anterior cruciate ligament reconstruction. *Arthroscopy* 1992;8:10-18.
58. Matawa MJ, Evans TA, Wright RW, Shively RA. Septic arthritis of the knee following anterior cruciate ligament reconstruction: results of a survey of sports medicine fellowship directors. *Arthroscopy* 1998;14:717-725
59. McAllister DR, Parker RD, Cooper AE, Recht MP, Abate J. Outcomes of postoperative septic arthritis after anterior cruciate ligament reconstruction. *Am J Sports Med* 1999;27:562-570
60. Mehta VM, Mandala C, Foster D, Petsche TS. Comparison of revision rates in bone-patella tendon-bone autograft and allograft anterior cruciate ligament reconstruction. *Orthopedics* 2010;33:12
61. Mei-Dan O, Mann G, Steinbacher G, Ballester SJ, Cugat RB, Alvarez PD. Septic arthritis with *Staphylococcus lugdunensis* following arthroscopic ACL revision with BPTB allograft. *Knee Surg Sports Traumatol Arthrosc* 2008;16:15-18.
62. Montgomery SC, Campbell J. Septic arthritis following arthroscopy and intra-articular steroids. *J Bone Joint Surg Br* 1989;71:540
63. Muscolo DL, Carbo L, Aponte-Tinao LA, Ayerza MA, Makino A. Massive bone loss from fungal infection after anterior cruciate ligament arthroscopic reconstruction. *Clin Orthop Relat Res* 2009;467:2420-2425.
64. Nag HL, Neogi DS, Nataraj AR, Ajay KV, Yadav CS, Singh U. Tubercular infection after arthroscopic anterior cruciate ligament reconstruction. *J Arthrosc Rel Surg* 2009;25(2):131-136.

65. Noyes FR, Barber S. The effect of an extra-articular procedure on allograft reconstruction for chronic rupture of the anterior cruciate ligament. *J Bone Joint Surg Am* 1991;73:882-892
66. Oh HL, Chen DB, Seeto BG, MacDessi SJ. Mycobacterium fortium infection after anterior cruciate ligament reconstruction using a polylactic acid bioabsorbable screw: Case report. *The Knee* 2010;17:176-178.
67. Olsen EJ. Use of soft tissue allografts in sports medicine. *Advances Operative Orthopaedics* 1993;1:111-28
68. Papakonstantinou O, Chung C, Chanchairujira K, Resnick D. Complications of anterior cruciate ligament reconstruction: MR imaging. *EurRadiol* 2003;13:1106-1117
69. Pierre PS. Complications of anterior cruciate ligament reconstruction. *Sports Med Arthrosc Rev* 2004; 12:185-195
70. Poehling GG, Curl WW, Lee CA, et al. Analysis of outcomes of anterior cruciate ligament repair with 5-year follow-up: allograft versus autograft. *Arthroscopy* 2005;21:774-85.
71. Prodromos C, Joyce B, Shi K. A meta-analysis of stability of autografts compared to allografts after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2007;15:851-856.
72. Schollin-Borg M, Michaelsson K, Rahme H. Presentation, outcome and cause of septic arthritis after anterior cruciate ligament reconstruction: a case control study. *Arthroscopy* 2003;19:941-947

73. Schub DL, Schmitz LM, Sakamoto FA, Winalski CS, Parker RD. Long-term outcomes of postoperative septic arthritis after anterior cruciate ligament reconstruction. *Am J Sports Med* 2012;40(12):2764-2770.
74. Schulz AP, Gotze S, Schmidt HGK, Jurgens C, Faschingbauer M. Septic arthritis of the knee after anterior cruciate ligament surgery: a stage-adapted treatment regimen. *Am J Sports Med* 2007;35:1064–1069
75. Shelton WR, Papendick L, Dukes AD. Autograft versus allograft anterior cruciate reconstruction. *Arthroscopy*. 1997;13(4):446-449
76. Shelton WR, Treacy SH, Dukes AD, Bomboy AL. Use of allografts in knee reconstruction: I. Basic science aspects and current status. *J Am Acad Orthop Surg*. 1998;6:165-8.
77. Simonds, Holmberg SD, Hurwitz RL, Coleman TR, Bottenfield S, Conley LJ, Kohlenberg SH, Castro KG, Dahan BA, Schable CA, et al. Transmission of human immunodeficiency virus type 1 from a seronegative organ and tissue donor. *N Engl J Med* 1992;326:726-32.
78. Smith RL, Schurman DJ, Kajiyama BA, Mell M, Gilkerson E. The effect of antibiotics on the destruction of cartilage in experimental infectious arthritis. *J Bone Joint Surg Am* 1987;69:1063-1068.
79. Spindler KP, Wright RW. Clinical Practice. Anterior cruciate ligament tear. *N Engl J Med*. 2008 Nov 13;359(20):2135-42.
80. Strickland SM, MacGillivray JD, Warren RF. Anterior cruciate ligament reconstruction with allograft tendons. *Orthop Clin N Am* 2003;34:41-47.

81. Sun K, Tian SQ, Zhang JH, Xia CS, Zhang CL, Yu TB. Anterior cruciate ligament reconstruction with bone-patellar tendon-bone autograft versus allograft. *Arthroscopy* 2009;25:750-759.
82. Tomford WW. Transmission of disease through transplantation of musculoskeletal allografts. *J Bone Joint Surg Am* 1995; 77:1742-1754.
83. Tuman J, Diduch DR, Baumfield JA, Rubino LJ, Hart JM. Joint infection unique to hamstring tendon harvester used during anterior cruciate ligament reconstruction surgery. *Arthroscopy* 2008;24:618-620.
84. Vallianatos PG, Tilentzoglou AC, Koutsoukou AD. Septic arthritis caused by *Erysipelothrix rhusiopathiae* infection after arthroscopically assisted anterior cruciate ligament reconstruction. *Arthroscopy* 2003;19(3):1-4.
85. Van Tongel A, Stuyck J, Bellemans J, Vandenneucker H. Septic arthritis after arthroscopic anterior cruciate ligament reconstruction: A retrospective analysis of incidence management and outcome. *Am J Sports Med* 2007;35:1059-1063.
86. Viola R, Marzano N, Vianello R. An unusual epidemic: staphylococcus-negative infections involving anterior cruciate ligament reconstruction with salvage of the graft and function. *Arthroscopy* 2000;16:173-177
87. Wang C, Ao Y, Wang J, Hu Y, Cui G, Yu J. Septic arthritis after arthroscopic anterior cruciate ligament reconstruction: A retrospective analysis of incidence, presentation, treatment, and cause. *Arthroscopy* 2009; 25:243-249.
88. Williams RJIII, Laurencin CT, Warren RF, Speciale AC, Brause BD, O'Brien S. Septic arthritis after arthroscopic anterior cruciate ligament reconstruction. Diagnosis and management. *Am J Sports Med* 1997;25(2):261-7.

89. Zalavras CG, Patzakis MJ, Tibone J, Weisman N, Holtom P. Treatment of persistent infection after anterior cruciate ligament surgery. *Clin Orthop Rel Res* 2005;439:52-55.

Table 1. Infection rate by graft type

Study (year)	Autograft (infections / total)	Allograft (infections / total)	
Barker et al (2010)	12/177 = 0.44%	6/1349 = 0.68%	
Indelli et al (2002)	4/1400 = 0.29%	2/2100 = 0.10%	
Greenberg et al (2010)	0/221 = 0%	0/640 = 0%	
Garras et al (2012)	2/253 = 0.79%	4/535 = 0.74%	
Katz et al (2008)	2/170 = 1.2%	4/628 = 0.63%	
Wang et al (2009)	20/3978 = 0.50%	1/90 = 1.11%	
Crawford et al (2005)	0/41 = 0%	11/290 = 3.8%	
TOTAL	40/7840 = 0.51%	28/5632 = 0.49%	Overall: 68/13472 = 0.50%