

3-5-2018

TNF α induces Ca²⁺ influx to accelerate extrinsic apoptosis in hepatocellular carcinoma cells

Jianjun Zhu

Fourth Military Medical University (China); North Sichuan Medical College

Mingpeng Jin

Fourth Military Medical University (China)

Jiaojiao Wang

Fourth Military Medical University (China)

Hui Zhang

Fourth Military Medical University (China)

Yousheng Wu

Fourth Military Medical University (China)

Follow this and additional works at: <https://jdc.jefferson.edu/medoncfp>



Part of the [Oncology Commons](#)

See next page for additional authors

[Let us know how access to this document benefits you](#)

Recommended Citation

Zhu, Jianjun; Jin, Mingpeng; Wang, Jiaojiao; Zhang, Hui; Wu, Yousheng; Li, Deyang; Ji, Xiaoying; Yang, Hushan; Yin, Chun; Ren, Tingting; and Xing, Jinliang, "TNF α induces Ca²⁺ influx to accelerate extrinsic apoptosis in hepatocellular carcinoma cells" (2018). *Department of Medical Oncology Faculty Papers*. Paper 76.

<https://jdc.jefferson.edu/medoncfp/76>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Medical Oncology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Jianjun Zhu, Mingpeng Jin, Jiaojiao Wang, Hui Zhang, Yousheng Wu, Deyang Li, Xiaoying Ji, Hushan Yang, Chun Yin, Tingting Ren, and Jinliang Xing

RESEARCH

Open Access



TNF α induces Ca²⁺ influx to accelerate extrinsic apoptosis in hepatocellular carcinoma cells

Jianjun Zhu^{1,2}, Mingpeng Jin¹, Jiaojiao Wang¹, Hui Zhang³, Yousheng Wu¹, Deyang Li¹, Xiaoying Ji¹, Hushan Yang⁴, Chun Yin¹, Tingting Ren^{1*} and Jinliang Xing^{1*}

Abstract

Background: Tumor necrosis factor- α has been proven an effective anticancer agent in preclinical studies. However, the translation of TNF α from research to clinic has been blocked by significant systemic toxicity and limited efficacy at maximal tolerated dose, which need urgently to be solved.

Methods: The level of cytosolic Ca²⁺ was assessed by Fura-2 in HCC cells. After changing cytosolic Ca²⁺ level by using agonists or inhibitors, cell apoptosis was detected by flow cytometry. We also detected the effect of ionomycin or parvalbumin on the anti-tumor activity of TNF α in a mice model. Lastly, we studied the roles of cytosolic Ca²⁺ in the mitochondrial-dependent intrinsic apoptosis pathway.

Results: Here, we demonstrated that TNF α induced extracellular Ca²⁺ influx into cytoplasm through transient receptor potential channel in HCC cells. Both cytosolic Ca²⁺ scavenger and Ca²⁺-binding protein PV effectively desensitized hepatocellular carcinoma cells to TNF α , whereas combination ionomycin or 1,4,5-inositol triphosphate significantly sensitized HCC cells to TNF α , indicating that the increased level of cytosolic Ca²⁺ was positively correlated with the TNF α -induced cell apoptosis in vitro. In a nude mice xenograft model, our data revealed that TNF α combined with ionomycin remarkably synergized the anti-tumor effect of TNF α . Furthermore, we found that TNF α -mediated extracellular Ca²⁺ influx accelerated TNF α -induced extrinsic apoptosis through activating calpain/IAP/caspase3 pathway.

Conclusions: Our study provides the evidence supporting a novel mechanism by which TNF α induces extracellular Ca²⁺ influx to enhance cell apoptosis and suggests that increasing the level of cytosolic Ca²⁺ might be an alternative strategy to improve the pro-apoptotic activity of TNF α in HCC cells, although suitable chemical or biological reagents need to be further tested.

Keywords: Cell apoptosis, TNF α , Ca²⁺ influx, Hepatocellular carcinoma

Background

TNF α is a 23KD type II transmembrane protein, which is arranged in stable homotrimers. It is primarily produced by macrophages and a variety of other cells, including NK cells, T lymphocytes, smooth muscle cells, fibroblasts and others [1]. Many preclinical data suggest that TNF α may be used as a highly specific anti-cancer drug against many types of tumors [1, 2]. Moreover, recombinant human

TNF α (rhTNF α) has been tested as a systemic treatment of cancer patients in several phase I and phase II clinical trials [3]. However, the initial enthusiasm for the development of TNF α as a systemic treatment has waned when facing significant toxicities and a lack of evidence for therapeutic benefit [2]. How to increase the anti-cancer activity of TNF α at a low-dose condition is the key problem which needs to be solved urgently.

Ca²⁺, as a common signal transduction factor, has played many important roles in the process of cell division, growth, and death [4–6]. The cytosolic Ca²⁺ level is always risen during the process of cell apoptosis [7, 8]. Recent studies have revealed that TNF α is related with

* Correspondence: rtt419@163.com; rtt419@fmmu.edu.cn;

xingjl@fmmu.edu.cn

¹State Key Laboratory of Cancer Biology and Experimental Teaching Center of Basic Medicine, Fourth Military Medical University, 169 Changle West Road, Xi'an 710032, China

Full list of author information is available at the end of the article



remodeling of cytosolic Ca^{2+} homeostasis in a variety of human cells [9, 10]. For example, a study with human pulmonary artery endothelial cells (HPAEC) has reported that TNF α exposure significantly increases TRPC1 expression level and thrombin-induced Ca^{2+} influx in TNF α -stimulated HPAEC is two-fold greater than that in control cells [11]. Moreover, Wang GJ et al. have reported that TNF α induces the increases of cytosolic Ca^{2+} level, CaMKII δ B and CaN expression levels, and thus promotes cardiac hypertrophy [12]. More recently, another study has showed that TNF α induces cell death by the activation of transient receptor potential melastatin (TRPM2) to increase cytosolic Ca^{2+} level, followed by caspase activation and PARP cleavage [13]. Although some studies have reported that the increase of cytosolic Ca^{2+} level induced by TNF α plays important roles in varieties of physical activities, including cell death, it is still largely unknown about the functional roles and mechanisms underlying cytosolic Ca^{2+} in TNF α -induced cell death and whether the remodeling of cytosolic Ca^{2+} facilitates the pro-apoptotic effect of TNF α .

Here, we showed that TNF α induced extracellular Ca^{2+} influx in HCC cells. Importantly, the increased level of cytosolic Ca^{2+} mediated by TNF α was positively correlated with TNF α -induced cell apoptosis. The molecular mechanisms underlying the cytosolic Ca^{2+} in regulating TNF α -induced cell apoptosis were also deeply explored. Furthermore, combination with ionomycin was proven to be able to enhance the anti-cancer activity of TNF α in the mouse model.

Methods

Cell culture and public dataset collection

Human cell lines SNU739, SNU368, SNU354, SNU878, JHH-2, Huh-1, HLE, HLF, SMMC772, MHCC97H and QSG-7701 were routinely cultured. The authentication information of cell lines was provided in supplementary files. The detailed information about TCGA database was listed in Additional file 1: Table S3.

Knockdown, forced expression of target genes

Small interfering RNAs (siRNAs) were synthesized by GenePharma Company (Shanghai, China). The sequences of siRNA and primers were listed in Additional file 1: Table S2.

qRT-PCR and western blot

RNA extraction, cDNA synthesis and qPCR reactions were performed in Additional file 1: Supplementary Methods. Primers used in this study were listed in Additional file 1: Table S2. The western blot assay was performed as described in Additional file 1: Supplementary Methods. The primary antibodies used in this study and the working concentration were listed in Additional file 1: Table S1.

Measurement of cytosolic Ca^{2+}

Cells were loaded with Fura-2/AM (Invitrogen) for 30 min at 37 °C, and examined with a confocal laser scanning microscope FV1000 (Olympus, Tokyo, Japan). After 40 s of baseline recording, TNF α (100 ng/mL) was added where appropriate, and confocal images were recorded every 2 s.

Cell apoptosis assay

Cell apoptosis was measured by Annexin V-FITC detection Kit (BestBio, Shanghai, China) according to the manufacturer's protocol. The percentages of total apoptotic cells (both early and late) defined as the AnnexinV-FITC-positive fraction were determined.

TUNEL assay

Terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling (TUNEL) assay (Roche Applied Science, Rotkreuz, Switzerland) was performed to analyze cell apoptosis in xenograft tissues according to the manufacturer's protocol. Images of TUNEL/DAPI-stained sections were grabbed by a confocal laser scanning microscope FV1000 (Olympus). The apoptosis ratio was calculated as the percentage of both TUNEL and DAPI-positive nuclei after at least 500 cells were counted.

Calpain activity assay

Calpain activity was assayed using a fluorometric kit (Abcam, Cambridge, UK) according to the manufacturer's protocol.

Nude mice xenograft model

Nude mice xenograft model was used to assess in vivo tumor growth as described in Additional file 1: Supplementary Methods. TNF α (40 $\mu\text{g}/\text{Kg}$) only, or ionomycin (3 mg/Kg) only, or TNF α (40 $\mu\text{g}/\text{Kg}$) combined with ionomycin (3 mg/Kg) were administered by tail vein injection every three days with vehicle (40% (wt/vol.) of 2-hydroxypropyl- β -cyclodextrin for one month.

Immunoprecipitation

For immunoprecipitation experiments, total cell protein or synthesized TNF α were incubated with 200 μL protein A beads (BEAVER, Suzhou, China) supplemented with an antibody overnight at 4 °C. After washing the protein A beads, normalized amounts of total lysates or immunoprecipitated samples were analyzed by SDS-PAGE and western blot.

Detection of mitochondrial membrane potential

Mitochondrial membrane potential was measured using the fluorescence probe TMRM (Invitrogen) according to the manufacturer's protocol. Cells were incubated with 10 nM TMRM for 10 min at 37 °C in the dark and images were captured by laser confocal microscope (FV1000,

Olympus) and analyzed by ImagePro image analysis software (Media Cybernetics, Silver Spring, MD, USA).

Detection of cytochrome c release

Cytochrome c release was measured by immunofluorescence staining assay. Briefly, cells were incubated with MitoTracker Red (1 μ M) for 45 min at 37 $^{\circ}$ C in the dark. After washing the dyes, cell samples were fixed with 4% paraformaldehyde for 30 min, permeabilized with 0.2% Triton X-100 for 10 min, blocked by 2% BSA for 30 min and incubated with Cytochrome c antibody (1:100) overnight at 4 $^{\circ}$ C. Cell samples were then incubated with fluorophore-conjugated secondary antibody (1:200) and visualized by a confocal laser scanning microscope FV1000 (Olympus).

Statistical analysis

SPSS 17.0 software (SPSS, Chicago, IL) was used for all statistical analyses and *P* values less than 0.05 was considered to be statistically significant. Unpaired t-tests were used for comparisons between two groups where appropriate after checking for normal distribution and equal variance of the

data. One-way ANOVA were used for comparisons among three or more groups. Correlations between measured variables were tested by Spearman's rank correlation analyses.

Results

TNF α induces extracellular Ca $^{2+}$ influx into HCC cells

We firstly analyzed the level of cytosolic Ca $^{2+}$ after TNF α treatment in SNU739 and HLF HCC cells, and found that fluorescence intensity of cytosolic Ca $^{2+}$ indicator Fura-2 was obviously increased after TNF α treatment, which presented a dose-dependent manner (Fig. 1a-d). In contrast, Fura-2 fluorescence was not changed after TNF α treatment when HCC cells were cultured in calcium-free medium (Fig. 1e, f), which indicated that TNF α induced extracellular Ca $^{2+}$ influx into HCC cells.

Ca $^{2+}$ influx induced by TNF α is mediated by TRP channel and independent of TNFR

To explore whether tumor necrosis factor receptors (TNFRs) participated in TNF α -mediated Ca $^{2+}$ influx in HCC cells, the expression of TNFR was measured at





Fig. 1 TNF α induces Ca $^{2+}$ influx in HCC cells. **a** and **b** Confocal microscope analysis of [Ca $^{2+}$] $_c$ using fluorescent probe Fura-2/AM in SNU739 and HLF cells with treatment as indicated. (Arrow: cells treated with TNF α or PBS). **c** and **d** Quantitative analysis of the maximal increased level of cytosolic Ca $^{2+}$ after TNF α treatment (**d**) and (**f**) Confocal microscope analysis of [Ca $^{2+}$] $_c$ in SNU739 and HLF cells with treatment as indicated before stimulation of 100 ng/mL TNF α . HBSS (Ca $^{2+}$ free): cells cultured in Ca $^{2+}$ free-HBSS before TNF α stimulation; HBSS (1.3 mM Ca $^{2+}$): cells cultured in HBSS containing 1.3 mM Ca $^{2+}$ before TNF α stimulation. Data were shown as mean \pm SD. All experiments were performed at least three times. * *P* < 0.05; ** *P* < 0.01

both the mRNA and protein level by real-time PCR and Western Blot, respectively. We found that TNFR1 but not TNFR2 was expressed in HCC cells (Additional file 2: Figure S1a, b). Moreover, we successfully silenced TNFR1 expression by siRNA in SNU739 and HLF cells, which was verified by real-time PCR and Western Blot. Our data showed that TNFR1 knockdown had no effect on the expression of TNFR2 at both the mRNA and protein levels in HCC cells, suggesting that a compensatory positive regulation of TNFR-2 expression may be excluded. We further confirmed that TRADD, which is a direct downstream molecular of TNFR1 and functions to transfer cell death signal after TNF α stimulation [14], was not able to effectively interact with TNFR1 after siTNFR1 treatment.

These data further demonstrated that TNFR1 was successfully knocked down and TNFR1-induced classical extrinsic pathway was inactivated (Additional file 2: Figure S1c-e). Furthermore, our data indicated that the expression level of TNFR1 had no effect on the TNF α -mediated Ca²⁺ influx in HCC cells (Fig. 2a). These results indicate that TNFR pathway is not involved in the process of TNF α -mediated Ca²⁺ influx in HCC cells.

In recently years, 4 kinds of calcium channels in mammal cells have been identified, including voltage-gated calcium channels (VGCC), ligand-gated calcium channels (LGCC), transient receptor potential (TRP), and store-operated calcium channels (SOCE). As LGCCs are only expressed in excitable cells, and our data in Fig. 1e and f



also showed that TNF α had no effect on the level of cytosolic Ca²⁺ in cells cultured in calcium-free medium, so the effects of LGCCs and SOCEs on TNF α -mediated Ca²⁺ influx in HCC cells were ruled out. We then explored whether VGCC and TRP channels participated in the process of TNF α -mediated Ca²⁺ influx in HCC cells. Our data showed that the extracellular Ca²⁺ influx in HCC cells was significantly inhibited by both two TRP channel blockers, CAI and SKF96365, whereas the VGCC channel blockers, verapamil and diltiazem had no effect on the TNF α -mediated Ca²⁺ influx in HCC cells (Fig. 2b, c), suggesting that TRP channels may participate in the process of TNF α -mediated Ca²⁺ influx in HCC cells.

It has been reported that a series of TRP channels, including TRPC1, TRPC6, TRPV5, TRPV6, TRPM3, TRPM7, and TRPP5, are implicated in cell apoptosis process [13–15]. Our results demonstrated that TRPM7 was the main channel type expressed in HCC cells (Fig. 2d), which was also demonstrated by bioinformatic analysis based on public transcriptome sequencing data from The Cancer Genome Atlas (TCGA) database (Additional file 2:

Figure S1 h). Moreover, we found that TRPM7 knockdown significantly inhibited TNF α -mediated Ca²⁺ influx in HCC cells (Fig. 2e, Additional file 2: Figure S1f, g). Furthermore, co-immunoprecipitation assay showed that TNF α did not directly interact with TRPM7 in SNU739 and HLF cells (Additional file 2: Figure S1i). These findings strongly suggest that TNF α -mediated Ca²⁺ influx may be mediated by TRPM7 and independent of TNFR.

Decreased cytosolic Ca²⁺ level attenuates TNF α -induced apoptosis

We next investigated the roles of TNF α -mediated Ca²⁺ influx in the apoptosis of HCC cells. As shown in Additional file 3: Figure S2a and c, both cytosolic Ca²⁺ scavenger (BAPTA-AM) and Ca²⁺-binding protein parvalbumin (PV) efficiently reduced the level of [Ca²⁺]_c after TNF α treatment in HCC cells. We further found that either Ca²⁺ scavengers or PV reduced the percentage of TNF α -induced apoptotic HCC cells (Fig. 3a, b, Additional file 3: Figure S2b, d). We then investigated whether TRP channels mediated the effects of TNF α -mediated Ca²⁺ influx on cell




Fig. 3 Decreased cytosolic Ca²⁺ level attenuates TNF α -induced apoptosis of HCC cells. **a-d** Cell apoptosis analysis by flow cytometry 24 h after treatment as indicated before stimulation of TNF α (100 ng/mL). BAPTA-AM: 10 μ M; CAI: 10 μ M; SKF96365: 100 μ M; PV: expression vector encoding parvalbumin; siTRPM7: siRNA against TRPM7. **e** Tumor growth curves of subcutaneous xenograft tumor model developed from stable HCC cell lines (Right); dissected tumors from sacrificed mice (Left) were shown (*n* = 4 for each group). EV: xenografts developed from SNU-739 cells transfected with empty vector; PV, xenografts developed from SNU-739 cells stably forced expressing parvalbumin; TNF α : 40 μ g/Kg. **f** TUNEL staining in tumor tissues of nude mice xenograft model with treatment as indicated. Blue: DAPI; Green: TUNEL positive nucleus. Quantification of cell death ratios were shown in the right. Scale Bar, 50 μ m. Data were shown as mean \pm SD. EV: empty vector; PV: parvalbumin. All experiments were performed at least three times. * *P* < 0.05; ** *P* < 0.01

apoptosis. Our data showed that the apoptotic rates of SNU739 and HLF cells were significantly decreased after treatment of TRP channel blocker, CAI or SKF96365 (Fig. 3c, Additional file 3: Figure S2f), and the same effect was also observed after knockdown of TRPM7 in SNU739 and HLF cells (Fig. 3d, Additional file 3: Figure S2g). Taken together, our results demonstrate that inhibiting the elevation of TNF α -mediated cytosolic Ca²⁺ attenuates TNF α -induced apoptosis *in vitro*.

To evaluate the role of TNF α -mediated Ca²⁺ influx in tumor growth *in vivo*, we established the subcutaneous nude mice model. The mice treated with TNF α exhibited significantly decreased growth capacity than those in the control group (Fig. 3e). Moreover, our results also indicated that the growth capacity of xenograft tumors developed from SNU739-PV cells were significantly enhanced than xenograft tumors developed from SNU739-EV cells with TNF α treatment. These results strongly suggest that PV protein may attenuate TNF α -induced cell apoptosis via buffering the TNF α -mediated Ca²⁺ influx in HCC cells. We further confirmed the effect of PV on TNF α -mediated cell apoptosis in xenograft tumor tissues using TUNEL staining (Fig. 3f). These *in vivo* data support the evidence that extracellular Ca²⁺ influx contributes to the TNF α -induced HCC cell apoptosis.

Increased cytosolic Ca²⁺ sensitizes HCC cells to TNF α -induced apoptosis

Furthermore, we investigated whether increased extracellular Ca²⁺ influx enhanced TNF α -induced apoptosis in HCC cells. Ionomycin (an ionophore with high affinity to calcium) and IP₃ (an agonist to promote calcium releasing from ER) were used to rapidly elevate [Ca²⁺]_c as described previously [16, 17]. As shown in Additional file 4: Figure S3a and d, the optimal concentrations of ionomycin and TNF α were arbitrarily determined as 1 μ M and 100 ng/mL, respectively, based on apoptosis rate of less than 10% in normal hepatic cell QSG-7701. Our results showed that ionomycin (1 μ M) treatment remarkably increased the level of [Ca²⁺]_c in HLF and QSG-7701 cells after TNF α treatment (Additional file 4: Figure S3b). The similar effects were observed in HLF and QSG-7701 cells after IP₃ (10 μ M) treatment (Additional file 4: Figure S3c). Our data further demonstrated that TNF α -induced cell apoptosis was significantly increased at a dose-response manner in HLF and QSG-7701 cells with treatment of either 1 μ M ionomycin or 10 μ M IP₃ (Fig. 4a, b, Additional file 4: Figure S3e, f). Taken together, these results indicate that the increased cytosolic Ca²⁺ sensitizes HCC cells and normal hepatocytes to TNF α -induced apoptosis. Moreover, our data indicated that the treatment with a combination of 100 ng/mL TNF α and 1 μ M ionomycin exhibited an apoptosis percentage of about 37% in HCC

cell HLF, but only about 14% in normal hepatocyte QSG-7701, which is two times as treatment with TNF α (100 ng/mL) only (Additional file 4: Figure S3d, e). We have evaluated the toxicity of TNF α only and the combination of TNF α with ionomycin in normal hepatic cell QSG-7701. Our data showed that treatment with 100 ng/mL TNF α and a combination of 100 ng/mL TNF α and 1 μ M ionomycin exhibited an apoptosis percentage of 6.5% and 14.6%, respectively, in normal hepatic cell QSG-7701 (Additional file 4: Figure S3d, e), suggesting that the possible toxicity of TNF α may be increased by a combined use of ionomycin, although the pro-apoptotic effect can be improved.


To evaluate whether ionomycin could sensitize HCC cells to TNF α -induced apoptosis *in vivo*, we treated xenograft tumors developed from SNU739 cells with ionomycin and TNF α . As shown in Fig. 4c, a significant decrease of growth capacity was observed in xenograft tumors treated with the combination of TNF α (40 μ g/kg) and ionomycin (3 mg/kg) for one month when compared with xenograft tumors with treatment of TNF α (40 μ g/kg) alone. We further confirmed the effect of combined TNF α and ionomycin treatment on cell apoptosis in xenograft tumor tissues using TUNEL staining (Fig. 4d). These *in vivo* data further indicated that the increased Ca²⁺ influx sensitizes HCC cells to TNF α -mediated cell apoptosis and the induction of Ca²⁺ influx by ionomycin may synergize the apoptotic effect of TNF α .

TNF α -induced apoptosis is positively correlated with the level of extracellular calcium influx

We then investigated the relationship between TNF α -induced cell apoptosis and the level of extracellular Ca²⁺ influx in 10 HCC cell lines with TNF α treatment (Additional file 5: Figure S4a, b). Our results showed a different degree of TNF α -mediated extracellular Ca²⁺ influx in 10 HCC cells, suggesting that TNF α -mediated extracellular Ca²⁺ influx was a common phenomenon in HCC cells (Fig. 5a). Moreover, we found that TNF α -induced cell apoptosis was obviously positively correlated with the level of extracellular Ca²⁺ influx (Fig. 5b, c).

TNF α -mediated Ca²⁺ influx activates calpain signaling to enhance extrinsic apoptosis

Previous studies have demonstrated that calpain, a class of Ca²⁺-dependent cysteine proteases, and one of its main substrates the inhibitor apoptosis protein (IAP) plays critical roles in TNF α -induced extrinsic apoptotic pathway [18]. Therefore, we investigated the effect of TNF α -mediated extracellular Ca²⁺ influx on calpain/cIAP/caspase3 pathway. Western blot analysis showed that TNF α had no effect on the expression of calpain. However, the enzymatic activity of calpain was significantly increased in HCC cells with TNF α treatment when compared with control group



(Fig. 6a). Our data also showed that the expression of cIAP2 and XIAP was significantly decreased by $TNF\alpha$ treatment, whereas the expression of cIAP1 was not affected. In addition, the expression of the cleaved caspase3 was significantly increased by $TNF\alpha$ treatment (Fig. 6b).

To further investigated the $TNFR1$ -independent mechanism, we re-evaluated the effect of $TNF\alpha$ -mediated Ca^{2+} influx on $TNF\alpha$ -mediated cell apoptosis in HCC cells with $TNFR1$ knockdown. Moreover, the calpain activity was significantly inhibited by calpeptin (40 μM) treatment, whereas $TNFR1$ knockdown had no effect on the calpain activity, suggesting that the activation of calpain was independent of $TNFR1$ (Fig. 6c). In contrast, the expression levels of cIAP2 and XIAP were remarkably increased and the cleaved caspase 3 was decreased by $TNFR1$ knockdown in HCC cells (Fig. 6d). In addition, the suppression of calpain activity by calpeptin significantly increased the expression of cIAP2 and XIAP and decreased the expression of cleaved caspase 3 in HCC cells with $TNFR1$ knockdown (Fig. 6d). Besides, similar results were found after the decrease of cytosolic $[Ca^{2+}]$ through the forced

expression of PV (Fig. 6e, f). In contrast, ionomycin treatment, which can cause the increase of cytosolic $[Ca^{2+}]$, significantly inhibited the expression of cIAP2 and XIAP and increased the calpain activity and expression of cleaved caspase 3 in HCC cells with $TNFR1$ knockdown (Fig. 6g, h). Overall, these results clearly show that cytosolic Ca^{2+} participates in the process of $TNF\alpha$ -induced cell apoptosis through calpain/IAP/caspase3 pathway in HCC cells and demonstrate that $TNF\alpha$ -induced cell apoptosis is affected by both the expressions of $TNFR1$ and the levels of cytosolic Ca^{2+} in HCC cells. Furthermore, we have also investigated the effect of $TNF\alpha$ on the extracellular Ca^{2+} influx, TRPM7 activity, and activation of calpain signaling in normal hepatic cell QSG-7701. As shown in Additional file 6: Figure S5a, the cytosolic Ca^{2+} level was significantly increased after $TNF\alpha$ treatment in QSG-7701 cells. The extracellular Ca^{2+} influx was significantly inhibited after TRPM7 knockdown, indicating that $TNF\alpha$ -mediated Ca^{2+} influx is also dependent on TRPM7 channel in QSG-7701 cells (Additional file 6: Figure S5b, c). In addition, our results also showed that the activity of




Fig. 5 The level of extracellular calcium influx is positively correlated with TNF α -mediated apoptosis. **a** Apoptosis analysis by flow cytometry 24 h after treatment as indicated. **b** Confocal microscope analysis of $[Ca^{2+}]_i$ level using fluorescent probe Fura-2/AM in 10 kinds of HCC cells with treatment as indicated. **c** Correlational analysis of the maximal increased level of cytosolic Ca^{2+} and apoptotic rates in HCC cells after TNF α (100 ng/mL) treatment. Data were shown as mean \pm SD. All experiments were performed at least three times




Fig. 6 Ca^{2+} influx induced by TNF α activated calpain signaling to enhances extrinsic apoptosis. **a** Western Blot analysis for Calpain expression and analysis of Calpain activity after TNF α (100 ng/mL) stimulation for 4 h in HCC cells with treatment as indicated. **b** Western Blot analysis for expression of cIAP1, cIAP2, XIAP and cleaved caspase-3 after TNF α (100 ng/mL) treatment. **c-h** Analysis of Calpain activity and the expression of cIAP1, cIAP2, XIAP and cleaved caspase-3 after treatment with TNF α stimulation for 4 h in HCC cells with treatment as indicated. TNF α : 100 ng/mL; siATNFR1: siRNA target TNFR1; Calpeptin: 40 μ M; PV: HCC cells transfected with expression vector encoding parvalbumin, Iono: 1 μ M ionomycin. Data were shown as mean \pm SD. All experiments were performed at least three times. * $P < 0.05$; ** $P < 0.01$


calpain was significantly increased after TNF α treatment in normal hepatic cells (Additional file 6: Figure S5d).

TNF α -mediated Ca²⁺ influx has no effect on mitochondria-dependent intrinsic apoptosis

To investigate whether TNF α -mediated Ca²⁺influx was associated with mitochondrial-dependent intrinsic apoptosis, we evaluated the mitochondrial membrane potential (MMP), the subcellular location of cytochrome c (Cyto c), and the translocation of Bcl-2 family members in HCC cells after TNF α treatment. As shown in Fig. 7a, TNF α -mediated Ca²⁺ influx had no effect on the MMP of HCC cells. There was also no obvious release of Cyto c from mitochondria to cytoplasm (Fig. 7b), and no obvious translocation of Bax and Bak from the cytoplasm to mitochondria after TNF α treatment (Fig. 7c). Overall, these data indicate that cell apoptosis induced by TNF α -mediated Ca²⁺ influx is independent of intrinsic apoptosis pathway.

Discussion

TNF α has been proven to be an effective anticancer agent in a series of preclinical studies [2]. However, the promise of systemic TNF α has not translated to patient therapy and the enthusiasm had been cubed due to the toxicity profile and lack of efficacy at maximum tolerated dose (MTD) of the patients [2]. In the past decades, lots of researchers have focused on the modifications of TNF α structure so as to increase its anti-tumor activity, although these modifications are not remarkably effective [1]. Recently, TNF α has been described to be related with levels of cytosolic Ca²⁺ in various cell types [7, 19], although its effect remains largely unclear. In this study, we for the first time demonstrated that TNF α induced extracellular Ca²⁺ influx in HCC cells. Bellomo et al. have reported that TNF α induces a sustained increase in intracellular free Ca²⁺ concentration in mammary adenocarcinoma [19]. Chang et al. have reported that the combined treatment of TNF α and



IFN- γ significantly increases the cytosolic Ca^{2+} concentration in pancreatic β cell [7]. However, Carrasquel et al. have reported that TNF α increases the basal level of $[\text{Ca}^{2+}]_c$ after a Ca^{2+} pulse in human sperm [20]. Motagally et al. have reported that the incubation with TNF α decreases depolarization-induced Ca^{2+} influx in postganglionic sympathetic neurons [10]. Moreover, a recent report has provided further supporting evidence, showing that the level of cytosolic Ca^{2+} is decreased in cardiocytes after TNF α treatment [12]. These findings suggest that the level of cytosolic Ca^{2+} is regulated by TNF α , but the dual roles of TNF α in regulating the level of cytosolic Ca^{2+} may be context dependent or cell type specific, which needs more comprehensive investigation.

Furthermore, both TRP channel inhibitor and TRPM7 knockdown significantly inhibited TNF α -mediated Ca^{2+} influx in HCC cells, indicating that the activity of TRPM7 is regulated by TNF α . Our results further showed that TNF α did not directly interact with TRPM7, indicating that TNF α was not a direct activator of TRPM7 channel. Numata et al. have reported that TRPM7 channel is activated by membrane stretch which is produced by negative pressure (3 cm H_2O) or hypotonic solution [21]. Several organic molecules have also been identified as the activators of TRPM7, including naltriben, mibefradil and bradykinin [22, 23]. Furthermore, Desai et al. have found that TRPM7 channel is activated by caspase-dependent cleavage [24]. However, our results showed that TRPM7 channel was immediately activated upon TNF α stimulation and no direct interaction was observed between TNF α and TRPM7. Therefore, the exact mechanism underlying the activation of TRPM7 channel by TNF α needs to be explored in future study.

Changes in the levels of intracellular Ca^{2+} provide dynamic and highly versatile signals that control kinds of cellular processes, although their importance is perhaps most strikingly exemplified by their functional role in life-and-death decisions [5]. Accumulating evidence have demonstrated the increased levels of cytosolic Ca^{2+} plays a critical role in cell death [8, 19]. Bellomo et al. have reported that the increased intra-nuclear free Ca^{2+} induced by TNF α enhances the activity of Ca^{2+} -dependent endonuclease, resulting in DNA fragmentation and cell apoptosis [19]. In consistence with above-mentioned study, we found that the decreased cytosolic Ca^{2+} level attenuates TNF α -induced cell apoptosis, whereas the increased cytosolic Ca^{2+} sensitizes HCC cells to TNF α -induced apoptosis, strongly suggesting that TNF α -induced apoptosis may be positively correlated with the level of extracellular Ca^{2+} influx in HCC cells.

Lots of proteases have been identified as the downstream molecules activated by cytosolic Ca^{2+} to trigger cell death, such as calpain, calcineurin, and DAP kinase

[8, 25]. Calpain is a Ca^{2+} -activated cysteine protease localized to the cytosol and mitochondria, which has been shown to regulate apoptosis and necrosis [25]. It has been demonstrated that calpain mediates the cisplatin-induced apoptosis in human lung adenocarcinoma cells through truncating Bid to tBid and then inducing the mitochondrial apoptotic pathway [26]. Recently, more and more substrates have been identified to be hydrolyzed specifically by calpain during apoptosis, such as human DNA polymerase epsilon, cain/cabin1, fodrin, p53, caspase 7 and caspase 3 [26–30]. Moreover, inhibitor apoptosis proteins (IAPs) has been found to be hydrolyzed by calpain to promote TNF α -induced classical extrinsic apoptosis [3, 18]. In the present study, we first reported that cytosolic Ca^{2+} /calpain/IAPs pathway plays a critical role in synergizing the pro-apoptotic effect of TNF α .

It is generally accepted that mitochondrial Ca^{2+} uptake functions in cell proliferation [5]. However, excessive Ca^{2+} load to the mitochondria may induce apoptosis [8]. Accumulated Ca^{2+} within mitochondria regulates production of ATP, and activates of metabolism-related enzymes involved in cell proliferation [5]. In contrast, mitochondrial Ca^{2+} loading also causes PTP opening to irreversibly commit cells to death by causing IMM depolarization, matrix swelling, release of stored Ca^{2+} and apoptogenic proteins [8]. These findings highlight a dual role of mitochondrial Ca^{2+} in energy provision and induction of cell death, which depend on the amount of mitochondrial Ca^{2+} uptake [31]. Actually, our data showed that MMP, subcellular location of cytochrome c, and the translocation of Bcl-2 family proteins were not obviously changed by TNF α treatment, strongly suggesting that TNF α -induced extracellular Ca^{2+} influx may not trigger mitochondria-dependent cell death in HCC.

Ionomycin is antibiotic produced by *Streptomyces globatus*, which is characterized as a calcium ionophore used to increase the cytosolic Ca^{2+} concentration in numerous studies, but the usage of ionomycin in vivo is rare [32]. Recently, Zheng et al. has reported that ionomycin treatment effectively improves the hyperglycaemia and insulin resistance in the mouse model of diabetes [33]. In the present study, our data showed that the combination of TNF α with ionomycin significantly improves the pro-apoptotic effect in HCC cells, although the possible toxicity of TNF α may be also increased. In future study, suitable chemical or biological reagents that specifically increase cytosolic Ca^{2+} need to be further screened and tested for better combination effect with TNF α in HCC treatments.

Conclusion

In summary, our data provides the strong evidence to support the notion that extracellular Ca^{2+} influx induced by

TNF α facilitates the TNF α -mediated extrinsic apoptosis through activating calpain/IAPs/caspase3 signaling pathway and suggests that increasing the level of cytosolic Ca²⁺ might be an alternative strategy to improve the pro-apoptotic activity of TNF α in HCC cells, although suitable chemical or biological reagents need to be further tested.

Additional files

Additional file 1: Supplemental materials and Methods. **Table S1.** Primary antibodies used for western blot. **Table S2.** Sequence of primers and siRNA. **Table S3.** Public datasets used for bioinformatic analysis. (DOC 97 kb)

Additional file 2: Figure S1. (a) and (b) The relative mRNA and protein expression of TNFR1 and TNFR2 measured by qRT-PCR or Western Blot in SNU739 and HLF cells. (c) and (d) qRT-PCR and Western Blot analysis of TNFR1 and TNFR2 mRNA and protein expression levels in SNU739 and HLF cells transfected with siRNA as indicated. (e) TNFR1 protein expression were determined by Co-immunoprecipitation (Co-IP) and western blot in HCC cells as described. (f) and (g) qRT-PCR and western blot analysis of TRPM7 mRNA and protein expression levels in SNU739 and HLF cells transfected with siRNA as indicated. (h) The relative mRNA expression levels of TRPC1, TRPC6, TRPM2, TRPM3, TRPM7, TRPV4, TRPV5, TRPV6, TRPP2, and TRPP5 in HCC tumor tissues were analyzed in public microarray data TCGA downloaded from the Gene Expression Omnibus (GEO) database. (i) The interaction effects between TNF α and TRPM7 were determined by Co-immunoprecipitation (Co-IP) and western blot in HCC cells as described. Data were shown as mean \pm SD. All experiments were performed at least three times. * $P < 0.05$; ** $P < 0.01$. (JPEG 945 kb)

Additional file 3: Figure S2. Decreased cytosolic Ca²⁺ level attenuates TNF α -induced apoptosis of HCC cells. (a) Confocal microscope analysis of [Ca²⁺]_c level using fluorescent probe Fura-2/AM in SNU739 and HLF cells with treatment as indicated. BAPTA-AM: 10 μ M. (b) Cell apoptosis analysis by flow cytometry 24 h after treatment as indicated before TNF α (100 ng/mL) stimulation. BAPTA-AM: 10 μ M. (c) Confocal microscope analysis of [Ca²⁺]_c level using fluorescent probe Fura-2/AM in HCC cells with treatment as indicated. EV: cells transfected with the empty vector; PV-OE: cells stably forced expressing Parvalbumin protein. (d) Apoptosis analysis by flow cytometry 24 h after treatment as indicated. (e) Western Blot analysis for Parvalbumin expression in SNU739 and HLF cells with treatment as indicated. (f) Cell apoptosis analysis by flow cytometry 24 h after treatment as indicated. CAI: 10 μ M; SKF96365: 100 μ M. (g) Cell apoptosis analysis by flow cytometry 24 h after treatment as indicated. siTRPM7: siRNA against TRPM7; si Ctrl: negative control siRNA. Data were shown as mean \pm SD. All experiments were performed at least three times. * $P < 0.05$; ** $P < 0.01$. (ZIP 1986 kb)

Additional file 4: Figure S3. Cytosolic Ca²⁺ sensitized HCC cells to TNF α -induced apoptosis. (a) Apoptosis analysis by flow cytometry 24 h after treatment as indicated. (b) and (c) Confocal microscope analysis of [Ca²⁺]_c level using fluorescent probe Fura-2/AM in HLF and QSG-7701 cells with treatment as indicated. TNF α (50 ng/mL) + Iono: 50 ng/mL TNF α combined with 1 μ M ionomycin, TNF α (100 ng/mL) + Iono: 100 ng/mL TNF α combined with 1 μ M ionomycin, TNF α (200 ng/mL) + Iono: 200 ng/mL TNF α combined with 1 μ M ionomycin. TNF α (50 ng/mL) + IP₃: 50 ng/mL TNF α combined with 10 μ M IP₃; TNF α (100 ng/mL) + IP₃: 100 ng/mL TNF α combined with 10 μ M IP₃; TNF α (200 ng/mL) + IP₃: 200 ng/mL TNF α combined with 10 μ M IP₃; (d)-(f) Apoptosis analysis by flow cytometry 24 h after treatment as indicated. Data were shown as mean \pm SD. All experiments were performed at least three times. * $P < 0.05$; ** $P < 0.01$. (ZIP 2448 kb)

Additional file 5: Figure S4. The level of extracellular calcium influx is positively correlated with TNF α -mediated apoptosis. (a) Confocal microscope analysis of [Ca²⁺]_c level using fluorescent probe Fura-2/AM in 10 kinds of HCC cells with treatment as indicated. (b) Apoptosis analysis by flow cytometry 24 h after treatment as indicated. All experiments were performed at least three times. (ZIP 2184 kb)

Additional file 6: Figure S5. The role of TNF α -mediated Ca²⁺ influx in normal hepatic cells. (a) and (c) Confocal microscope analysis of [Ca²⁺]_c

level using fluorescent probe Fura-2/AM in QSG-7701 cells with treatment as indicated. TNF α : 100 ng/mL; siTRPM7: siRNA target TRPM7. (b) qRT-PCR and western blot analysis of TRPM7 mRNA and protein expression levels in QSG-7701 cells transfected with siRNA as indicated. (d) Analysis of Calpain activity after TNF α stimulation for 4 h in QSG-7701 cells with treatment as indicated. Data were shown as mean \pm SD. All experiments were performed at least three times. * $P < 0.05$; ** $P < 0.01$. (JPEG 664 kb)

Abbreviations

cIAP1/2: cellular inhibitor apoptosis protein 1/2; Cyto c: cytochrome c; HCC: hepatocellular carcinoma; IAP: inhibitor apoptosis protein; IFN- γ : interferon gamma; IHC: immunohistochemistry; IMM: inner mitochondrial membrane; LGCC: ligand-gated calcium channels; MMP: mitochondrial membrane potential; MTD: maximum tolerated dose; PARP: poly-ADP-ribose polymerase; PTP: permeability transition pore; PV: parvalbumin; SOCE: store-operated calcium channels; TNFR: tumor necrosis factor receptor; TNF α : tumor necrosis factor alpha; TRADD: tumor necrosis factor receptor type 1-associated death domain protein; TRP: transient receptor potential; TRPM7: transient receptor potential melastatin 7; VGCC: voltage-gated calcium channels; XIAP: X-linked inhibitor apoptosis protein

Acknowledgements

The authors thank Gang Wang for the help in animal study.

Ethical approval

The study was approved by the Ethics Committee of the Fourth Military Medical University. Animal research was approved by the Institutional Animal Care and Use Committee of Fourth Military Medical University.

Funding

This work was supported by National Natural Science Foundation of China [No. 8157272, and 81320108021], and The State Key Laboratory of Cancer Biology, [No. CBSKL20170].

Availability of data and materials

The datasets analyzed during the current study are available in the [TCGA] repository, [<http://cancergenome.nih.gov/>].

Authors' contributions

JZ performed most experiments, wrote the manuscript; JW and JM participated in the cell apoptosis analysis; MJ and HZ participated in the in vivo study; DL, XJ and YW performed data analysis; CY participated in the Co-IP assay; HY supervised the work and provided administrative support; JX and TR designed the overall study, supervised the experiments, analyzed the results, and wrote the paper. All authors read and approved the final manuscript.

Consent for publication

Not applicable

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹State Key Laboratory of Cancer Biology and Experimental Teaching Center of Basic Medicine, Fourth Military Medical University, 169 Changle West Road, Xi'an 710032, China. ²Department of Human Anatomy, Premedical College, North Sichuan Medical College, Nanchong 637000, China. ³Department of Pain Treatment, Tangdu Hospital, Fourth Military Medical University, Xian 710038, China. ⁴Division of Population Science, Department of Medical Oncology, Kimmel Cancer Center, Thomas Jefferson University, Philadelphia, PA 19107, USA.

Received: 28 November 2017 Accepted: 17 February 2018

Published online: 05 March 2018

References

- Anderson GM, Nakada MT, Dewitte M. Tumor necrosis factor- α in the pathogenesis and treatment of cancer. *Curr Opin Pharmacol*. 2004;4(4):314–20.
- Roberts NJ, Zhou S, Jr LAD, Holdhoff M. Systemic use of tumor necrosis factor alpha as an anticancer agent. *Oncotarget*. 2011;2(10):739–51.
- Mocellin S, Rossi CR, Pilati P, Nitti D. Tumor necrosis factor, cancer and anticancer therapy. *Cytokine Growth F R*. 2005;16(1):35–53.
- Kanno N, Glaser S, Chowdhury U, Phinizz JL, Baiocchi L, Francis H, Lesage G, Alpini G. Gastrin inhibits cholangiocarcinoma growth through increased apoptosis by activation of Ca²⁺-dependent protein kinase C- α . *J Hepatol*. 2001;34(2):284–91.
- Cook SJ. Ca²⁺ signalling checkpoints in cancer: remodelling Ca²⁺ for cancer cell proliferation and survival. *Nat Rev Cancer*. 2008;8(5):361.
- Stewart TA, Yapa KT, Monteith GR. Altered calcium signaling in cancer cells. *Biochim Biophys Acta*. 2015;1848(10):2502–11.
- Chang I, Cho N, Kim S, Kim JY, Kim E, Woo JE, Nam JH, Kim SJ, Lee MS. Role of calcium in pancreatic islet cell death by IFN- γ /TNF- α . *J Immunol*. 2004;172(11):7008.
- Hajnoczky G, Davies E, Madesh M. Calcium signaling and apoptosis. *Biochem Biophys Res Commun*. 2003;304(3):445.
- Cheng J, Tang W, Su Z, Guo J, Tong L, Wei Q. Calcineurin subunit B promotes TNF- α -induced apoptosis by binding to mitochondria and causing mitochondrial Ca²⁺ overload. *Cancer Lett*. 2012;321(2):169–78.
- Motagally MA, Lukewich MK, Chisholm SP, Neshat S, Lomax AE. Tumour necrosis factor alpha activates nuclear factor kappaB signalling to reduce N-type voltage-gated Ca²⁺ current in postganglionic sympathetic neurons. *J Physiol*. 2009;587(Pt 11):2623.
- Paria BC, Vogel SM, Ahmmed GU, Alamgir S, Shroff J, Malik AB, Tirupathi C. Tumor necrosis factor- α -induced TRPC1 expression amplifies store-operated Ca²⁺ influx and endothelial permeability. *Am J Physiol Lung Cell Mol Physiol*. 2004;287(6):1303–13.
- Wang GJ, Wang HX, Yao YS, Guo LY, Liu P. The role of Ca²⁺/calmodulin-dependent protein kinase II and calcineurin in TNF- α -induced myocardial hypertrophy. *Braz J Med Biol Res*. 2012;45(11):1045–51.
- Zhang W, Hirschler-Laszkiwicz I, Tong Q, Conrad K, Sun SC, Penn L, Barber DL, Stahl R, Carey DJ, Cheung JY. TRPM2 is an ion channel that modulates hematopoietic cell death through activation of caspases and PARP cleavage. *Am J Physiol Cell Physiol*. 2006;290(4):C1146.
- Shapovalov G, Lehen'kyi V, Skryma R, Prevarskaya N. TRP channels in cell survival and cell death in normal and transformed cells. *Cell Calcium*. 2011;50(3):295.
- Narayanan KL, Irmady K, Subramaniam S, Unsicker K, Von BUHO. Evidence that TRPC1 is involved in hippocampal glutamate-induced cell death. *Neurosci Lett*. 2008;446(2–3):117.
- Gao W, Liu J, Hu M, Huang M, Cai S, Zeng Z, Lin B, Cao X, Chen J, Zeng JZ. Regulation of proteolytic cleavage of retinoid X receptor- α by GSK-3 β . *Carcinogenesis*. 2013;34(6):1208–15.
- Taylor CW, Konieczny V. IP3 receptors: take four IP3 to open. *Sci Signal*. 2016;9(422):e1.
- Yeo JK, Cha SD, Cho CH, Kim SP, Cho JW, Baek WK, Suh MH, Kwon TK, Park JW, Suh SI. Se-methylselenocysteine induces apoptosis through caspase activation and Bax cleavage mediated by calpain in SKOV-3 ovarian cancer cells. *Cancer Lett*. 2002;182(1):83–92.
- Bellomo G, Perotti M, Taddei F, Mirabelli F, Finardi G, Nicotera P, Orrenius S. Tumor necrosis factor alpha induces apoptosis in mammary adenocarcinoma cells by an increase in intranuclear free Ca²⁺ concentration and DNA fragmentation. *Cancer Res*. 1992;52(5):1342.
- Carrasquel G, Camejo MI, Michelangeli F, Ruiz MC. Effect of tumor necrosis factor- α on the intracellular Ca²⁺ homeostasis in human sperm. *Am J Reprod Immunol*. 2013;70(2):153–61.
- Numata T, Shimizu T, Okada Y. TRPM7 is a stretch- and swelling-activated cation channel involved in volume regulation in human epithelial cells. *Am J Physiol Cell Physiol*. 2007;292(1):C460.
- V C, S F, T G. Assessment of TRPM7 functions by drug-like small molecules. *Cell Calcium*. 2017;67:166–173. <https://doi.org/10.1016/j.ccca.2017.03.004>.
- Clark K, Langeslag M, Leeuwen BV, Ran L, Ryazanov AG, Figdor CG, Moolenaar WH, Jalink K, Leeuwen FNV. TRPM7, a novel regulator of actomyosin contractility and cell adhesion. *EMBO J*. 2006;25(2):290–301.
- Desai BN, Krapivinsky G, Navarro B, Krapivinsky L, Carter BC, Febyay S, Delling M, Penumaka A, Ramsey IS, Manasian Y. Cleavage of TRPM7 releases the kinase domain from the ion channel and regulates its participation in Fas-induced apoptosis. *Dev Cell*. 2012;22(6):1149–62.
- Smith MA, Schnellmann RG. Calpains, mitochondria, and apoptosis. *Cardiovasc Res*. 2012;96(1):32.
- Cabon L, Galánmalo P, Bouharrou A, Delavallée L, Brunellenavas MN, Lorenzo HK, Gross A, Susin SA. BID regulates AIF-mediated caspase-independent necroptosis by promoting BAX activation. *Cell Death Differ*. 2012;19(2):245–56.
- Gafni J, Cong X, Chen SF, Gibson BW, Ellerby LM. Calpain-1 cleaves and activates Caspase-7. *J Biol Chem*. 2009;284(37):25441–9.
- Kim MJ, Jo DG, Hong GS, Kim BJ, Lai M, Cho DH, Kim KW, Bandyopadhyay A, Hong YM, Kim DH, et al. Calpain-dependent cleavage of cain/cabin1 activates calcineurin to mediate calcium-triggered cell death. *Proc Natl Acad Sci U S A*. 2002;99(15):9870–5.
- Liu L, Xing D, Chen WR, Chen T, Pei Y, Gao X. Calpain-mediated pathway dominates cisplatin-induced apoptosis in human lung adenocarcinoma cells as determined by real-time single cell analysis. *Int J Cancer*. 2008;122(10):2210–22.
- Waterhouse NJ, Finucane DM, Green DR, Elce JS, Kumar S, Alnemri ES, Litwack G, Khanna K, Lavin MF, Watters DJ. Calpain activation is upstream of caspases in radiation-induced apoptosis. *Cell Death Differ*. 1998;5(12):1051–61.
- Ilmarinen P, Moilanen E, Kankaanranta H. Mitochondria in the Center of Human Eosinophil Apoptosis and Survival. *Int J Mol Sci*. 2014;15(3):3952–69.
- Wang Y, Kuramitsu Y, Baron B, Kitagawa T, Akada J, Tokuda K, Cui D, Nakamura K. PERK/CHOP contributes to the CGK733-induced vesicular calcium sequestration which is accompanied by non-apoptotic cell death. *Oncotarget*. 2015;6(28):25252–65.
- Zheng W, Feng X, Qiu L, Pan Z, Wang R, Lin S, Hou D, Jin L, Li Y. Identification of the antibiotic ionomycin as an unexpected peroxisome proliferator-activated receptor γ (PPAR γ) ligand with a unique binding mode and effective glucose-lowering activity in a mouse model of diabetes. *Diabetologia*. 2013;56(2):401–11.

Submit your next manuscript to BioMed Central
and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

