

8-1-2021

## Digital Literacy at an Urban Cancer Center: Implications for Technology Use and Vulnerable Patients

Amy Leader

Thomas Jefferson University, amy.leader@jefferson.edu

Lisa M. Capparella

Thomas Jefferson University, lisa.capparella@jefferson.edu

L. Waldman, BA

Thomas Jefferson University, lauren.waldman@jefferson.edu


Rebecca Cammy

Thomas Jefferson University, rebecca.cammy@jefferson.edu

Alison Petok

Follow this and additional works at: <https://jdc.jefferson.edu/kimmelccfp>

Thomas Jefferson University, alison.petok@jefferson.edu

 Part of the [Oncology Commons](#), and the [Public Health Commons](#)

**[Let us know how access to this document benefits you](#)**

*See next page for additional authors*

---

### Recommended Citation

Leader, Amy; Capparella, Lisa M.; Waldman, BA, L.; Cammy, Rebecca; Petok, Alison; Dean, Rebecca; Shimada, Ayako; Yocavitch, Liana; Rising, Kristin L.; Garber, Gregory; Worster, Brooke; and Dicker, Adam, "Digital Literacy at an Urban Cancer Center: Implications for Technology Use and Vulnerable Patients" (2021). *Kimmel Cancer Center Faculty Papers*. Paper 79.

<https://jdc.jefferson.edu/kimmelccfp/79>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Kimmel Cancer Center Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

---

**Authors**

Amy Leader; Lisa M. Capparella; L. Waldman, BA; Rebecca Cammy; Alison Petok; Rebecca Dean; Ayako Shimada; Liana Yocavitch; Kristin L. Rising; Gregory Garber; Brooke Worster; and Adam Dicker

# Digital Literacy at an Urban Cancer Center: Implications for Technology Use and Vulnerable Patients

Amy E. Leader, DrPH, MPH<sup>1</sup>; Lisa M. Capparella, MSS<sup>1</sup>; Lauren B. Waldman, BA<sup>1</sup>; Rebecca B. Cammy, MSW<sup>1</sup>; Alison R. Petok, MSW, MPH<sup>1</sup>; Rebecca Dean, MSW<sup>2</sup>; Ayako Shimada, MS<sup>1</sup>; Liana Yocavitch, MPH<sup>1</sup>; Kristin L. Rising, MD, MSHP<sup>1</sup>; Gregory D. Garber, MSW<sup>1</sup>; Brooke Worster, MD<sup>1</sup>; and Adam P. Dicker, MD, PhD<sup>1</sup>



**PURPOSE** eHealth literacy, or the ability to seek, find, understand, and appraise health information from electronic sources, has become increasingly relevant in the era of COVID-19, when so many aspects of patient care became dependent on technology. We aimed to understand eHealth literacy among a diverse sample of patients with cancer and discuss ways for health systems and cancer centers to ensure that all patients have access to high-quality care.

**METHODS** A cross-sectional survey of patients with cancer and caregivers was conducted at an NCI-designated cancer center to assess access to the Internet, smartphone ownership, use of mobile apps, willingness to engage remotely with the health care team, and use of the patient portal. Descriptive statistics and bivariate analyses were used to assess frequencies and significant differences between variables.

**RESULTS** Of 363 participants, 55% (n = 201) were female, 71% (n = 241) identified as non-Hispanic White, and 29% (n = 85) reported that their highest level of education was a high school diploma. Most (90%, n = 323) reported having access to the Internet and most (82%, n = 283) reported owning a smartphone. Younger patients or those with a college degree were significantly more likely to own a smartphone, access health information online, know how to download an app on their own, have an interest in communicating with their health care team remotely, or have an account on the electronic patient portal.

**CONCLUSION** As cancer centers increasingly engage patients through electronic and mobile applications, patients with low or limited digital literacy may be excluded, exacerbating current cancer health disparities. Patient-, provider- and system-level technology barriers must be understood and mitigated.

JCO Clin Cancer Inform 5:872-880. © 2021 by American Society of Clinical Oncology

Licensed under the Creative Commons Attribution 4.0 License

## BACKGROUND

The COVID-19 pandemic is an unprecedented and historic event that caused a fundamental shift in how health care is delivered to and received by patients. The rapid uptake of telemedicine and patient portals,<sup>1,2</sup> both for continued health care and access to COVID testing and vaccines, is convenient and efficient for providers but requires resources and digital health literacy from patients. Research is just beginning to understand how reliance on these technologies is affecting vulnerable and under-resourced patients.<sup>3,4</sup> Here, we describe an assessment at our cancer center of our patients' and caregivers' ability to access and use common technology platforms and describe the implications of our findings in the context of health care in a postpandemic world.

Universal health literacy has been identified as a public health goal for the 21st century.<sup>5</sup> In today's digital society, one of the most relevant aspects of health literacy is electronic health literacy, or eHealth

literacy. eHealth literacy has been defined as the ability to seek, find, understand, and appraise health information from electronic sources and apply the knowledge gained to addressing or solving a health problem.<sup>6</sup> Although sufficient health literacy has been associated with positive health outcomes, the impact of eHealth literacy on health outcomes has been less explored,<sup>7</sup> but is assumed to be equally important. The pathway from eHealth literacy to health outcomes may be through patient engagement,<sup>8</sup> such that patients with higher eHealth literacy can more effectively self-assess symptoms, communicate with providers, digest electronic information about their health, and manage biometric measures and medications.

A cancer diagnosis is often accompanied by an onslaught of new information and new health care teams. A recent study found that more than 90% of patients with cancer turn to the Internet to access information about cancer before speaking to a health care professional.<sup>9</sup> Research shows that although the interest in using technology to manage cancer care is

## ASSOCIATED CONTENT

### Data Supplement

Author affiliations and support information (if applicable) appear at the end of this article.

Accepted on July 12, 2021 and published at [ascopubs.org/journal/cci](https://ascopubs.org/journal/cci) on August 24, 2021; DOI <https://doi.org/10.1200/CCI.21.00039>

## CONTEXT

### Key Objective

The COVID-19 pandemic has caused increased reliance on technology to access cancer care. It is important to understand the distribution of eHealth literacy, or the ability to seek, find, or understand health information from electronic sources, in a cancer patient population to know where disparities may exist. We surveyed 363 patients with cancer and caregivers to document their eHealth literacy, as well as predictors of eHealth literacy.

### Knowledge Generated

We found that older patients, those with a lower educational level, and those from a minority race or ethnicity had the lowest levels of eHealth literacy. They were the least likely to have access to or use technology in their everyday life and use it in managing their cancer care.

### Relevance

Cancer centers must be cognizant of these disparities when using technology with patients, to not widen cancer health disparities.

high, the actual adoption of such technology among patients with cancer is much lower and disparities in use exist.<sup>10,11</sup> Disparities in eHealth literacy and use are most frequently seen among older patients, those who have lower socioeconomic status, and those who identify with a racial or ethnic minority group.<sup>12</sup> Finally, patients with lower health literacy report lower levels of shared decision making<sup>13,14</sup>; by extension, it can be assumed that patients with lower eHealth literacy would also report lower levels of shared decision making, although this has yet to be explored.

The purpose of this study was to assess differences in patient access to and use of digital technology and electronic health information in their daily lives and in relation to their cancer care. The study was stimulated by an interest in using digital health to help our patients. As we began to consider remote monitoring for patient-reported outcomes, fundamental questions regarding digital literacy were raised. We surveyed a diverse sample of patients and caregivers who received care from a large, urban NCI-designated cancer center. We expected to see the greatest disparities in eHealth literacy among our older patients with cancer, those who have fewer years of education, and those who identify with a minority racial or ethnic group. The results of the study can inform best practices for ensuring that all patients with cancer have access to information and their health care providers, as well as informing future interventions to reduce the digital literacy divide among medically underserved patients.

## METHODS

### Participants

As patients checked in for scheduled appointments in either Medical Oncology or Radiation Oncology, they were given a paper-based survey to complete either on their own or with assistance if they had difficulties or questions. A

smaller sample of patients completed the survey on an electronic tablet in our Center Support and Welcome Center and were offered assistance or a paper survey if they preferred. All patients with cancer and caregivers were eligible to participate in the survey if they were older than age 18 years and able to read and speak English. All participants provided verbal consent before beginning the survey. Survey data were collected anonymously. Participants were not compensated for their time. We received expedited approval from our institution's institutional review board to conduct the survey.

### Survey

The 24-item survey (Data Supplement) was created by the research team to meet the needs of the cancer center and was divided into four content areas: access and use of the Internet for health information; smartphone ownership; use of mobile health apps and wearable technology; and use of technology to interact with health care professionals. For the first content area, we asked questions about where patients and caregivers typically access the Internet, the type of device they typically use to access the Internet, whether they have ever visited a web site to learn about cancer, and whether they have ever visited our cancer center's web site to learn about resources available to them. For the second content area, we asked participants whether they have a cell phone or smartphone, the brand of the phone, how much time they typically spend on their phone, and what they do on their phone. For the third content area, we asked participants how many apps are currently on their phone or tablet, if any of the apps are health-related, if they can find and download an app on their own, if they have ever used an app to help with a decision related to their cancer care, and if they routinely use any wearable technology. For the fourth content area, we asked whether participants would be comfortable communicating with their provider over their phone in a

telemedicine visit and whether they use our cancer center's electronic portal to manage their cancer care. Finally, we collected demographic information from each participant to assess whether they were a patient or a caregiver, their age, sex, race or ethnicity, zip code, and educational achievement. The survey took about 10 minutes to complete.

### Data Analysis

Descriptive statistics were computed for all variables. Some demographic variables were collapsed based on sample distribution and recoded for bivariate analyses. Age was categorized as below age 60, between age 60-69, and older than age 70 years. Race and ethnicity were categorized as non-Hispanic White or minority race or ethnicity (included non-Hispanic Black, Hispanic, and Asian participants). Differences in technology use by sex, ages, race or ethnicities, and education levels were tested by chi-square tests. Statistical significance was defined as  $P < .05$  on a two-tailed distribution. To evaluate the associations between (1) find and download an app unassisted, (2) willing to do telehealth, and (3) use of electronic patient portal and participants' characteristics (age, education level, and race), three separate logistic regression model analyses were conducted. For characteristics with more than two categories, type-III  $P$  values representing the overall association of the characteristic with the digital literacy measure were calculated as were pairwise comparisons with a reference category. The significance level of all tests was set a priori to the .05 level. All analyses were performed with SAS 9.4. Missing data, which were determined to be minimal and at random, were excluded from analyses on a question-by-question basis.

## RESULTS

### Description of Participants

Four hundred and fifty ( $n = 450$ ) patients and caregivers were approached to complete the survey and 363 participants completed the survey. Three hundred and forty-seven ( $n = 347$ , 96%) were completed on paper and 16 (4%) were completed on tablet computer. Three hundred forty-six ( $n = 346$ , 96%) were patients, 13 (4%) were caregivers, and four were unknown. Almost half of the participants ( $n = 151$ , 42%) were younger than age 60 years, whereas an additional one quarter ( $n = 84$ , 23%) were older than age 70 years. Fifty-five percent of participants ( $n = 201$ ) were female, whereas 45% of participants ( $n = 160$ ) were male. The majority of participants were non-Hispanic White ( $n = 241$ , 66%), whereas 23% of participants ( $n = 85$ ) were non-Hispanic Black. Educational attainment was divided among participants: 27% ( $n = 99$ ) had a high school diploma, 17% ( $n = 60$ ) had a trade or an associate degree, 23% ( $n = 85$ ) obtained a bachelor's degree, and 15% ( $n = 53$ ) had an advanced degree. Characteristics of participants are displayed in [Table 1](#).

### Access and Use of the Internet for Health Information

Thirty-seven participants (10%) reported that they do not use the Internet at all. Among those who do, patients and caregivers predominantly accessed the Internet at home ( $n = 309$ , 86%), followed by at work ( $n = 77$ , 20%) and at a public place ( $n = 22$ , 6%). Participants reported that they accessed the Internet in multiple ways, using computers ( $n = 248$ , 70%), cell phones ( $n = 191$ , 54%), and tablets ( $n = 125$ , 35%). Most participants ( $n = 252$ , 69%) reported that they had visited a web site to learn more about their cancer. These data are presented in [Table 2](#).

### Smartphone Ownership

Nearly all participants had a cell phone (98%,  $n = 354$ ); for 283 (80%) of those participants, their phone is a smartphone. Among the 283 participants who own a smartphone, 55% ( $n = 160$ ) reported spending less than 2 hours per day on their phone. Thirty percent reported using their phones 2-4 hours per day ( $n = 86$ ) and 15% ( $n = 44$ ) reported using their phones for more than 4 hours per day. Participants reported using their smartphones primarily for communication ( $n = 269$ , 95%), accessing the Internet ( $n = 201$ , 71%), and navigating to places ( $n = 170$ , 60%).

**TABLE 1.** Participant Characteristics ( $n = 363$ )

Characteristic	Participants, No. (%)
Role	
Patient	346 (96)
Caregiver	13 (4)
Missing	4 (1)
Age, years	
< 60	151 (42)
60-69	127 (35)
> 70	84 (23)
Missing	1 (< 1)
Sex	
Female	201 (55)
Male	160 (45)
Missing	2 (< 1)
Ethnicity	
Non-Hispanic White	241 (66)
Non-Hispanic Black	85 (23)
Hispanic	7 (2)
Asian	9 (2)
Missing	21 (6)
Education level	
High school or equivalent	99 (27)
Some college or associate degree	60 (17)
Bachelor's degree	85 (23)
Greater than bachelor's degree	53 (15)
Missing	66 (18)

**TABLE 2.** eHealth Literacy Among Participants (n = 363)

Characteristic	Participants, No. (%)
Own a cell phone	
Yes	354 (98)
No	5 (1)
Missing	4 (1)
Own a smartphone	
Yes	283 (78)
No	64 (9)
Missing	16 (4)
Routinely access the Internet	
Yes	325 (90)
No	37 (10)
Missing	1 (< 1)
Where access the Internet (check all that apply)	n = 445 responses
At home	309 (86)
At work	77 (20)
In public	22 (6)
No access	37 (10)
Devices used to access the Internet (check all that apply)	n = 564 responses
Computers	248 (70)
Cell phones	191 (54)
Tablets	125 (35)
Visited a web site for cancer information	
Yes	252 (69)
No	106 (29)
Missing	5 (2)
Find and download a smartphone app unassisted	
Yes	249 (69)
No	107 (29)
Missing	7 (2)
Willing to do visit by telehealth	
Yes	244 (67)
No	110 (30)
Missing	9 (3)
Use of electronic patient portal	
Yes, routinely use	162 (45)
Yes, registered but do not use	49 (14)
No, not registered	111 (31)
Missing	41 (10)
Use wearable technology	
Yes	35 (10)
No	260 (72)
Missing	68 (19)

### Use of Mobile Health Apps and Wearable Technology

Only about 70% of participants (n = 249) reported that they can find and download them on their own. Most (62%, n = 225) did not have a health-related app on their phone or tablet. Only 10% (n = 35) reported using wearable technology, including brands like Fitbit and Apple Watch. These data are presented in [Table 2](#).

### Using Technology to Interact with Health Care Professionals

Sixty-seven percent of participants (n = 244) said that they would feel comfortable communicating with their doctor or nurse using a smartphone or tablet (ie, a telehealth visit). Just under half of the participants (45%, n = 162) said they use the cancer center's patient portal fairly often, 14% (n = 49) said they signed up for access to the patient portal but do not use it often, and 31% (n = 111) said they do not use the patient portal. These data are presented in [Table 2](#).

### Differences in Technology Use by Age, Educational Attainment, and Race or Ethnicity

There were significant differences in the use of technology by age. The most notable were that older participants were less likely to access the Internet, own a smartphone, download an app by themselves, or have an interest in communicating electronically with providers. Almost all participants, across all educational levels, reported owning a cell phone (98.6%, n = 355). However, there were significant differences in the use of technology by educational attainment. The largest disparities were that those with only a high school education were less likely to access the Internet, visit a web site for health purposes, be able to download an app by themselves, have an interest in communicating electronically with their providers, or have any wearable technology. Finally, there were significant differences by race or ethnicity of the respondent. Participants from minority races and ethnicities were significantly less likely to routinely access the Internet, visit a web site to look for cancer information, be able to find and download an app on their own, and use the patient portal. There were no reported differences in technology use by sex. All these data can be found in [Table 3](#).

### Prediction Models for Important eHealth Skills

We aimed to understand which factors would be predictive for three key eHealth literacy skills: finding and downloading an app unassisted, willingness to do a telehealth appointment, and using a patient portal. Age and educational attainment were significantly predictive across all three models, with those who were younger or had more education being more likely to use digital technology. In

**TABLE 3.** eHealth Literacy by Participant Age, Educational Attainment, and Race

Participant Characteristic	Age (years)				Educational Attainment					Race		
	< 60	60-69	> 70	$\chi^2$ P	High School Graduate or Less	Trade School or Some College	Bachelor's Degree	Advanced Degree	$\chi^2$ P	White	Other Races Combined	$\chi^2$ P
Own a cell phone, No. (% yes)	149 (99.3)	125 (99.2)	80 (96.4)	3.89 (.143)	99 (100.0)	60 (100.0)	84 (98.9)	51 (96.2)	5.69 (.127)	237 (99.2)	99 (98.0)	0.798 (> .05)
Own a smartphone, No. (% yes)	135 (90.6)	97 (80.2)	50 (65.8)	20.77 (< .01)	68 (73.1)	46 (79.3)	78 (95.1)	46 (88.5)	27.05 (< .01)	189 (90.6)	79 (80.2)	9.14 (< .05)
Routinely access the Internet, No. (% yes)	146 (96.7)	112 (88.2)	67 (79.8)	17.39 (< .01)	76 (76.8)	55 (91.7)	84 (98.8)	52 (98.1)	30.39 (< .01)	224 (92.9)	85 (84.2)	6.30 (< .05)
Visited a web site for cancer info, No. (% yes)	111 (73.5)	91 (72.8)	50 (61)	4.54 (.10)	48 (49.5)	45 (75.0)	75 (88.2)	47 (86.8)	41.99 (< .01)	181 (75.1)	56 (57.7)	9.96 (< .05)
Find and download an app unassisted, No. (% yes)	126 (83.4)	84 (66.7)	39 (49.4)	33.70 (< .01)	53 (54.1)	44 (73.3)	74 (88.1)	42 (80.8)	30.20 (< .01)	176 (74.3)	63 (63.0)	8.74 (< .05)
Willing to do visit by telehealth, No. (% yes)	122 (81.9)	85 (68.5)	37 (45.7)	32.12 (< .01)	51 (53.7)	42 (70.0)	70 (83.3)	41 (77.4)	20.46 (< .01)	169 (71.3)	65 (65.7)	1.10 (> .05)
Use of electronic patient portal, No. (% yes)	71 (52.6)	62 (54.9)	29 (39.2)	4.88 (.087)	30 (35.3)	28 (50.0)	46 (61.3)	31 (64.6)	24.16 (< .01)	127 (59.3)	28 (30.4)	24.81 (< .01)
Use wearable technology, No. (% yes)	24 (19.7)	8 (7.5)	3 (4.5)	12.50 (< .01)	2 (2.4)	5 (9.6)	16 (22.2)	6 (13.3)	15.02 (< .01)	25 (13.0)	7 (8.1)	1.36 (> .05)

**TABLE 4.** Logistic Regression Model Analyses for Select eHealth Literacy Skills

Effect	OR (95% CI)	P
Estimated OR of being able to find and download an app unassisted		
Age, years		
< 60	Ref	< .001
60-69	0.48 (0.24 to 0.95)	.035
> 70	0.18 (0.08 to 0.38)	< .001
Educational attainment		
HS graduate or less	Ref	< .001
Some college or tech school	2.17 (1.02 to 4.61)	.044
Bachelor's degree	5.98 (2.65 to 13.46)	< .001
Advanced degree	3.67 (1.54 to 8.73)	.003
Race		
White	Ref	
Other races combined	0.55 (0.29 to 1.06)	.074
Estimated OR of willing to do telehealth		
Age, years		
< 60	Ref	< .001
60-69	0.64 (0.33 to 1.22)	.174
> 70	0.19 (0.09 to 0.40)	< .001
Educational attainment		
HS graduate or less	Ref	< .001
Some college or tech school	1.96 (0.94 to 4.09)	.074
Bachelor's degree	4.44 (2.11 to 9.35)	< .001
Advanced degree	3.50 (1.50 to 8.13)	.004
Race		
White	Ref	
Other races combined	0.68 (0.36 to 1.28)	.235
Estimated OR of using electronic patient portal		
Age, years		
< 60	Ref	.013
60-69	1.08 (0.59 to 1.98)	.798
> 70	0.38 (0.18 to 0.79)	.009
Educational attainment		
HS grad or less	Ref	.006
Some college or tech school	1.26 (0.60 to 2.65)	.544
Bachelor's degree	2.60 (1.31 to 5.15)	.006
Advanced degree	3.21 (1.44 to 7.14)	.004
Race		
White	Ref	
Other races combined	0.26 (0.13 to 0.48)	< .001

Abbreviations: OR, odds ratio; ref, reference.

one model, using a patient portal, those who were not White were significantly less likely to use a patient portal for their health care needs. The results of the three models are found in [Table 4](#).

## DISCUSSION

Our study aimed to document eHealth literacy among a diverse sample of patients with cancer and caregivers at an urban, academically based cancer center. We found that



although access to the Internet and smartphone use was relatively high among our patients and caregivers, disparities in access and use of the technology exist among our most vulnerable patients with cancer, most notably those who are older and of lower educational attainment. Although this has been documented previously,<sup>15,16</sup> it remains worrisome as it is evidence that we are expanding, rather than closing, the digital divide among patients and caregivers who need access to information and technology to manage their disease and interact with their care team.

Given these findings, the question of how to provide the same level of access to health information and health care to all patients, regardless of eHealth literacy level, is an important one. Patient navigators, which are deployed by many health systems to provide an extra level of patient support, may be able to overcome some of the technical barriers of cancer care.<sup>17</sup> Whether it is teaching them how to download and use a health app to track their cancer care or create an account on the patient portal, patient navigators or support personnel may be one answer to ensuring equal access for all. Our cancer center is currently offering classes to teach patients and caregivers how to sign up and access their electronic health records. We created a telehealth task force to walk patients through the process of conducting a telehealth visit. Other suggestions include linking patients to community resources that provide free or low-cost home or publicly available Internet access.<sup>18</sup> Recently, many of these recommendations were included in an 18-point Digital Universal Precautions for health care organizations that want to make digital health accessible and meaningful for all patients, regardless of digital health literacy.<sup>3</sup>

Because of the COVID-19 pandemic, providers and health systems were forced to reimagine health care, much of it transitioning to virtual platforms. In our study, we found that patients who are older and have lower educational attainment are less willing to participate in telehealth visits; patients of racial and ethnic minority heritages are less likely to use our patient portal to manage their care. Knowing these patients with cancer are already at increased risk for poor cancer outcomes and now are at increased risk of poor outcomes from the pandemic,<sup>19</sup> leaving these patients behind with our continued use of technology could have a disastrous impact. To ensure that the expansion of telemedicine does not exacerbate health disparities, four key actions were recently proposed: (1) proactively exploring disparities in telemedicine access; (2) developing solutions to mitigate barriers; (3) removing health system–created barriers to access; and (4) advocating for policies and infrastructure that facilitate equitable access.<sup>20</sup> This encompasses the need for health systems to

create and use technology that patients want to adopt to manage their health care needs.<sup>21</sup>

The research described here can be used as a starting point toward developing a tool to identify those with low eHealth literacy, to be able to deploy resources and support to those who need assistance in navigating health care in a digital world. Although a few digital literacy screening tools exist or are in development,<sup>22-25</sup> none have the ability to screen for skills that we feel are essential in this post-pandemic era such as downloading an app, conducting a telehealth visit, or logging into a patient portal. While we acknowledge that screening tools should not replace efforts to provide universal access to literacy-appropriate health care,<sup>26</sup> identifying and assisting those at greatest risk of not being able to access care remains an important aspect of supportive cancer care.

There are limitations to our study. While comprising a large sample of patients with cancer and caregivers, it was a convenience sample of patients and the results may not be generalizable to the overall cancer patient population. We limited our sample to English-speaking patients and were not able to capture eHealth and digital literacy disparities that are known to exist in the Hispanic and Latino population.<sup>27</sup> These data are self-reported and do not explore why some participants are not using certain technologies. The survey that participants completed was created by the research team to provide data on questions and concerns that were most pressing to the cancer center and deviated from using known scales of eHealth literacy that, although validated,<sup>22-25</sup> were not as relevant to our needs.

Going forward, as health systems and cancer centers unveil new programs and options for patients, they must be cognizant that not all patients have access to these technologies nor the capabilities or knowledge to use them to their full capacity. Unfortunately, there are few evidence-based approaches to increasing eHealth literacy in socially disadvantaged groups.<sup>28</sup> As more appointments are scheduled and occur online and through telehealth, as more cancer centers rely on patient-generated health data to track symptoms and evaluate quality of care,<sup>29</sup> and as more information is posted exclusively on web sites and through social media platforms, there is a segment of the patient population that will not be privy to these services. Leadership must remain cognizant of disparities in eHealth literacy and ensure that until every patient can fully participate, a low-tech option must be available. Directing all patient traffic to a web site, a portal, or an app may be easier and more cost effective for the health system but it does so at a cost to vulnerable patients.

## AFFILIATIONS

<sup>1</sup>Sidney Kimmel Cancer Center, Thomas Jefferson University, Philadelphia, PA

<sup>2</sup>School of Social Policy and Practice, University of Pennsylvania, Philadelphia, PA

## CORRESPONDING AUTHOR

Amy E. Leader, DrPH, MPH, Division of Population Science, Department of Medical Oncology, Sidney Kimmel Cancer Center, Thomas Jefferson University, 834 Chestnut St, Suite 314, Philadelphia, PA 19107; e-mail: amy.leader@jefferson.edu.

## SUPPORT

Supported by NIH/NCI 5P30CA056036-17.

## AUTHOR CONTRIBUTIONS

**Conception and design:** Amy E. Leader, Rebecca B. Cammy, Alison R. Petok, Gregory D. Garber, Brooke Worster, Adam P. Dicker

**Administrative support:** Lisa M. Capparella

**Provision of study materials or patients:** Lisa M. Capparella

**Collection and assembly of data:** Amy E. Leader, Lisa M. Capparella, Lauren B. Waldman, Rebecca B. Cammy, Rebecca Dean, Gregory D. Garber, Brooke Worster, Adam P. Dicker

**Data analysis and interpretation:** Amy E. Leader, Lauren B. Waldman, Ayako Shimada, Liana Yocavitch, Kristin L. Rising, Gregory D. Garber, Brooke Worster, Adam P. Dicker

**Manuscript writing:** All authors

**Final approval of manuscript:** All authors

**Accountable for all aspects of the work:** All authors

## AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The following represents disclosure information provided by the authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate

Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to [www.asco.org/rwc](http://www.asco.org/rwc) or [ascopubs.org/cci/author-center](http://ascopubs.org/cci/author-center).

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](http://OpenPayments)).

**Kristin L. Rising**

**Research Funding:** Ortho Clinical Diagnostics

**Gregory D. Garber**

**Consulting or Advisory Role:** Ethos Cannabis, Bristol Myers Squibb Foundation

**Brooke Worster**

**Consulting or Advisory Role:** Ethos Cannabis

**Research Funding:** Ethos Cannabis

**Adam P. Dicker**

**Stock and Other Ownership Interests:** Oncohost, Self Care Catalysts

**Consulting or Advisory Role:** EMD Serono, Janssen, Roche, Cybexa Therapeutics, Oncohost, Third Bridge, Accordant, Alcimed, Orano Med, IBA

**Patents, Royalties, Other Intellectual Property:** We recently filed a patent Doped BEO Compounds for Optically Stimulated Luminescence (OSL) and Thermoluminescence (TL) Radiation Dosimetry

**Expert Testimony:** Wilson Sonsini

**Travel, Accommodations, Expenses:** Oncohost

**Other Relationship:** European Commission

**Uncompensated Relationships:** Google, Dreamit Ventures

No other potential conflicts of interest were reported.

## ACKNOWLEDGMENT

The authors thank all the patients and caregivers at our cancer center who took the time to complete the survey.

## REFERENCES

1. Wosik J, Fudim M, Cameron B, et al: Telehealth transformation: COVID-19 and the rise of virtual care. *J Am Med Inform Assoc* 27:957-962, 2020
2. Bokolo Anthony Jnr: Use of telemedicine and virtual care for remote treatment in response to COVID-19 pandemic. *J Med Syst* 44:132, 2020
3. Smith B, Magnani JW: New technologies, new disparities: The intersection of electronic health and digital health literacy. *Int J Cardiol* 292:280-282, 2019
4. Abdel-Rahman O: Patient-related barriers to some virtual healthcare services among cancer patients in the USA: A population-based study. *J Comp Eff Res* 10:119-126, 2021
5. Nutbeam D: Health literacy as a public health goal: A challenge for contemporary health education and communication strategies into the 21<sup>st</sup> century. *Health Promot Int* 15:259-267, 2000
6. Norman CD, Skinner HA: eHealth literacy: essential skills for consumer health in a networked world. *J Med Internet Res* 8:e9, 2006
7. Neter E, Brainin E: Association between health literacy, eHealth literacy, and health outcomes among patients with long-term conditions: A systematic review. *Eur Psychol* 24:68-81, 2019
8. Barelo S, Triberti S, Graffigna G, et al: eHealth for patient engagement: a systematic review. *Front Psychol* 8:2013, 2016
9. Hoogland AI, Mansfield J, Lafranchise EA, et al: eHealth literacy in older adults with cancer. *J Geriatr Oncol* 11:1020-1022, 2020
10. Jiang Y, West BT, Barton DL, et al: Acceptance and use of eHealth/mHealth applications for self-management among cancer survivors. *Stud Health Technol Inform* 245:131-135, 2017
11. Poddar R, Thomas A, MiMeglio M, et al: Access to Internet, smartphone usage, and acceptability of mobile health technology among cancer patients. *Support Care Cancer* 28:5455-5461, 2020
12. Neter E, Brainin E: e-Health literacy, extending the digital divide to the realm of health information. *J Med Internet Res* 14:e19, 2012
13. Brabers AE, Rademakers JJ, Groenewegen PP, et al: What role does health literacy play in patients' involvement in medical decision-making? *PLoS One* 12:e0173316, 2017
14. Chang HL, Li FS, Lin CF: Factors influencing implementation of shared medical decision making in patients with cancer. *Patient Prefer Adherence* 13:1995-2005, 2019
15. Chesser A, Burke A, Reyes J, et al: Navigating the digital divide: A systematic review of eHealth literacy in underserved populations in the United States. *Inform Health Soc Care* 41:1-19, 2016
16. Tarver WL, Haggstrom DA: The use of cancer-specific patient-centered technologies among underserved populations in the United States: Systematic review. *J Med Internet Res* 21:e10256, 2019

17. National Academies of Science: Establishing Effective Patient Navigation Programs in Oncology: Proceedings from a Workshop. Washington DC, National Academies Press, 2018
18. Salovey P, Williams-Piehota P, Mowad L, et al: Bridging the digital divide by increasing computer and cancer literacy: Community technology centers for head-start parents and families. *J Health Commun* 14:228-245, 2009
19. APM Research Lab Staff: The color of coronavirus.: COVID-19 deaths by race and ethnicity in the U.S. APM Research Lab. American Public Media. <https://www.apmresearchlab.org/covid/deaths-by-race>
20. Nouri S, Khoong EC, Lyles CR, et al: Addressing equity in telemedicine for chronic disease management during the COVID-19 pandemic. *NEJM Catal* 4, 2020. [10.1056/CAT.20.0123](https://doi.org/10.1056/CAT.20.0123)
21. Rodriguez JA, Clark CR, Bates DW: Digital health equity as a necessity in the 21st century cures act era. *JAMA* 323:2381-2382, 2020
22. Norman CD, Skinner HA: eHEALS: the eHealth Literacy Scale. *J Med Internet Res* 8:e27, 2006
23. van der Vaart R, Drossaert C: Development of the digital health literacy instrument: Measuring a broad spectrum of health 1.0 and health 2.0 skills. *J Med Internet Res* 19:e27, 2017
24. Seçkin G, Yeatts D, Hughes S, et al: Being an informed consumer of health information and assessment of electronic health literacy in a national sample of Internet users: Validity and reliability of the e-HLS Instrument. *J Med Internet Res* 18:e161, 2016
25. Karnoe A, Furstrand D, Christensen KB, et al: Assessing competencies needed to engage with digital health services: Development of the eHealth Literacy Assessment Toolkit. *J Med Internet Res* 20:e178, 2018
26. Kronzer VL: Screening for health literacy is not the answer. *BMJ* 354:i3699, 2016
27. Millar RJ, Sahoo S, Yamashita T, et al: Literacy skills, language use, and online health information seeking among Hispanic adults in the United States. *Patient Educ Couns* 103:1595-1600, 2020
28. Jacobs RJ, Lou JQ, Ownby RL, et al: A systematic review of eHealth interventions to improve health literacy. *Health Inform J* 22:81-98, 2016
29. Jim HS, Hoogland AI, Brownstein NC, et al: Innovations in research and clinical care using patient-generated health data. *CA Cancer J Clin* 70:182-199, 2020

