

1-1-2023

Testing for Homologous Recombination Repair or Homologous Recombination Deficiency for Poly (ADP-ribose) Polymerase Inhibitors: A Current Perspective

Thomas J Herzog

Ignace Vergote

Leonard G Gomella

Tsveta Milenkova

Tim French

See next page for additional authors

Follow this and additional works at: <https://jdc.jefferson.edu/urologyfp>



Part of the [Urology Commons](#)

[Let us know how access to this document benefits you](#)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Urology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Thomas J Herzog, Ignace Vergote, Leonard G Gomella, Tsveta Milenkova, Tim French, Raffi Tonikian, Christian Poehlein, and Maha Hussain



Current Perspective

Testing for homologous recombination repair or homologous recombination deficiency for poly (ADP-ribose) polymerase inhibitors: A current perspective



Thomas J. Herzog^{a,*}, Ignace Vergote^b, Leonard G. Gomella^c,
Tsveta Milenkova^d, Tim French^d, Raffi Tonikian^{e,1}, Christian Poehlein^e,
Maha Hussain^f

^a University of Cincinnati Cancer Center, University of Cincinnati Medical Center, 234 Goodman St, Cincinnati, OH 45219, USA

^b Department of Gynecology and Obstetrics, Division of Gynecologic Oncology, Leuven Cancer Institute, Catholic University Leuven, Herestraat 49, 3000 Leuven, Belgium, European Union

^c Department of Urology, Sidney Kimmel Cancer Center—Jefferson University Health, 1025 Walnut St Suite 1100, Philadelphia, PA 19107, USA

^d AstraZeneca, 316 Hills Rd, Cambridge CB2 8PA, UK

^e Merck & Co., Inc., 90 E Scott Ave, Rahway, NJ 07065, USA

^f Robert H. Lurie Comprehensive Cancer Center of Northwestern University, 420 E Superior St, Chicago, IL 60611, USA

Received 5 July 2022; received in revised form 20 October 2022; accepted 22 October 2022

Available online 2 November 2022

KEYWORDS

Homologous recombination;
Poly(ADP-ribose) polymerase inhibitors;
Ovarian neoplasms;
Prostate neoplasms;
Breast neoplasms;

Abstract Poly (ADP-ribose) polymerase inhibitors (PARPis) have demonstrated clinical activity in patients with *BRCA1* and/or *BRCA2* mutated breast, ovarian, prostate, and pancreatic cancers. Notably, BRCA mutations are associated with defects in the homologous recombination repair (HRR) pathway. This homologous recombination deficiency (HRD) phenotype can also be observed as genomic instability in tumour cells. Accordingly, PARPi sensitivity has been observed in various tumours with HRD, independent of BRCA mutations. Currently, four PARPis are approved by regulatory agencies for the treatment of cancer across multiple tumour types. Most indications are specific to tumours with a confirmed

* Corresponding author: University of Cincinnati Cancer Center, University of Cincinnati Medical Center, 234 Goodman St, Cincinnati, OH 45219, USA

E-mail address: herzogtj@ucmail.uc.edu (T.J. Herzog).

¹ Current affiliation: Biologiques AbCellera Québec Inc., Montréal, QC, Canada.

Metastatic pancreatic neoplasms

BRCA mutation, mutations in other HRR-related genes, HRD evidenced by genomic instability, or evidence of platinum sensitivity. Regulatory agencies have also approved companion and complementary diagnostics to facilitate patient selection for each PARPi indication. This review aims to summarise the biological basis, clinical validation, and clinical relevance of the available diagnostic methods and assays to assess HRD.

© 2022 Merck Sharp & Dohme LLC., a subsidiary Merck & Co., Inc., Rahway, NJ, USA and The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Genome stability is essential to life, and the cell has evolved multiple DNA damage response (DDR) pathways to detect and repair DNA damage [1,2]. Defects in these pathways can lead to mutations, genomic instability, cell death and tumourigenesis [1,2]. Poly (ADP-ribose) polymerases (PARPs) play an essential role in DDR pathways and represent a target for anticancer therapy [1,3]. In the past decade, PARP inhibitors (PARPis) have emerged as a new class of anticancer drugs in multiple cancers, including ovarian (niraparib, olaparib and rucaparib), metastatic castration-resistant prostate (mCRPC; olaparib and rucaparib), breast (olaparib and talazoparib) and metastatic pancreatic (olaparib) [4–11]. Predicting benefit from PARPis is critical for optimal clinical use, and several biomarker tests are available to determine patient selection for PARPis across treatment settings, with more tests in development [3]. These tests identify tumours with defects in DNA repair pathways, most specifically the homologous recombination repair (HRR) pathway, which make them vulnerable to PARPis. Choice of the appropriate biomarker test depends on tumour type and may even differ by line of therapy for the same tumour type, thereby presenting challenges for the clinician. This review aims to enhance the understanding of different biomarker tests to assess homologous recombination deficiency (HRD) and their

relevance for PARPi treatment selection in different tumour types.

2. HRD in cancer

Numerous genes have been implicated in the HRR pathway, including those encoding sensor proteins involved in DSB detection (i.e. γ H2AX, ATM and ATR), signal mediator proteins (i.e. BRCA1, BRCA2 and PALB2), and an effector protein (RAD51) that promotes strand invasion and replication fork stabilisation (Fig. 1) [12]. The best-characterised HRR genes in relation to cancer are BRCA1 and BRCA2, which have been associated with breast, ovarian, pancreatic and prostate cancers [13].

HRD is characterised by an inability to repair DNA DSBs through the HRR pathway. Causes of HRD are not always known but include loss-of-function mutations and epigenetic modifications in HRR-related genes, particularly BRCA1 and/or BRCA2 [14]. HRD typically results in accumulation of genomic damage, which may lead to mitotic catastrophe and cell death as well as carcinogenesis [15]. Consequently, cancer cells can demonstrate genomic instability [16,17], which may manifest as loss of heterozygosity (LOH; existence of a single allele resulting from a cross-chromosomal event that leads to the loss of entire genes and the surrounding chromosomal region [17]); telomeric allelic imbalance



Fig. 1. Measuring HRD. (A) The identification of mutations in BRCA and other HRR-related genes and detecting genomic instability are two principal ways to detect HRD. (B) Notably, LOH, TAI and LST are all measures of genomic instability. Abbreviations: HRD, homologous recombination deficiency; HRR, homologous recombination repair; LOH, loss of heterozygosity; LST, large-scale state transitions; TAI, telomeric allelic imbalance.

(TAI; a discrepancy in the 1:1 allele ratio in the telomere of the chromosome owing to reciprocal translocations [18]); and large-scale state transitions (LST; chromosomal breaks between adjacent regions of at least 10 Mb [19]).

3. PARPi: indications

In 2014, olaparib became the first PARPi approved by the US Food and Drug Administration (FDA) specifically for treating germline BRCA-mutated metastatic ovarian cancer after ≥ 3 lines of chemotherapy [20]. Currently, four PARPis (olaparib, rucaparib, talazoparib and niraparib) are approved by regulatory agencies for the treatment of multiple tumour types. Most indications are specific to tumours with a confirmed BRCA mutation, other HRR mutation, or HRD genomic instability (Supplementary Table 1), but PARPi therapy is also broadly indicated for patients with platinum-sensitive disease relapse [4–10]. Four companion diagnostics are approved in conjunction with the drug approvals to facilitate patient selection.

4. Biomarker tests for PARPi

PARPi sensitivity has been well established for tumours that harbour BRCA mutations [21]. The prevalence of BRCA mutations have been reported to range from 1% to 15% (*BRCA1*) and 2%–6% (*BRCA2*) in advanced/metastatic breast cancer, 12%–15% (*BRCA1*) and 5%–7% (*BRCA2*) in ovarian cancer, 0.3%–1% (*BRCA1*) and 5%–6% (*BRCA2*) in metastatic prostate cancer and 0.3%–2.3% (*BRCA1*) and 0.7%–5.7% (*BRCA2*) in pancreatic cancer [3,22–24]. However, clinical data have shown that PARPi therapy can be efficacious for patients who test negative with existing HRD diagnostics as well as for patients whose tumours are driven by other HRR mutations or HRD genomic instability that is unrelated to BRCA mutations [25–29]. Accordingly, PARPi biomarker tests that are currently available can be categorised into three approaches depending on how they aim to detect the presence of HRD in tumour cells, which include identifying the presence of a mutation that causes HRD (e.g. BRCA mutations), detecting genomic instability (indicative of HRD) or directly testing HRR function in a cellular assay [3,22,23].

The first approach is evaluating tumour mutation status by gene sequencing of a panel of specified genes known to cause HRD. *BRCA1* and *BRCA2* are the most well-characterised HRR genes in terms of their relationship to cancer and should always be evaluated. This approach can be expanded to include multiple other genes related to the HRR pathway, but, in some diseases, such as first-line ovarian cancer, non-BRCA HRR mutations do not predict PARPi response, while HRD genomic instability has been predictive (Fig. 1)

[3,22,23,30]. Emerging evidence suggests that other HRR gene alterations may be associated with different PARPi sensitivity.

The second approach investigates whether the consequences of HRD, namely genomic instability, are present by identifying chromosomal aberrations (i.e. genomic scars), which are changes in the genome as a consequence of the dysfunctional HRR pathway (both direct and indirect) including LOH, TAI and LST [3,22,23]. Given the recognition of BRCA mutations as an archetypal cause of HRD, BRCA testing is invariably conducted. As such, some assays combine the measurement of both the HRR mutations (i.e. BRCA) and HRD genomic instability approaches.

The third approach directly evaluates HRR function in tumour cells to determine if there is proficiency. An example of a functional assay is the RAD51 foci assay, which measures the capacity of tumour cells to recruit nuclear RAD51 foci during the S/G2 cell cycle phase in the presence of DSBs using immunofluorescence [31]. Functional assays provide a measure of HRR competency, which may be important when, for example, a reversion mutation results in a tumour switching from HRD status to being HRR proficient [32]. Because of technical complexity, functional assays such as RAD51 foci are not routinely used clinically.

Table 1 [33–39] summarises the HRD diagnostics for various PARPi indications. These assays use a variety of sample types (blood or tumour tissue). Other tests for HRR mutations and HRD genomic instability, including locally developed tests, are being established, validated and commercialised.

5. Clinical validity and utility of HRR mutations and tests for HRD in various cancers

In the first-line maintenance treatment setting of ovarian cancer, the benefit of PARPis in BRCA-mutated tumours was demonstrated in randomised phase 3 trials (Table 2 [28,30,40–49]).

HRD positivity (defined as a tumour with a BRCA mutation or a genomic instability score above a specified threshold for the HRD assay used) was also associated with a PFS benefit with PARPis in randomised phase 3 trials in the first-line maintenance treatment setting in advanced ovarian cancer (Table 2 [28,30,40–49]). A recent exploratory analysis evaluating HRR mutations other than BRCA demonstrated that only a small percentage (4%) of patients with HRD-positive tumours without a BRCA mutation harbour other non-BRCA HRR mutations commonly included in the HRR mutation testing panel [30]. A PFS benefit with olaparib plus bevacizumab versus placebo plus bevacizumab was not observed in the subgroup of patients with non-BRCA HRR mutations; however, in a small number

Table 1
HRD diagnostics for PARPis.

Diagnostic	Sample used	Biomarkers evaluated	Cancer type (regulatory-approved PARPi)	Test category
FoundationOne®CDx is a comprehensive genomic profiling test that detects substitutions, insertion and deletion alterations and copy number alterations in 324 genes and select gene rearrangements, including a number of HRR genes, as well as genomic signatures including MSI and TMB. LOH is evaluated for patients with ovarian cancer. FoundationOne CDx defines positive HRD status as a BRCA mutation in the tumour and/or LOH-high. On this device, assessment of LOH is designated by the FDA as a complementary, not companion, diagnostic [33]	DNA from FFPE tumour tissue	<i>BRCA1</i> and/or <i>BRCA2</i> alterations HRR genes (<i>BRCA1</i> , <i>BRCA2</i> , <i>ATM</i> , <i>BARD1</i> , <i>BRIP1</i> , <i>CDK12</i> , <i>CHEK1</i> , <i>CHEK2</i> , <i>FANCL</i> , <i>PALB2</i> , <i>RAD51B</i> , <i>RAD51C</i> , <i>RAD51D</i> and <i>RAD54L</i>) alteration	Ovarian cancer (olaparib, rucaparib) Prostate cancer (olaparib)	HRR mutation test
FoundationOne®LiquidCDx evaluates specific HRR gene variants in circulating cell-free DNA [34]	cfDNA isolated from plasma derived from anticoagulated peripheral whole blood	<i>BRCA1</i> and <i>BRCA2</i> alterations <i>BRCA1</i> , <i>BRCA2</i> and <i>ATM</i> alterations <i>BRCA1</i> and <i>BRCA2</i> alterations	Ovarian cancer (rucaparib) Prostate cancer (olaparib) Prostate cancer (rucaparib)	HRR mutation test
Myriad BRACAnalysisCDx® evaluates <i>BRCA1</i> and <i>BRCA2</i> variants in genomic DNA from blood [35]	Genomic DNA from whole blood specimens	Deleterious or suspected deleterious mutations in <i>BRCA1</i> and <i>BRCA2</i> genes Deleterious or suspected deleterious mutations in <i>BRCA1</i> and <i>BRCA2</i> genes Deleterious or suspected deleterious mutations in <i>BRCA1</i> and <i>BRCA2</i> genes Deleterious or suspected deleterious mutations in <i>BRCA1</i> and <i>BRCA2</i> genes	Breast cancer (olaparib, talazoparib) Ovarian cancer (olaparib or rucaparib) Pancreatic cancer (olaparib) Prostate cancer (olaparib)	BRCA (HRR) mutation test
Myriad MyChoice®CDx is a comprehensive genomic profiling test specifically for determining HRD status in ovarian cancer [36]	DNA from FFPE tumour tissue	Myriad HRD (defined as deleterious or suspected deleterious mutations in <i>BRCA1</i> and <i>BRCA2</i> genes and/or positive Genomic Instability Score)	Ovarian cancer (olaparib, niraparib)	BRCA (HRR) mutation test and HRD genomic instability test

(continued on next page)

Table 1 (continued)

Diagnostic	Sample used	Biomarkers evaluated	Cancer type (regulatory-approved PARPi)	Test category
CE-IVD (European Union)^a				
Diagnostic	Sample used	Biomarkers evaluated	Cancer type	Test category
Myriad MyChoice [®] CDx PLUS is a next-generation sequencing based in vitro diagnostic device that provides sequencing and large rearrangement analyses on a panel of genes and/or detects genomic instability [37]	DNA from FFPE tumour tissue	Tumour genomic instability and/or detect sequence variants and large rearrangements in up to 15 genes (<i>ATM</i> , <i>BARD1</i> , <i>BRCA1</i> , <i>BRCA2</i> , <i>BRIPI</i> , <i>CDK12</i> , <i>CHEK1</i> , <i>CHEK2</i> , <i>FANCL</i> , <i>PALB2</i> , <i>PPP2R2A</i> , <i>RAD51B</i> , <i>RAD51C</i> , <i>RAD51D</i> and <i>RAD54L</i>)	Ovarian cancer	HRR mutation test and HRD genomic instability test
SOPHiA DDM [™] Dx Homologous Recombination Deficiency Solution ^b is a deep learning–powered diagnostic application leveraging low-pass whole genome sequencing in conjunction with a convolutional neural network–based deep-learning algorithm to produce the Genomic Integrity Index. This score measures the extent of genomic scarring across the genome as a result of mutations within genes associated with HRR, yielding HRD impact on tumour samples [38]	DNA from FFPE tumour tissue	<i>AKT1</i> , ^c <i>ATM</i> , <i>BARD1</i> , <i>BRCA1</i> , <i>BRCA2</i> , <i>BRIPI</i> , <i>CCNE1</i> , <i>CDK12</i> , <i>CHEK1</i> , <i>CHEK2</i> , <i>ESR1</i> , ^c <i>FANCA</i> , <i>FANCD2</i> , <i>FANCL</i> , <i>FGFR1</i> , ^c <i>FGFR2</i> , ^c <i>FGFR3</i> , ^c <i>MRE11</i> , <i>NBN</i> , <i>PALB2</i> , <i>PIK3CA</i> , ^c <i>PPP2R2A</i> , <i>PTEN</i> , <i>RAD51B</i> , <i>RAD51C</i> , <i>RAD51D</i> , <i>RAD54L</i> and <i>TP53</i>	Ovarian cancer	HRR mutation test and HRD genomic instability test
AmoyDx [®] HRD Focus Panel is a next-generation sequencing-based in vitro diagnostic assay intended for qualitative determination of HRD status via detection and classification of single nucleotide variants and insertions and deletions in protein coding regions and intron/exon boundaries of the <i>BRCA1</i> and <i>BRCA2</i> genes and the determination of the Genomic Scar Score, which is an algorithmic measurement of genomic instability status [39]	DNA from FFPE tumour tissue	<i>BRCA1</i> , <i>BRCA2</i> and Genomic Scar Score	Ovarian cancer	BRCA (HRR) mutation test and HRD genomic instability test

Abbreviations: CE-IVD, CE-marked in vitro diagnostics; FFPE, formalin-fixed paraffin embedded; HRD, homologous recombination deficiency; HRR, homologous recombination repair; LOH, loss of heterozygosity; LST, large-scale state transitions; MSI, microsatellite instability; NGS, next-generation sequencing; PARPi, poly (ADP-ribose) polymerase inhibitor; TAI, telomeric allelic imbalance; TMB, tumour mutational burden.

^a A number of *BRCA1/BRCA2* tests are also designated CE-IVD.

^b HRR gene panel analysis of this test is not covered by the CE-IVD claim.

^c Hotspot coverage only.

Table 2

Clinical trials of PARPis in ovarian cancer evaluating the predictive value of BRCA mutations, HRR mutations or HRD status.

Study	Study design	Treatments	Inclusion criteria	HRRm or HRD status	Primary endpoint
First-line maintenance therapy					
PAOLA-1/ENGOT-ov25 (NCT02477644) [44]	Phase 3, randomised, double-blind, multicenter	Olaparib 300 mg BID + bevacizumab (<i>n</i> = 537) versus placebo + bevacizumab (<i>n</i> = 269)	Newly diagnosed, advanced high-grade serous or endometrioid ovarian, primary peritoneal or fallopian tube cancer; complete or partial response after platinum-taxane chemotherapy plus bevacizumab	Deleterious germline <i>BRCA1</i> or <i>BRCA2</i> mutation (MyChoice® HRD Plus assay) HRD-positive defined as HRD score ≥ 42	<u>PFS</u> Overall: 22.1 versus 16.6 mo (HR 0.59; 95% CI: 0.49–0.72) BRCA-mutated HRD-positive: 37.2 versus 17.7 mo (HR 0.33; 95% CI: 0.25–0.45) Non-BRCA HRD-positive: 28.1 versus 16.6 mo (HR 0.43; 95% CI: 0.28–0.66)
SOLO1 (NCT01844986) [45]	Phase 3, randomised, double-blind, multicenter	Olaparib 300 mg BID (<i>n</i> = 260) versus placebo (<i>n</i> = 131)	Newly diagnosed, advanced high-grade serous or endometrioid ovarian, primary peritoneal or fallopian tube cancer; complete or partial response after platinum-based chemotherapy without bevacizumab	Deleterious or suspected deleterious germline or somatic <i>BRCA1</i> and/or <i>BRCA2</i> mutation (BRCAAnalysis test)	<u>PFS</u> NR versus 13.8 mo (HR 0.30; 95% CI: 0.23–0.41) <u>PFS rate at 3 years</u> 60% versus 27%
PRIMA/ENGOT-OV26/ GOG-3012 (NCT02655016) [28]	Phase 3, randomised, double-blind, multicenter	Niraparib 300 mg/day (200 mg/day if bodyweight <77 kg; <i>n</i> = 487) versus placebo (<i>n</i> = 246)	Newly diagnosed, advanced high-grade serous or endometrioid ovarian, peritoneal or fallopian tube cancer; complete or partial response after platinum-based chemotherapy	HRD-positive defined as HRD score ≥ 42 and/or a deleterious BRCA mutation (MyChoice test)	<u>PFS</u> Overall: 13.8 versus 8.2 mo (HR 0.62; 95% CI: 0.50–0.76) HRD-positive: 21.9 versus 10.4 mo (HR 0.43; 95% CI: 0.31–0.59)
ATHENA-MONO/GOG-3020/ENGOT-ov45 (NCT03522246) [46]	Phase 3, randomised, double-blind, multicenter	Rucaparib 600 mg BID (<i>n</i> = 427) versus placebo (<i>n</i> = 111)	Newly diagnosed, advanced high-grade epithelial ovarian, fallopian tube or primary peritoneal cancer; complete or partial response after platinum-based chemotherapy	Tumour HRD test status (BRCA mutations and genomic LOH; FoundationOne CDx)	<u>PFS</u> ITT: 20.2 versus 9.2 mo (HR 0.52; 95% CI: 0.40–0.68) HRD-positive: 28.7 versus 11.3 mo (HR, 0.47; 95% CI: 0.31–0.72)
Relapsed/recurrent treatment strategies					
QUADRA (NCT02354586) [47]	Phase 2, open-label, single-arm, multicenter	Niraparib 300 mg QD (<i>N</i> = 463)	Metastatic, relapsed high-grade serous epithelial ovarian, fallopian tube or primary peritoneal cancer; ≥ 3 prior lines of chemotherapy	Germline <i>BRCA1</i> and/or <i>BRCA2</i> variants and HRD score (MyChoice HRD test)	<u>PFS</u> HRD-positive tumours sensitive to platinum-based therapy and naive to PARPi: 5.5 mo
Study 19 (NCT00753545) [40]	Phase 2, randomised, double-blind, multicenter	Olaparib 400 mg BID (<i>n</i> = 136) versus placebo (<i>n</i> = 129)	Recurrent high-grade serous ovarian, fallopian tube or primary peritoneal cancer; platinum-sensitive; ≥ 2 courses of platinum-based chemotherapy	<i>BRCA1</i> and/or <i>BRCA2</i> mutation status not required	<u>PFS</u> ITT: 8.4 versus 4.8 mo (HR 0.35; 95% CI: 0.25–0.49) BRCA-mutated: HR 0.18 (95% CI: 0.10–0.31) Non-BRCA–mutated: HR 0.54 (95% CI: 0.34–0.85) Non-BRCA–mutated HRD-positive: HR 0.48 (95% CI: 0.18–1.27) Non-BRCA–mutated HRD-negative: HR 0.60 (95% CI: 0.31–1.17)
SOLO2/ENGOT-Ov21 (NCT01874353) [30]	Phase 3, randomised, double-blind, multicenter	Olaparib 300 mg BID (<i>n</i> = 196) versus placebo (<i>n</i> = 99)	Relapsed, high-grade serous or endometrioid ovarian, fallopian tube or primary peritoneal cancer; platinum-sensitive; ≥ 2 prior lines of platinum-based chemotherapy; <i>BRCA1</i> and/or <i>BRCA2</i> mutation	Deleterious <i>BRCA1</i> and/or <i>BRCA2</i> mutation (BRCAAnalysis assay)	<u>PFS</u> 19.1 versus 5.5 mo (HR 0.30; 95% CI: 0.22–0.41)

(continued on next page)

Table 2 (continued)

Study	Study design	Treatments	Inclusion criteria	HRRm or HRD status	Primary endpoint
SOLO3 (NCT02282020) [49]	Phase 3, randomised, open-label, multicenter	Olaparib 300 mg BID (<i>n</i> = 178) versus nonplatinum chemotherapy (<i>n</i> = 88)	Relapsed, high-grade serous or endometrioid ovarian, primary peritoneal and/or fallopian tube cancer; platinum-sensitive; ≥ 2 prior lines of platinum-based chemotherapy; gBRCA mutation	Deleterious or suspected deleterious germline <i>BRCA1</i> or <i>BRCA2</i> mutation (BRCAAnalysis assay)	<u>PFS</u> 13.4 versus 9.2 mo (HR 0.62; 95% CI: 0.43–0.91)
NOVA/ENGOT-OV16 (NCT01847274) [41]	Phase 3, randomised, double-blind, multicenter	Niraparib 300 mg (gBRCA mutation cohort, <i>n</i> = 138; non-BRCA cohort, <i>n</i> = 234) versus placebo (gBRCA mutation cohort, <i>n</i> = 65; non-gBRCA mutation cohort, <i>n</i> = 116)	Recurrent, high-grade serous ovarian, fallopian tube or primary peritoneal cancer; platinum-sensitive; ≥ 2 prior lines of platinum-based chemotherapy	Germline BRCA mutation (BRCAAnalysis assay) HRD status also assessed (MyChoice HRD assay)	<u>PFS</u> gBRCA-mutated cohort: 21.0 versus 5.5 mo (HR 0.27; 95% CI: 0.17–0.41) Non-gBRCA–mutated HRD-positive cohort: 12.9 versus 3.8 mo (HR 0.38; 95% CI: 0.24–0.59) Overall non-gBRCA–mutated cohort: 9.3 versus 3.9 mo (HR 0.45; 95% CI: 0.34–0.61)
ARIEL3 (NCT01968213) [42,43]	Phase 3, randomised, double-blind, multicenter	Rucaparib 600 mg BID (<i>n</i> = 375) versus placebo (<i>n</i> = 189)	Recurrent, high-grade serous or endometrioid ovarian, fallopian tube or primary peritoneal cancer; platinum-sensitive; ≥ 2 prior lines of platinum-based chemotherapy	<i>BRCA1</i> or <i>BRCA2</i> mutation non-BRCA HRR-related gene mutation (T5 next-generation sequencing assay; germline mutations identified with BRCAAnalysis CDx test)	<u>Investigator-assessed PFS (primary analysis)</u> Overall: 10.8 versus 5.4 mo (HR 0.36; 95% CI: 0.30–0.45) BRCA mutated cohort: 16.6 versus 5.4 mo (HR 0.23; 95% CI: 0.16–0.34) HRD cohort: 13.6 versus 5.4 mo (HR 0.32; 95% CI: 0.24–0.42) <u>Chemotherapy-free interval (post-progression)</u> Overall: 14.3 versus 8.8 mo (HR 0.43; 95% CI: 0.35–0.53) BRCA-mutated cohort: 20.8 versus 8.7 mo (HR 0.28; 95% CI: 0.19–0.41) HRD cohort: 18.0 versus 9.1 mo (HR 0.40; 95% CI: 0.31–0.53) <u>Investigator-assessed PFS2 (post-progression)</u> Overall: 21.0 versus 16.5 mo (HR 0.66; 95% CI: 0.53–0.82) BRCA-mutated cohort: 26.8 versus 18.4 mo (HR 0.56; 95% CI: 0.38–0.83) HRD cohort: 25.3 versus 18.4 mo (HR 0.66; 95% CI: 0.49–0.87)
ARIEL4 (NCT02855944) [48]	Phase 3, randomised, open-label, multicenter	Rucaparib 600 mg BID (<i>n</i> = 233) versus chemotherapy (<i>n</i> = 116)	Relapsed, high-grade epithelial ovarian, fallopian tube or primary peritoneal cancer; PARPi-naive; ≥ 2 prior lines of chemotherapy	Deleterious <i>BRCA1</i> and/or <i>BRCA2</i> mutation <i>BRCA</i> reversion mutations were also assessed	<u>PFS</u> Significantly longer for rucaparib versus chemotherapy <i>BRCA</i> reversion mutation subgroup: 2.9 versus 5.5 mo (HR 2.769; 95% CI: 0.989–7.755)

Abbreviations: BID, twice daily; gBRCA germline BRCA mutation; HR, hazard ratio; HRD, homologous recombination deficiency; HRRm, homologous recombination repair gene mutation; ITT, intention-to-treat; PARPi, poly (ADP-ribose) polymerase inhibitor; PFS, progression-free survival.

of these patients, HRD-positive status by genomic instability testing was predictive of a PFS benefit [30].

In patients with relapsed platinum-sensitive ovarian cancer, HRR mutation, HRD status and response to immediate platinum chemotherapy are all predictive of a PFS benefit with PARPis, as demonstrated by the findings from four clinical trials (Table 2 [28,30,40–49]).

The European Society for Medical Oncology Translational Research and Precision Medicine Working Group recently assessed evidence of predictive biomarkers in high-grade serous ovarian cancer and recommended the use of germline or somatic BRCA mutation testing and a validated scar-based (genomic instability) test for HRD to select patients for PARPi treatment in the first-line maintenance and in the platinum-sensitive relapsed settings [3]. This group concluded that evidence is currently insufficient to determine the clinical validity of non-BRCA HRR mutation, next-generation sequencing-based mutational signature, *BRCA1* or *RAD51C* promoter methylation and functional assays for predicting a PARPi response [3]. Data on the clinical validity of somatic BRCA mutations, HRR mutations and HRD genomic instability tests in other cancers besides mCRPC are limited.

Olaparib, rucaparib and niraparib have demonstrated antitumour activity in people with BRCA-mutated mCRPC [50,51]. The PROfound and TRITON2 trials explored treatment outcomes in patients with mutations in other HRR genes [52,53]. PROfound was a randomised, open-label phase 3 trial evaluating olaparib versus physician's choice of hormone therapy (enzalutamide or abiraterone) in mCRPC with a qualifying alteration in any one of 15 HRR genes [52]. In patients with ≥ 1 alteration in *BRCA1*, *BRCA2* or *ATM*, PFS was significantly prolonged for the olaparib versus the control group (median, 7.4 versus 3.6 months; hazard ratio [HR] 0.34; 95% CI: 0.25–0.47) and with statistically significant OS benefit (median, 19.1 versus 14.7 months; HR 0.69; 95% CI: 0.50–0.97; $p = 0.0175$) [54]. A PFS benefit was also observed in the overall population (any HRR mutation) with an HR of 0.49 (95% CI: 0.38–0.63), and a trend toward improved OS was also seen (HR 0.79; 95% CI: 0.61–1.03; nominal $p = 0.0515$). The HR for PFS in patients with HRR mutations other than *BRCA1*, *BRCA2* or *ATM* was 0.88. In the phase 2 TRITON2 trial, rucaparib was evaluated in patients with mCRPC and a non-BRCA HRR mutation. The trial found limited radiographic and prostate-specific antigen responses to rucaparib in patients with an alteration in *ATM*, *CDK12* or *CHEK2*; patients with alterations in other HRR mutations (e.g. *PALB2*) may benefit from rucaparib, but the patient subgroup sizes were too small to make definitive conclusions; further investigation was deemed warranted [53].

In advanced/metastatic breast cancer, olaparib has demonstrated a PFS benefit versus chemotherapy in patients with a germline BRCA mutation in two randomised,

open-label phase 3 clinical trials [55,56]. In OlympiAD, olaparib significantly prolonged median PFS versus chemotherapy (7.0 versus 4.2 months; HR 0.58; 95% CI: 0.43–0.80) in patients with a germline BRCA mutation and human epidermal growth factor receptor type 2-negative metastatic breast cancer treated with ≤ 2 previous chemotherapies. Olaparib has also been associated with a median PFS of 6.3 months (90% CI: 4.4–not available) in 16 patients with metastatic breast cancer and a somatic BRCA mutation [57]. In the EMBRACA study, talazoparib significantly prolonged median PFS versus chemotherapy in patients with a germline BRCA mutation and advanced breast cancer treated with ≤ 3 previous chemotherapies (8.6 versus 5.6 months; HR 0.54; 95% CI: 0.41–0.71) [56].

Approval of olaparib maintenance therapy for pancreatic cancer was based on results from the pivotal phase 3 POLO trial in patients with a germline *BRCA1* or *BRCA2* mutation and metastatic disease that had not progressed during first-line platinum-based chemotherapy. A PFS benefit in favour of olaparib versus placebo was reported (median, 7.4 versus 3.8 months; HR 0.53; 95% CI: 0.35–0.82) [58].

6. Conclusions

Patients with HRD tumours caused by a BRCA mutation represent the subgroup with the best-documented clinical benefit from a PARPi, and whether non-BRCA HRR mutations or HRD genomic instability tests best predict PARPi response by disease site requires further research in many indications. BRCA mutations remain the best-characterised measure of HRD and should always be assessed regardless of other assays that might be used (e.g. assessment of other HRR-related genes or HRD genomic instability). However, patients with HRD tumours without a BRCA mutation can benefit from PARPis in first-line ovarian cancer. HRD is a phenotype that can be measured in different ways, such as by identifying genomic scars or mutational signatures or by measuring HRR function. Further, HRD genomic instability gives a historical rather than a functional view of the genome, hence the interest in functional assays such as that for RAD51 foci. However, varying degrees of evidence exist for the predictive value of these different measures. Inconsistent use of terminology adds to the confusion over these tests. The regulatory-approved companion diagnostics are specific to each PARPi, tumour type, and treatment setting. HRR mutations and HRD genomic instability tests have different predictive power in different clinical settings and are not interchangeable. In newly diagnosed advanced ovarian cancer, HRD positivity by genomic instability tests, but not non-BRCA HRR mutations, are predictive of benefit from PARPi maintenance treatment after response to first-line platinum chemotherapy. In the relapsed setting, selection for benefit

from PARPi is based on platinum sensitivity. HRR mutation tests have been validated in patients with prostate cancer whereas there are few data on the clinical validity of HRD genomic instability testing. Future clinical and translational studies are needed to further elucidate the mechanisms by which PARPis interact with HRD to inform treatment strategies and patient selection.

Funding

This work was funded by Merck Sharp & Dohme LLC, a subsidiary of Merck & Co., Inc., Rahway, NJ, USA, and AstraZeneca, Cambridge, UK.

Data sharing

Merck Sharp & Dohme LLC, a subsidiary of Merck & Co., Inc., Rahway, NJ, USA (MSD) is committed to providing qualified scientific researchers access to anonymised data and clinical study reports from the company's clinical trials for the purpose of conducting legitimate scientific research. MSD is also obligated to protect the rights and privacy of trial participants and, as such, has a procedure in place for evaluating and fulfilling requests for sharing company clinical trial data with qualified external scientific researchers. The MSD data sharing website (available at: http://engagezone.msd.com/ds_documentation.php) outlines the process and requirements for submitting a data request. Applications will be promptly assessed for completeness and policy compliance. Feasible requests will be reviewed by a committee of MSD subject matter experts to assess the scientific validity of the request and the qualifications of the requestors. In line with data privacy legislation, submitters of approved requests must enter into a standard data-sharing agreement with MSD before data access is granted. Data will be made available for request after product approval in the US and EU or after product development is discontinued. There are circumstances that may prevent MSD from sharing requested data, including country or region-specific regulations. If the request is declined, it will be communicated to the investigator. Access to genetic or exploratory biomarker data requires a detailed, hypothesis-driven statistical analysis plan that is collaboratively developed by the requestor and MSD subject matter experts; after approval of the statistical analysis plan and execution of a data-sharing agreement, MSD will either perform the proposed analyses and share the results with the requestor or will construct biomarker covariates and add them to a file with clinical data that is uploaded to an analysis portal so that the requestor can perform the proposed analyses.

Author contributions

Thomas J. Herzog: Formal analysis; validation; draft writing; other – critically reviewing or revising the

manuscript for important intellectual content and final approval.

Ignace Vergote: conceptualisation; validation; draft writing; other – critically reviewing or revising the manuscript for important intellectual content and final approval.

Leonard G. Gomella: conceptualisation; validation; other – critically reviewing or revising the manuscript for important intellectual content and final approval.

Tsveta Milenkova: conceptualisation; validation; other – critically reviewing or revising the manuscript for important intellectual content and final approval.

Tim French: conceptualisation; formal analysis; validation; draft writing; other – critically reviewing or revising the manuscript for important intellectual content and final approval.

Raffi Tonikian: formal analysis; other – critically reviewing or revising the manuscript for important intellectual content and final approval.

Christian Poehlein: conceptualisation; data curation; formal analysis; validation; draft writing; other – critically reviewing or revising the manuscript for important intellectual content and final approval.

Maha Hussain: other – critically reviewing or revising the manuscript for important intellectual content and final approval.

Conflict of interest statement

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: TH reports scientific advisory boards for AstraZeneca, Caris, Clovis, Eisai, Epsilogen, Genentech, GSK, Immunogen, J&J, Merck, Mersana, and Seagen.

IV reports consulting for Agenus, Akesobio China, Amgen (Europe) GmbH, AstraZeneca, Bristol Myers Squibb, Clovis Oncology Inc., Carrick Therapeutics, Deciphera Pharmaceuticals, Eisai, Elevar Therapeutics, F. Hoffmann-La Roche Ltd, Genmab, GSK, Immunogen Inc., Jazzpharma, Karyopharm, Mersana, Millennium Pharmaceuticals, MSD, Novocure, Novartis, Octimet Oncology NV, Oncoinvent AS, Seagen, Sotio a.s., Verastem Oncology and Zentalis; contracted research (via KU Leuven) for Oncoinvent AS and Genmab; grant (i.e. corporate sponsored research) for Amgen and Roche and accommodations and/or travel expenses for Amgen, MSD, Tesaro, AstraZeneca and Roche.

LGG reports scientific advisory boards for AstraZeneca, Merck, Lantheus and Exelixis.

TM is an employee of AstraZeneca.

TF is an employee and stockholder of AstraZeneca.

RT is a former employee of Merck Sharp & Dohme LLC, a subsidiary of Merck & Co., Inc., Rahway, NJ, USA, and stockholder of Merck & Co., Inc., Rahway, NJ, USA.

CP is an employee of Merck Sharp & Dohme LLC, a subsidiary of Merck & Co., Inc., Rahway, NJ, USA, and stockholder of Merck & Co., Inc., Rahway, NJ, USA.

MH reports scientific advisory boards for AstraZeneca, Merck, Janssen, Novartis, TEMPUS, Bayer, GSK, Pfizer and BMS.

Acknowledgements

This work was supported by Merck Sharp & Dohme LLC, a subsidiary of Merck & Co., Inc., Rahway, NJ, USA, and AstraZeneca LTD, Cambridge, UK, which are codeveloping olaparib. The funders participated in study design, data analysis and interpretation, and article writing. All authors had full access to the data and had final responsibility for the decision to submit the article for publication. Medical writing and/or editorial assistance was provided by Lei Bai, PhD, Max Chang, PhD, Holly C. Cappelli, PhD, CMPP, and Christina Vitolo of ApotheCom (Yardley, PA) and was funded by Merck Sharp & Dohme LLC, a subsidiary of Merck & Co., Inc., Rahway, NJ, USA, and AstraZeneca LTD, Cambridge, UK, which are codeveloping olaparib.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ejca.2022.10.021>.

References

- Lord CJ, Ashworth A. The DNA damage response and cancer therapy. *Nature* 2012;481:287–94.
- Jiang M, Jia K, Wang L, Li W, Chen B, Liu Y, et al. Alterations of DNA damage repair in cancer: from mechanisms to applications. *Ann Transl Med* 2020;8:1685.
- Miller RE, Leary A, Scott CL, Serra V, Lord CJ, Bowtell D, et al. ESMO recommendations on predictive biomarker testing for homologous recombination deficiency and PARP inhibitor benefit in ovarian cancer. *Ann Oncol* 2020;31:1606–22.
- Glaxo Smithkline. ZEPHYRUS (niraparib) capsules, for oral use. 9/2022. Research Triangle Park, NC: Glaxo SmithKline, Inc.; 2022.
- AstraZeneca Pharmaceuticals LP. LYNPARZA® (olaparib) tablets, for oral use. 8/2022. Wilmington, DE: AstraZeneca Pharmaceuticals LP; 2022.
- Clovis Oncology I. RUBRACA® (rucaparib) tablets, for oral use. 3/2022. Boulder, CO: Clovis Oncology, Inc.; 2022.
- Pfizer Labs I. TALZENNA- talazoparib capsule. 9/2021. Ny, NY: Pfizer Labs, Inc.; 2021.
- Limited GI. Zephyrus 100 mg hard capsules (SPC). 8/2022. Dublin, Ireland: GlaxoSmithKline (Ireland) Limited; 2022.
- AB A. Lynparza 100 mg film-coated tablets (SPC) Lynparza 150 mg film-coated tablets (SPC). 10/2022. Sweden: AstraZeneca AB: Södertälje; 2022.
- Ltd COI. Rubraca 200 mg film-coated tablets (SPC) Rubraca 250 mg film-coated tablets (SPC) Rubraca 300 mg film-coated tablets (SPC). 5/2022. Dublin, Ireland: Clovis Oncology Ireland Ltd; 2022.
- EEIG PEM. Talzenna 0.25 mg hard capsules (SPC) Talzenna 1 mg hard capsules (SPC). 12/2021. Belgium: Pfizer Europe MA EEIG: Brussels; 2021.
- Pellegrino B, Mateo J, Serra V, Balmaña J. Controversies in oncology: are genomic tests quantifying homologous recombination repair deficiency (HRD) useful for treatment decision making? *ESMO Open* 2019;4:e000480.
- Mersch J, Jackson MA, Park M, Nebgen D, Peterson SK, Singletary C, et al. Cancers associated with BRCA1 and BRCA2 mutations other than breast and ovarian. *Cancer* 2015;121:269–75.
- Konstantinopoulos PA, Ceccaldi R, Shapiro GI, D'Andrea AD. Homologous recombination deficiency: exploiting the fundamental vulnerability of ovarian cancer. *Cancer Discov* 2015;5:1137–54.
- Castedo M, Perfettini JL, Roumier T, Andreau K, Medema R, Kroemer G. Cell death by mitotic catastrophe: a molecular definition. *Oncogene* 2004;23:2825–37.
- Frey MK, Pothuri B. Homologous recombination deficiency (HRD) testing in ovarian cancer clinical practice: a review of the literature. *Gynecol Oncol Res Pract* 2017;4:4.
- Abkevich V, Timms KM, Hennessy BT, Potter J, Carey MS, Meyer LA, et al. Patterns of genomic loss of heterozygosity predict homologous recombination repair defects in epithelial ovarian cancer. *Br J Cancer* 2012;107:1776–82.
- Birkbak NJ, Wang ZC, Kim JY, Eklund AC, Li Q, Tian R, et al. Telomeric allelic imbalance indicates defective DNA repair and sensitivity to DNA-damaging agents. *Cancer Discov* 2012;2:366–75.
- Popova T, Manié E, Rieunier G, Caux-Moncoutier V, Tirapo C, Dubois T, et al. Ploidy and large-scale genomic instability consistently identify basal-like breast carcinomas with BRCA1/2 inactivation. *Cancer Res* 2012;72:5454–62.
- Kim G, Ison G, McKee AE, Zhang H, Tang S, Gwise T, et al. FDA approval summary: olaparib monotherapy in patients with deleterious germline BRCA-mutated advanced ovarian cancer treated with three or more lines of chemotherapy. *Clin Cancer Res* 2015;21:4257–61.
- Mateo J, Lord CJ, Serra V, Tutt A, Balmaña J, Castroviejo-Bermejo M, et al. A decade of clinical development of PARP inhibitors in perspective. *Ann Oncol* 2019;30:1437–47.
- Armstrong N, Ryder S, Forbes C, Ross J, Quek RG. A systematic review of the international prevalence of BRCA mutation in breast cancer. *Clin Epidemiol* 2019;11:543–61.
- Doan DK, Schmidt KT, Chau CH, Figg WD. Germline genetics of prostate cancer: prevalence of risk variants and clinical implications for disease management. *Cancers* 2021;13.
- Rosen MN, Goodwin RA, Vickers MM. BRCA mutated pancreatic cancer: a change is coming. *World J Gastroenterol* 2021;27:1943–58.
- Grellety T, Peyraud F, Sevenet N, Tredan O, Dohollou N, Barouk-Simonet E, et al. Dramatic response to PARP inhibition in a PALB2-mutated breast cancer: moving beyond BRCA. *Ann Oncol* 2020;31:822–3.
- Pilié PG, Gay CM, Byers LA, O'Connor MJ, Yap TA. PARP inhibitors: extending benefit beyond BRCA-mutant cancers. *Clin Cancer Res* 2019;25:3759–71.
- Kuemmel S, Harrach H, Schmutzler RK, Kostara A, Ziegler-Löhr K, Dyson MH, et al. Olaparib for metastatic breast cancer in a patient with a germline PALB2 variant. *NPJ Breast Cancer* 2020;6:31.
- González-Martín A, Pothuri B, Vergote I, DePont Christensen R, Graybill W, Mirza MR, et al. Niraparib in patients with newly diagnosed advanced ovarian cancer. *N Engl J Med* 2019;381:2391–402.
- Li N, Zhu J, Yin R, Wang J, Zhu J, Pan L, et al. Efficacy and safety of niraparib as maintenance treatment in patients with newly diagnosed advanced ovarian cancer using an individualized starting dose (PRIME study): a randomized, double-blind, placebo-controlled, phase 3 trial. SGO Annual Meeting on Women's cancer. Abstract 244. Presented March 19, 2022. XX; 2022.

- [30] Pujade-Lauraine E, Brown J, Barnicle A, Rowe P, Lao-Sirieix P, Criscione S, et al. Homologous recombination repair mutation gene panels (excluding BRCA) are not predictive of maintenance olaparib plus bevacizumab efficacy in the first-line PAOLA-1/ENGOT-ov25 trial. *Gynecol Oncol* 2021;162:S26–7.
- [31] Blanc-Durand F, Yaniz E, Genestie C, Rouleau E, Berton D, Lortholary A, et al. Evaluation of a RAD51 functional assay in advanced ovarian cancer, a GINECO/GINEGEPS study. *J Clin Oncol* 2021;39:5513.
- [32] Waks AG, Cohen O, Kochupurakkal B, Kim D, Dunn CE, Buendia Buendia J, et al. Reversion and non-reversion mechanisms of resistance to PARP inhibitor or platinum chemotherapy in BRCA1/2-mutant metastatic breast cancer. *Ann Oncol* 2020;31:590–8.
- [33] Foundation Medicine Inc. FoundationOne®CDx technical information. Cambridge, MA: Foundation Medicine Inc.; 2021.
- [34] Foundation Medicine Inc. FoundationONE Liquid CDx technical information. Cambridge, MA: Foundation Medicine Inc.; 2021.
- [35] Myriad Genetics Laboratories Inc. BRACAnalysis CDx technical information. Salt Lake City, UT: Myriad Genetics Laboratories, Inc.; 2021.
- [36] Myriad genetic Laboratories Inc. MyChoice® CDx technical information. Salt Lake City, UT: Myriad Genetics Laboratories, Inc.; 2021.
- [37] Myriad genetic Laboratories Inc. MyChoice® CDx plus technical information. Salt Lake City, UT: Myriad Genetics Laboratories, Inc.; 2022.
- [38] Genetics S. SOPHiA DDM™ Dx homologous recombination deficiency solution. Lausanne, Switzerland: SOPHiA Genetics; 2022.
- [39] Amoy diagnostics Co. L. NGS assays. Xiamen, China: Amoy Diagnostics Co., Ltd.; 2012.
- [40] Ledermann J, Harter P, Gourley C, Friedlander M, Vergote I, Rustin G, et al. Olaparib maintenance therapy in platinum-sensitive relapsed ovarian cancer. *N Engl J Med* 2012;366:1382–92.
- [41] Mirza MR, Monk BJ, Herrstedt J, Oza AM, Mahner S, Redondo A, et al. Niraparib maintenance therapy in platinum-sensitive, recurrent ovarian cancer. *N Engl J Med* 2016;375:2154–64.
- [42] Coleman RL, Oza AM, Lorusso D, Aghajanian C, Oaknin A, Dean A, et al. Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* 2017;390:1949–61.
- [43] Ledermann JA, Oza AM, Lorusso D, Aghajanian C, Oaknin A, Dean A, et al. Rucaparib for patients with platinum-sensitive, recurrent ovarian carcinoma (ARIEL3): post-progression outcomes and updated safety results from a randomised, placebo-controlled, phase 3 trial. *Lancet Oncol* 2020;21:710–22.
- [44] Ray-Coquard I, Pautier P, Pignata S, Pérol D, González-Martín A, Berger R, et al. Olaparib plus bevacizumab as first-line maintenance in ovarian cancer. *N Engl J Med* 2019;381:2416–28.
- [45] Moore K, Colombo N, Scambia G, Kim BG, Oaknin A, Friedlander M, et al. Maintenance olaparib in patients with newly diagnosed advanced ovarian cancer. *N Engl J Med* 2018;379:2495–505.
- [46] Monk BJ, Parkinson C, Lim MC, O'Malley DM, Oaknin A, Wilson MK, et al. A randomized, phase III trial to evaluate rucaparib monotherapy as maintenance treatment in patients with newly diagnosed ovarian cancer (ATHENA-MONO/GOG-3020/ENGOT-ov45). *J Clin Oncol* 2022;Jco2201003.
- [47] Moore KN, Secord AA, Geller MA, Miller DS, Cloven N, Fleming GF, et al. Niraparib monotherapy for late-line treatment of ovarian cancer (QUADRA): a multicentre, open-label, single-arm, phase 2 trial. *Lancet Oncol* 2019;20:636–48.
- [48] Kristeleit R. Rucaparib versus chemotherapy in patients with advanced, relapsed ovarian cancer and a deleterious BRCA mutation: efficacy and safety from ARIEL4, a randomized phase III study. SGO Virtual Annual Meeting on Women's cancer; 2021; Virtual: March 19-25. 2021.
- [49] Penson RT, Valencia RV, Cibula D, Colombo N, Leath CA, Bidzinski M, et al. Olaparib versus nonplatinum chemotherapy in patients with platinum-sensitive relapsed ovarian cancer and a germline BRCA1/2 mutation (SOLO3): a randomized phase III trial. *J Clin Oncol* 2020;38:1164–74.
- [50] Mateo J, Porta N, Bianchini D, McGovern U, Elliott T, Jones R, et al. Olaparib in patients with metastatic castration-resistant prostate cancer with DNA repair gene aberrations (TOPARP-B): a multicentre, open-label, randomised, phase 2 trial. *Lancet Oncol* 2020;21:162–74.
- [51] Smith MR, Scher HI, Sandhu S, Efstathiou E, Lara Jr PN, Yu EY, et al. Niraparib in patients with metastatic castration-resistant prostate cancer and DNA repair gene defects (GALAHAD): a multicentre, open-label, phase 2 trial. *Lancet Oncol* 2022;23:362–73.
- [52] de Bono J, Mateo J, Fizazi K, Saad F, Shore N, Sandhu S, et al. Olaparib for metastatic castration-resistant prostate cancer. *N Engl J Med* 2020;382:2091–102.
- [53] Abida W, Campbell D, Patnaik A, Shapiro JD, Sautois B, Vogelzang NJ, et al. Non-BRCA DNA damage repair gene alterations and response to the PARP inhibitor rucaparib in metastatic castration-resistant prostate cancer: analysis from the phase ii TRITON2 study. *Clin Cancer Res* 2020;26:2487–96.
- [54] de Bono JS, Mateo J, Fizazi K, Saad F, Shore N, Sandhu S, et al. Final overall survival (OS) analysis of PROfound: olaparib vs physician's choice of enzalutamide or abiraterone in patients (pts) with metastatic castration-resistant prostate cancer (mCRPC) and homologous recombination repair (HRR) gene alterations. *Ann Oncol* 2020;S507–49.
- [55] Robson M, Im SA, Senkus E, Xu B, Domchek SM, Masuda N, et al. Olaparib for metastatic breast cancer in patients with a germline BRCA mutation. *N Engl J Med* 2017;377:523–33.
- [56] Litton JK, Rugo HS, Ettl J, Hurvitz SA, Gonçalves A, Lee K-H, et al. Talazoparib in patients with advanced breast cancer and a germline BRCA mutation. *N Engl J Med* 2018;379:753–63.
- [57] Tung NM, Robson ME, Venz S, Santa-Maria CA, Nanda R, Marcom PK, et al. TBCRC 048: phase II study of olaparib for metastatic breast cancer and mutations in homologous recombination-related genes. *J Clin Oncol* 2020;38:4274–82.
- [58] Golan T, Hammel P, Reni M, Van Cutsem E, Macarulla T, Hall MJ, et al. Maintenance olaparib for germline BRCA-mutated metastatic pancreatic cancer. *N Engl J Med* 2019;381:317–27.