

3-15-2015

G protein $\beta\gamma$ subunits regulate cardiomyocyte hypertrophy through a perinuclear Golgi phosphatidylinositol 4-phosphate hydrolysis pathway.

S Malik

Department of Pharmacology and Physiology, University of Rochester

R G deRubio

Department of Pharmacology and Physiology, University of Rochester

M Trembley

Department of Pharmacology and Physiology, University of Rochester

R Irannejad

Department of Psychiatry, University of California, San Francisco

Follow this and additional works at: <https://jdc.jefferson.edu/bmpfp>

 Philip B Wedegaertner

Part of the [Medical Molecular Biology Commons](#)
Department of Biochemistry and Molecular Biology, Thomas Jefferson University

[Let us know how access to this document benefits you](#)

See next page for additional authors

Recommended Citation

Malik, S; deRubio, R G; Trembley, M; Irannejad, R; Wedegaertner, Philip B; and Smrcka, A V, "G protein $\beta\gamma$ subunits regulate cardiomyocyte hypertrophy through a perinuclear Golgi phosphatidylinositol 4-phosphate hydrolysis pathway." (2015). *Department of Biochemistry and Molecular Biology Faculty Papers*. Paper 85.

<https://jdc.jefferson.edu/bmpfp/85>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Biochemistry and Molecular Biology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

S Malik, R G deRubio, M Trembley, R Irannejad, Philip B Wedegaertner, and A V Smrcka

G protein $\beta\gamma$ subunits regulate cardiomyocyte hypertrophy through a perinuclear Golgi phosphatidylinositol 4-phosphate hydrolysis pathway

S. Malik^a, R. G. deRubio^a, M. Trembley^a, R. Irannejad^b, P. B. Wedegaertner^c, and A. V. Smrcka^a

^aDepartment of Pharmacology and Physiology, University of Rochester, Rochester, NY 14642; ^bDepartment of Psychiatry, University of California, San Francisco, San Francisco, CA 94158; ^cDepartment of Biochemistry and Molecular Biology, Thomas Jefferson University, Philadelphia, PA 19107

ABSTRACT We recently identified a novel GPCR-dependent pathway for regulation of cardiac hypertrophy that depends on Golgi phosphatidylinositol 4-phosphate (PI4P) hydrolysis by a specific isoform of phospholipase C (PLC), PLC ϵ , at the nuclear envelope. How stimuli are transmitted from cell surface GPCRs to activation of perinuclear PLC ϵ is not clear. Here we tested the role of G protein $\beta\gamma$ subunits. G $\beta\gamma$ inhibition blocked ET-1-stimulated Golgi PI4P depletion in neonatal and adult ventricular myocytes. Blocking G $\beta\gamma$ at the Golgi inhibited ET-1-dependent PI4P depletion and nuclear PKD activation. Translocation of G $\beta\gamma$ to the Golgi stimulated perinuclear Golgi PI4P depletion and nuclear PKD activation. Finally, blocking G $\beta\gamma$ at the Golgi or PM blocked ET-1-dependent cardiomyocyte hypertrophy. These data indicate that G $\beta\gamma$ regulation of the perinuclear Golgi PI4P pathway and a separate pathway at the PM is required for ET-1-stimulated hypertrophy, and the efficacy of G $\beta\gamma$ inhibition in preventing heart failure maybe due in part to its blocking both these pathways.

Monitoring Editor

Thomas F. J. Martin
University of Wisconsin

Received: Oct 24, 2014

Revised: Dec 23, 2014

Accepted: Jan 14, 2015

INTRODUCTION

G protein-coupled receptors (GPCRs) regulate many aspects of cardiac function, including cardiac contractility and hypertrophic growth. One of the key GPCR-regulated pathways in cardiomyocytes is the phosphoinositide-specific phospholipase C (PI-PLC) signaling pathway, which hydrolyzes plasma membrane (PM) phosphatidylinositol 4,5-bisphosphate (PIP₂) to produce inositol tris-

phosphate (IP₃) and diacylglycerol (DAG). IP₃-dependent calcium release in the nucleus and DAG-dependent activation of protein kinase D (PKD) are two of the critical signals involved in regulation of cardiac hypertrophic growth (Vega *et al.*, 2004; Wu *et al.*, 2006). Gq has been shown to drive cardiac hypertrophy (Knowlton *et al.*, 1993; D'Angelo *et al.*, 1997) and is known to activate PLC β (Smrcka *et al.*, 1991; Taylor *et al.*, 1991). Indeed, PLC β 1 has been directly implicated in α 1-adrenergic receptor-dependent hypertrophic growth of neonatal rat ventricular myocytes (NRVMs; Filtz *et al.*, 2009).

Our laboratory has identified unique roles for a distinct isoform of PI-PLC, PLC ϵ , in regulation of various aspects of cardiac function (Smrcka *et al.*, 2012). PLC ϵ provides a link between β -adrenergic receptor activation and type 2 ryanodine receptor (Ryr2)-dependent calcium release in the regulation of cardiac contractility (Wang *et al.*, 2005; Oestreich *et al.*, 2007; Oestreich *et al.*, 2009). PLC ϵ is also required for development of the pressure overload model of cardiac hypertrophy in mice and is downstream of multiple receptor-dependent stimuli in isolated ventricular myocytes (Zhang *et al.*, 2011, 2013). Like all PI-PLCs, PLC ϵ hydrolyzes PIP₂, but it has distinct regulatory features, in that it is directly activated by small GTPases, including Ras, Rap, and Rho (Smrcka *et al.*, 2012). PLC ϵ can also be activated by G protein $\beta\gamma$ subunits, although direct interactions have not been demonstrated (Wing *et al.*, 2001). These regulatory

This article was published online ahead of print in MBoC in Press (<http://www.molbiolcell.org/cgi/doi/10.1091/mbc.E14-10-1476>) on January 21, 2015.

Address correspondence to: Alan V. Smrcka (Alan_Smrcka@urmc.rochester.edu).

Abbreviations used: ANF, atrial natriuretic factor; AVM, adult ventricular myocyte; cpTOME, 8-(4-chlorophenylthio)-2'-O-methyladenosine-3',5'-monophosphate; DAG, diacylglycerol; Epac, exchange factor activated by cAMP; ET-1, endothelin-1; FAPP-PH, four-phosphate adaptor protein-pleckstrin homology domain; GPCR, G protein-coupled receptor; GRK2ct, C terminus of G protein-coupled receptor kinase 2; mAkap, muscle-specific A kinase anchoring protein; nDKAR, nuclear targeted D kinase reporter; NRVM, neonatal rat ventricular myocyte; PI4P, phosphatidylinositol 4-phosphate; PIP₂, phosphatidylinositol 4,5-bisphosphate, inositol 1,4,5 trisphosphate; PI-PLC, phosphatidylinositol-specific phospholipase C; PKC, protein kinase C; PKD, protein kinase D; PM, plasma membrane; PTX, pertussis toxin.

© 2015 Malik *et al.* This article is distributed by The American Society for Cell Biology under license from the author(s). Two months after publication it is available to the public under an Attribution-Noncommercial-Share Alike 3.0 Unported Creative Commons License (<http://creativecommons.org/licenses/by-nc-sa/3.0>). "ASCB," "The American Society for Cell Biology," and "Molecular Biology of the Cell" are registered trademarks of The American Society for Cell Biology.

Supplemental Material can be found at:
<http://www.molbiolcell.org/content/suppl/2015/01/18/mbc.E14-10-1476v1.DC1.html>

features potentially place it downstream of any signals that can activate these G proteins.

We recently identified a novel mechanism for PLC ϵ -dependent regulation of cardiac hypertrophy (Zhang *et al.*, 2013). PLC ϵ scaffolded at the nuclear envelope in cardiac myocytes by direct binding to muscle-specific A kinase anchoring protein (mAKAP) hydrolyzes perinuclear Golgi phosphatidylinositol 4-phosphate (PI4P) to generate local DAG and regulate nuclear PKD. PKD is a key kinase that phosphorylates histone deacetylase (HDAC) to stimulate hypertrophic gene expression, and deletion of PKD inhibits development of pressure overload and angiotensin II-induced cardiac hypertrophy (Vega *et al.*, 2004; Fielitz *et al.*, 2008). In cardiac myocytes, the Golgi apparatus intimately surrounds the nuclear envelope, and local signals generated at the Golgi in close proximity to the nucleus can regulate nuclear PKD activation. We recently proposed that Golgi-localized perinuclear DAG generated by mAKAP-scaffolded PLC ϵ is necessary to maintain activation of PKD at the nucleus, since DAG is not diffusible from membranes (Zhang *et al.*, 2013). In these studies, treatment of cardiac myocytes with endothelin-1 (ET-1) stimulated the PLC ϵ pathway to regulate nuclear PKD. The question that arises is how the signal is transmitted from PM ET-1A receptors to PLC ϵ at the perinuclear Golgi apparatus. ET-1AR can couple to multiple different G proteins, including G12/13 and Gq, which in turn lead to activation of intracellular signaling proteins such as Ras or Rho that could directly activate PLC ϵ .

A pathway involving PKD activation at the Golgi has been shown to be involved in vesicular transport and secretion in cultured cell lines and involves G protein $\beta\gamma$ subunits (Jamora *et al.*, 1999; Diaz Anel and Malhotra, 2005; Campelo and Malhotra, 2012). Some of these studies suggest that G $\beta\gamma$ directly binds and activates PKD, but there is evidence that G $\beta\gamma$ -dependent regulation of PLC activity is required to produce DAG at the Golgi for PKC and PKD activation (Diaz Anel, 2007; Irannejad and Wedegaertner, 2010). The substrate for PLC in this reaction was presumed to be PIP $_2$ but was not defined. Exactly how G $\beta\gamma$ is regulated at the Golgi is unclear, but it has been shown that G $\beta\gamma$ can translocate from the PM to the Golgi to regulate Golgi fission (Chisari *et al.*, 2007; Saini *et al.*, 2007, 2010).

The similarities between the Golgi trafficking and cardiac hypertrophy pathways and the fact that PLC ϵ can be regulated by G $\beta\gamma$ suggests the idea that G $\beta\gamma$ mediates the signaling downstream of ET-1 to regulate PLC ϵ -dependent PI4P hydrolysis, PKD activation, and cardiomyocyte hypertrophy. Blocking G $\beta\gamma$ signaling with either protein- or small molecule-based inhibitors has profound salutary effects on heart failure *in vivo* and in cardiomyocytes in culture (Koch *et al.*, 1995; Rockman *et al.*, 1998; Casey *et al.*, 2010; Rengo *et al.*, 2011). Here we provide evidence that G $\beta\gamma$ is required for ET-1-dependent perinuclear PI4P hydrolysis, nuclear PKD activation, and NRVM hypertrophy. This indicates that one of the potential mechanisms by which G $\beta\gamma$ blockade inhibits development of heart failure is through inhibition of this novel pathway.

RESULTS

ET-1-dependent perinuclear PI4P hydrolysis relies on G $\beta\gamma$ signaling

As we previously reported, in NRVMs transfected with the PI4P-specific probe green fluorescent protein-four-phosphate adapter protein-pleckstrin homology domain (GFP-FAPP-PH), a perinuclear fluorescent ring is observed by confocal fluorescence microscopy that strongly colocalizes with the Golgi apparatus and is identified with an antibody recognizing a Golgi-specific 58K protein (Figure 1A; Zhang *et al.*, 2013). The localization of PI4P also appears to overlap with mAKAP associated with the nuclear




FIGURE 1: PI4P-associated GFP-FAPP-PH fluorescence colocalizes with a Golgi marker and is closely associated with nuclear envelope-localized mAKAP. (A) NRVMs were transduced with adenovirus expressing GFP-FAPP-PH. Cells were fixed and stained for Golgi-specific 58K protein (Alexa Fluor 546 secondary antibody) and imaged for GFP and red fluorescence. (B) Same as A, except that cells were stained for mAKAP instead of Golgi and treated with and without 10 μ M brefeldin A for 20 min as indicated. (C) Cells were treated for the indicated times with brefeldin A, fixed, and costained for mAKAP and Golgi-specific 58K protein. Scale bars, 10 μ m.

envelope (Figure 1B). Treatment with brefeldin A disrupts the localization of the perinuclear PI4P probe (Figure 1B) and the Golgi-specific 58K protein (Figure 1C) but does not affect the distribution of mAKAP, supporting the idea that PI4P and mAKAP are in distinct but closely associated compartments—Golgi and nuclear envelope—that are not readily distinguished by conventional light microscopy.

To determine whether G $\beta\gamma$ is involved in perinuclear Golgi PI4P hydrolysis, we monitored perinuclear fluorescence intensity associated with GFP-FAPP-PH in NRVMs in the presence and absence of




FIGURE 2: G $\beta\gamma$ inhibitors block ET-1-stimulated perinuclear PI4P depletion in NRVMs. (A) NRVMs transfected with GFP-FAPP-PH were pretreated with 10 μ M gallein or PBS vehicle control, followed by stimulation with 100 nM ET-1. Perinuclear FAPP fluorescence intensity was measured over time by confocal microscopy. (B) NRVMs were treated as in A, except that 10 μ M cpTOME was added to stimulate PI4P hydrolysis. (C) NRVMs were cotransfected with GFP-FAPP-PH and GRK2ct or vehicle control, followed by ET-1 stimulation as in A. (D) NRVMs were transfected as in A were pretreated for 18 h with 100 ng/ml pertussis toxin. All traces represent pooled data from at least three cells from three separate myocyte preparations \pm SEM. The last five values from each time course were averaged and tested for significance using Student's *t* test. ****p* < 0.005; ns, not significant.

ET-1. Stimulation with ET-1 leads to a time-dependent decrease in perinuclear PI4P-associated GFP-FAPP-PH fluorescence, as previously reported (Figure 2A). Pretreatment of NRVMs with 10 μ M gallein, a G $\beta\gamma$ blocker (Bonacci *et al.*, 2006), completely inhibited ET-1-dependent decrease in PI4P fluorescence. To control for potential nonspecific effects of gallein, we treated NRVMs with the Epac activator cpTOME to stimulate PLC ϵ . cpTOME binding to Epac leads to activation of Rap, which in turn directly activates PLC ϵ , resulting in perinuclear PI4P depletion (Bos, 2003; Zhang *et al.*, 2013). Because this pathway is independent of G $\beta\gamma$ signaling, gallein would be predicted to have no effect on cpTOME-dependent depletion of perinuclear PI4P. Treatment of NRVMs with cpTOME leads to PI4P depletion, as previously reported, and pretreatment with 10 μ M gallein does not alter this response (Figure 2B), indicating relative specificity of gallein for ET-1-dependent PI4P depletion.

To provide further evidence for G $\beta\gamma$ signaling controlling perinuclear PI4P hydrolysis, we transfected NRVMs with the C-terminus

of GRK2 (GRK2ct), a well-established G $\beta\gamma$ blocker. In cells transfected with GRK2ct, ET-1-dependent PI4P depletion was completely abolished (Figure 2C). Taken together, these data indicate that G $\beta\gamma$ signaling is required for ET-1-dependent stimulation of perinuclear PI4P hydrolysis. G $\beta\gamma$ signaling is often associated with signaling via Gi proteins, but ET-1A receptors couple primarily to Gq. To determine whether Gi proteins were involved in the ET-1-mediated response, we pretreated cells with pertussis toxin (PTX) before ET-1. PTX had no effect on ET-1-stimulated PI4P hydrolysis, indicating that Gi is not involved in this response.

Subcellular requirement for G $\beta\gamma$ -dependent PI4P hydrolysis

ET-1A receptors are present in the sarcolemmal membrane, whereas ET-1B receptors are present on intracellular membranes in ventricular myocytes (Bkaily *et al.*, 2011). Acute stimulation of NRVMs with the ET-1 peptide, however, should stimulate only receptors that are at least initially present on the cell surface. Previous data with the ET1A receptor-selective blocker BQ-123 indicate that ET-1-dependent PI4P depletion is dependent on the ET-1A receptor (Zhang *et al.*, 2013). Thus the question arises of whether G $\beta\gamma$ at the PM or at the Golgi is required for ET-1-dependent PI4P depletion. The Wedegaertner laboratory developed Golgi- and PM-targeted GRK2ct by fusing GRK2ct with the Golgi-targeting sequence KDELrD193N or the 66-amino acid PM-targeting motif of Rit GTPase (Irannejad and Wedegaertner, 2010). These constructs have been extensively characterized to localize to the appropriate compartments in HeLa cells. When transfected into NRVMs, Golgi GRK2 was clearly targeted to the perinuclear Golgi apparatus and colocalized exclusively with a Golgi marker (Figure 3A).

PM-targeted GRK2ct was more diffuse, but because of the flat nature of these cells, a clear PM-associated fluorescent ring around the periphery of the cell is difficult to observe (Figure 3B); however, PM-targeted GRK2ct had similar distribution as PM-targeted myristoylated and palmitoylated-yellow fluorescent protein (lyn-YFP), which is used extensively as a PM marker in other cell types (Teruel *et al.*, 1999; Figure 3B). It is also clear that the Golgi-targeted GRK2ct shows no PM association and the PM-targeted GRK2ct shows no Golgi localization, speaking to the specificity of these inhibitors for these compartments. When NRVMs were transfected with PM-targeted GRK2ct, there was no effect on ET-1-stimulated Golgi PI4P hydrolysis (Figure 3, C and E). This is in stark contrast to the data in Figure 2C showing that cytosolic GRK2ct expression blocks ET-1-dependent PI4P depletion, indicating that restriction of GRK2ct to the PM prevents access to the pool of G $\beta\gamma$ that is relevant to stimulation of PI4P hydrolysis. On the other hand, transfection of NRVMs with Golgi-GRK2ct significantly inhibited ET-1-dependent

PI4P hydrolysis, indicating that G β γ signaling is required at the Golgi to stimulate perinuclear PI4P hydrolysis (Figure 3, D and E). Western blots of HEK293 cells or NRVMs transfected with PM or Golgi GRK2ct show similar levels of expression (Supplemental Figure S1).

We proposed that the role of perinuclear PI4P hydrolysis is to generate local DAG to maintain activation of a nuclear pool of PKD. PKD is then involved in phosphorylating HDAC and activation of hypertrophic gene expression. Because Golgi-GRK2ct blocked ET-1–dependent PI4P hydrolysis, we predicted that Golgi-GRK2ct would inhibit activation of the nuclear pool of PKD. NRVMs were cotransfected with a fluorescence resonance energy transfer (FRET)–based reporter of PKD activity that is specifically targeted to the nucleus, nDKAR (Kunkel *et al.*, 2007), with either Golgi- or PM-targeted GRK2ct. ET-1 treatment of NRVMs led to a decrease in nDKAR FRET (indicative of phosphorylation by nuclear PKD), which was completely inhibited by cotransfection with Golgi-targeted GRK2ct (Figure 3, F–H). Targeting GRK2ct to the PM partially inhibited nuclear PKD activation (Figure 3, F–H). These data indicate that G β γ at the perinuclear Golgi in cardiac myocytes is required for nuclear PKD activation and that G β γ activation at the PM is also partially required for nuclear PKD activation.

Rapamycin-induced γ translocation to the Golgi induces PI4P hydrolysis and PKD activation

As an alternate approach to examining the role of G β γ in different compartments, we used a rapamycin-inducible translocation system to specifically direct G β γ to either the PM or the Golgi (Irannejad and Wedegaertner, 2010). In this system, FRB is fused to the C-terminus of the G γ subunit and FKBP is fused to either amino acids 1–11 of Lyn for PM targeting or KDELR-D193N for Golgi targeting. To examine the effects of G β γ targeting on perinuclear PI4P hydrolysis, we cotransfected NRVMs with G β γ ₁₂-FRB, GFP-FAPP-PH, and either PM or Golgi FKBP. Rapamycin-inducible translocation of G β γ ₁₂-FRB to either PM (with Lyn-FKBP–cyan fluorescent protein [CFP]) or Golgi (KDELR-D193N-FKBP-CFP) has been demonstrated in HeLa cells (Irannejad and Wedegaertner, 2010). Figure 4A, top, shows that GFP-G β becomes enriched in the perinuclear Golgi with rapamycin treatment. Figure 4A, bottom, shows localization of Golgi-targeted FKBP before and after 20 min of rapamycin treatment. PM-dependent translocation was more difficult to observe, perhaps because of the diffuse PM localization observed in NRVMs (unpublished data; see Figure 2B). For measurement of PI4P depletion, 24 h after cotransfection of FRB and FKBP constructs, rapamycin was added, and perinuclear PI4P-associated GFP-FAPP-PH fluorescence was monitored. Targeting G β γ -FRB to the Golgi produced a strong stimulation of PI4P hydrolysis, whereas PM targeting of G β γ had no effect (Figure 4B). In other cell types, forced translocation of G β γ to the Golgi caused Golgi breakdown and vesiculation that could affect apparent PI4P-associated GFP-FAPP-PH fluorescence (Jamora *et al.*, 1999; Irannejad and Wedegaertner, 2010). As shown in Figure 4A, rapamycin-induced translocation to the Golgi in NRVMs does not affect overall Golgi structure, since the Golgi-targeted KDELR-D193N-FKBP-CFP localization and associated morphology are not altered by rapamycin treatment under conditions in which PI4P fluorescence is dramatically decreased.

Targeting of G β γ -FRB to the Golgi, but not the PM, also stimulated nuclear PKD activation (Figure 4C). Rapamycin alone had no effect on PI4P hydrolysis or nuclear PKD activation (unpublished data). Although rapamycin dependent recruitment of G β γ to the PM is not sufficient to activate nuclear PKD, it is possible that rapamycin-dependent recruitment of G β γ to the PM may not be capable of generating G β γ signaling at the PM. Together these data indicate

that targeting G β γ to the Golgi is sufficient to stimulate perinuclear PI4P hydrolysis and PKD activation.

Golgi-targeted GRK2ct blocks ET-1–stimulated cardiomyocyte hypertrophy

Overexpression of GRK2ct in cardiac myocytes has been well documented to block development of cardiomyocyte hypertrophy *in vitro* and *in vivo* (Koch *et al.*, 1995; Rockman *et al.*, 1998; Rengo *et al.*, 2011). We showed that mAkap-scaffolded PLC ϵ hydrolyzes perinuclear PI4P at the perinuclear Golgi apparatus in cardiac myocytes, and this is required for activation of nuclear PKD and cardiomyocyte growth. The data presented thus far indicate that G β γ can mediate activation of this hypertrophic pathway at the Golgi. This suggests the possibility that one of the mechanisms for the antihypertrophic actions of GRK2ct is to block the Golgi perinuclear PLC ϵ /PI4P/PKD pathway. To test this, we transfected NRVMs with plasmids expressing YFP alone or YFP plus GRK2ct, Golgi GRK2ct, or PM-GRK2ct. Twenty-four hours after infection, cells were treated with ET-1 for 24 h to induce hypertrophy (Figure 5). Because transfection efficiency of NRVMs was low, fluorescent cells were identified and hypertrophy was assessed by staining for increases in intracellular atrial natriuretic factor (ANF) levels with an anti-ANF antibody. ET-1 stimulated a robust increase in the level of ANF staining (Figure 5). In Figure 5A, the GRK2ct-transfected cell (green) shows no ET-1–induced ANF staining (red), whereas a neighboring untransfected cell shows robust ET-1–dependent ANF induction. Figure 5B shows representative images of cells transfected with each of the constructs, and Figure 5C shows quantitation of these data. Surprisingly, expression of either Golgi-GRK2ct or PM-GRK2ct prevented ET-1–dependent ANF induction. As another measure of hypertrophy, the increase in cell size associated with ET-1–stimulated hypertrophy was assessed after adenoviral transduction of NRVMs with YFP, PM-GRK2ct, or Golgi-GRK2ct. Both Golgi- and PM-targeted GRK2ct blocked the ET-1–stimulated increase in cell size (Figure 6).

Neonatal cardiac myocytes are a standard model for examining signaling pathways associated with cardiac hypertrophy, but a more physiological model is the adult ventricular myocyte (AVM). AVMs were isolated from C57Bl6 mice and infected with adenoviruses expressing a PIP₂ detector, tubby-GFP, or a PI4P detector, FAPP-PH-GFP, as was done previously for NRVMs. Cells transfected with Tubby GFP showed localized PIP₂-associated fluorescence at the sarcolemma (Figure 7A, top), whereas FAPP-PH-GFP detected PI4P at the perinuclear Golgi (Figure 7A, bottom). Some FAPP-PH-GFP fluorescence was also observed in a pattern corresponding to Z-lines or T-tubules (Figure 7A, bottom). Thus, as in NRVMs, PIP₂ is enriched at the PM in AVMs and is not observable at the perinuclear Golgi, indicating that the available PLC substrate at the Golgi is PI4P. We tested whether ET-1 treatment would lead to depletion of perinuclear PI4P as seen with NRVMs. ET-1 stimulated a time-dependent decrease in perinuclear PI4P-associated fluorescence that was inhibited by treatment with gallein (Figure 7B).

To determine whether Golgi-GRK2ct can block ET-1–stimulated hypertrophy in AVMs, we infected cells with adenoviruses expressing either PM- or Golgi-targeted GRK2ct, followed by treatment with ET-1 for 24 h. Hypertrophy was assessed by ANF staining. As in the NRVMs, stimulation with ET-1 caused a robust increase in ANF expression that was inhibited by expression of either PM- or Golgi-targeted GRK2ct (Figure 7, C and D).

Because either Golgi or PM GRK2ct can block hypertrophy, G β γ -mediated pathways at the PM or the Golgi are both required for cardiomyocyte hypertrophy. Golgi-targeted GRK2ct would not be




FIGURE 3: $G\beta\gamma$ signaling at the Golgi apparatus stimulates perinuclear PI4P depletion in NRVMs and nuclear PKD activation. (A) NRVMs were transfected with Golgi-GRK2ct. GRK2ct localization was analyzed by immunocytochemistry with a GRK2 antibody with an Alexa Fluor 546–conjugated secondary antibody and imaged in the red channel. The Golgi-specific antibody (anti-Golgi 58K protein) was used to identify the Golgi apparatus, with Alexa Fluor 488 secondary antibody imaged in the green channel. (B) Cells cotransfected with PM-GRK2ct, and myristoylated YFP to mark the PM, were analyzed for YFP fluorescence and GRK2ct localization as in A. (C) Cells were cotransfected with GFP-FAPP-PH and either LacZ or PM-GRK2ct constructs. Perinuclear GFP-FAPP-PH fluorescence was monitored with time after ET-1 addition. (D) Same as C, except that Golgi-targeted GRK2ct was used. (E) Quantitation of data in C and




FIGURE 4: Targeting Gβγ to the Golgi apparatus stimulates PI4P hydrolysis and nuclear PKD activation. (A) Cells were transfected with mCherry-Gβ₁, G_γ2FRB, and CFP-Golgi-FKBP. mCherry fluorescence (top two) and CFP fluorescence (bottom two) before and 10 min after rapamycin addition. (B) NRVMs were transfected with GFP-FAPP-PH, Gβ₁, G_γ2FRB, and either CFP-Golgi-FKBP or PM-FKBP. At time 0, 24 h after transfection, 10 μM rapamycin was added, and perinuclear PI4P fluorescence was monitored with time. (C) Cells were transfected with Gβ₁, G_γ2FRB, and nDKAR and either PM-FKBP or Golgi FKBP. At time 0, 24 h after transfection, 10 μM rapamycin was added, and nDKAR YFP/CFP fluorescence in the nucleus was monitored. All traces represent pooled data from at least three cells from three separate myocyte preparations ± SEM. Scale bars, 10 μm.

expected to influence PM GPCR desensitization mechanisms such as GRK2 recruitment and instead involve inhibition of the hypertrophic Golgi PLCε/PI4P signaling pathway. PM-targeted GRK2ct has the potential block the Golgi PLCε/PI4P pathway by preventing Gβγ translocation from the PM to the Golgi, but the data in Figure 2C indicate that PM-GRK2ct has no effect on Golgi PI4P hydrolysis. This indicates that two independent PM and Golgi Gβγ-mediated pathways are

required for hypertrophy development in cardiac myocytes. (Zhang et al., 2013) and is blocked by a Golgi-targeted Gβγ blocker (Golgi-GRK2ct). 3) Gβγ is a known upstream regulator of PLCε (Wing et al., 2001). 4) PLCε is required for development of cardiac hypertrophy in vivo and in vitro (Zhang et al., 2011, 2013). Although it is formally possible that blockade of Gβγ signaling at the Golgi inhibits hypertrophy development through a parallel pathway that does not involve regulation of

required for hypertrophy development in cardiac myocytes.

DISCUSSION

Expression of GRK2ct in cardiomyocytes to block Gβγ signaling inhibits development of hypertrophy in vivo and in vitro (Koch et al., 1995; Rockman et al., 1998; Casey et al., 2010). Our laboratory has developed small molecule-based inhibitors of Gβγ that block interactions between Gβγ and various downstream protein targets, including GRK2 (Bonacci et al., 2006). Gallein administration prevents development of heart failure in mice in several different models of hypertrophy (Casey et al., 2010). A model for the action of gallein and GRK2ct involves inhibition of GRK2 recruitment by Gβγ downstream of the βAR. On the other hand, GRK2ct and gallein/M119 are general Gβγ inhibitors that block multiple signaling pathways. This raises the possibility that the efficacy of these reagents in prevention of heart failure may rest, at least in part, in their ability to inhibit other signaling pathways downstream of Gβγ. Indeed, pathways involving Gβγ-dependent regulation of ERK phosphorylation (Lorenz et al., 2009) and phosphoinositide 3-kinase γ (PI3Kγ) recruitment (Naga Prasad et al., 2000; Nienaber et al., 2003; Prasad et al., 2003) have potential involvement in regulation of cardiac hypertrophy.

Here we identified another Gβγ pathway that is required for cardiac hypertrophy, which we propose involves Gβγ-dependent activation of PLCε at the perinuclear Golgi, downstream of the Gq- and G13-coupled ET-1A receptor. This proposal is based on the following observations: 1) Two different blockers of hypertrophy in vivo and in vitro—gallein and GRK2ct—block ET-1-dependent activation of Golgi PI4P hydrolysis in cardiac myocytes. 2) ET-1-dependent PI4P depletion at the Golgi and subsequent PKD activation requires PLCε (Zhang et al., 2013) and is blocked by a Golgi-targeted Gβγ blocker (Golgi-GRK2ct). 3) Gβγ is a known upstream regulator of PLCε (Wing et al., 2001). 4) PLCε is required for development of cardiac hypertrophy in vivo and in vitro (Zhang et al., 2011, 2013). Although it is formally possible that blockade of Gβγ signaling at the Golgi inhibits hypertrophy development through a parallel pathway that does not involve regulation of

D. Curves in C and D were fitted with single-exponential decay curves using GraphPad Prism 6 and analyzed for statistical significance. **p* < 0.05 and *****p* < 0.001 relative to ET-1 control. (F) NRVMs were cotransfected with nDKAR and LacZ, PM-GRK2ct, or Golgi-GRK2ct, and the nDKAR YFP/CFP ratio in the nucleus was monitored over time after addition of ET-1. (G) Changes in YFP/CFP ratio, normalized to time 0, ± SEM for each trace in F were pooled from 35 to 40 min, averaged, and analyzed by a one-way analysis of variance (ANOVA). (H) Curves from F were fitted and analyzed as in E. All traces (C, D, F, G) are pooled data from at least three cells from three separate myocyte preparations ± SEM. Scale bars, 10 μm.




FIGURE 5: Blocking $G\beta\gamma$ signaling in the Golgi or PM prevents ET-1–stimulated hypertrophic gene expression in NRVMs. (A) NRVMs were cotransfected with YFP and GRK2ct, followed by treatment with 100 nM ET-1 or vehicle. After 24 h, cells were fixed and stained for ANF (red). With ET-1 treatment, the intensity of red staining increases in untransfected cells but not in cells expressing YFP and GRK2ct. Scale bar, 40 μ m. (B) Cells were transfected with YFP alone or YFP plus GRK2ct, Golgi GRK2ct, or PM-GRK2ct. Cells were treated as in A. Scale bar, 20 μ m. (C) The intensity of ANF staining was quantitated with and without ET-1 only in transfected cells identified by YFP fluorescence. Quantitation is based on combined data (mean \pm SEM) from four separate experiments. **** $p < 0.0001$, statistically different from YFP control. None of the other samples is statistically different from YFP control (one-way ANOVA).

PLC, the simplest model that can account for the data involves activation of PLC ϵ by $G\beta\gamma$ at the Golgi.

We also show that expression of either PM- or Golgi-targeted GRK2ct inhibits development of cardiac hypertrophy but that PM targeting of GRK2ct does not affect perinuclear PI4P hydrolysis. This indicates that there are independent roles for $G\beta\gamma$ at the PM and the Golgi and that requirement for $G\beta\gamma$ at the PM is not to regulate the Golgi PI4P hydrolysis pathway. We showed that the PLC/PI4P hydrolysis pathway at the Golgi is required for ET-1–dependent hypertrophy but not that it is sufficient to drive hypertrophy. We propose that a second parallel signal is dependent on $G\beta\gamma$ activity at the PM and is also required for stimulation of hypertrophy. This could involve GRK2, PLC, or an undefined effector at the PM.

A major open question is how ET-1, through activation of ET-1AR at the PM, leads to $G\beta\gamma$ signaling at the perinuclear Golgi apparatus. One possibility is diffusion-based translocation of $G\beta\gamma$ from the PM to the Golgi. Translocation of fluorescently labeled G protein $\beta\gamma$ subunits from the PM to the Golgi was demonstrated (Saini *et al.*, 2007). This was proposed to rely on dissociation of $G\alpha$ from $G\beta\gamma$,

where modification of both $G\alpha$ and $G\gamma$ with lipids is sufficient to maintain PM localization of a $G\alpha\beta\gamma$ heterotrimer. The model suggests that when these subunits dissociate, the single lipid modification on the γ subunit is not sufficient to retain the $G\beta\gamma$ complex at the PM, and a new equilibrium is established in which $G\beta\gamma$ can equilibrate to internal membranes such as the Golgi (O'Neill *et al.*, 2012). Alternatively, transport of $G\beta\gamma$ in endosomes with or without internalization of cell surface ET-1A receptors could result in accumulation of free $G\beta\gamma$ in the Golgi. Finally, it is possible that activation of ET1A signaling at the PM could generate a signal at the PM that could diffuse to the Golgi to activate heterotrimers resident at the Golgi. A possible candidate for this might be an activator of G protein signaling (AGS) protein that can stimulate $G\beta\gamma$ release from $G\alpha$ subunits in a receptor-independent manner (Blumer *et al.*, 2007). It was recently that AGS3 can translocate from the PM to the Golgi in response to receptor activation (Oner *et al.*, 2013). Defining which, if any, of these mechanisms mediates this signal requires further investigation.

These observations have clinical significance with respect to design of inhibitors to inhibit development of heart failure. Recent




FIGURE 6: Blocking G $\beta\gamma$ signaling in the Golgi or PM prevents ET-1-stimulated cardiomyocyte hypertrophic cell growth. (A) NRVMs were infected with adenoviruses expressing YFP or YFP and the indicated targeted GRK2ct constructs. Cells were treated with 100 nM ET-1 or vehicle. After 48 h, cells were visualized in the YFP channel. Scale bar, 40 μ m. (B) Cells treated as in A quantitated for cell area using ImageJ (National Institutes of Health, Bethesda, MD). Quantitation is based on combined data (mean \pm SEM) from three separate experiments; 50 cells each experiment. Data were analyzed by one-way ANOVA; *** $p < 0.005$.

efforts have focused on development of inhibitors of GRK2 that improve cardiac performance and may prevent development of heart failure (Thal *et al.*, 2011, 2012). The data presented here suggest that although directly targeting GRK2 may be effective, a strategy that targets G $\beta\gamma$ may have greater efficacy due to its ability to block multiple hypertrophic signaling pathways. The list of G $\beta\gamma$ -dependent pathways that regulate cardiac hypertrophy continues to grow and include PI3K γ (Prasad *et al.*, 2003; Perrino *et al.*, 2006) and ERK activation (Lorenz *et al.*, 2009; Ruppert *et al.*, 2013), to which we now add G $\beta\gamma$ -regulated PLC activity at the Golgi. We also observed that G $\beta\gamma$ inhibition at the PM inhibits hypertrophy in NRVMs but have not characterized the underlying signaling mechanism. As new

therapeutic approaches to heart failure are considered, a global G $\beta\gamma$ inhibition strategy might have high efficacy due to multiple target inhibition, but there may be advantages to targeting G $\beta\gamma$ signaling at specific cellular locations such as the Golgi apparatus that could avoid potential side effects of global G $\beta\gamma$ inhibition or even G $\beta\gamma$ signaling at the PM. This remains to be explored in more-physiological animal models of heart failure.

MATERIALS AND METHODS

Materials

ET-1 was purchased from Sigma-Aldrich (St. Louis, MO). Gallein was from TCI America (Portland, OR), dissolved in phosphate-buffered saline (PBS), and titrated to pH 7.8 with NaOH to create a 15 mg/ml stock that was stored in aliquots at -20°C . Rapamycin and cpTOME were from EMD Millipore (Billerica, MA).

Adenoviral constructs and plasmids

The EGFP-FAPP-PH domain (N-terminal EGFP fusion) for detection of PI4P was as previously described (Balla *et al.*, 2005, 2009). GFP-Tubby (Santagata *et al.*, 2001; Balla *et al.*, 2009) and GFP-PLC δ -PH (Balla and Varnai, 2002) for detection of PIP $_2$ were also previously described. nDKAR (Kunkel *et al.*, 2007) was provided by Alexandra Newton (University of California, San Diego, CA). nDKAR was inserted into an adenovirus vector by standard methods. G γ_2 -C68S-FRB, PM-FKBP (LF2C, Lyn1-11-FKBP(X2)-CFP), and Golgi-FKBP (KDELr-D193N-FKBP(x2)-CFP), Golgi-GRK2ct (KDELrD193N-GRK2(495-689)), and PM-GRK2ct (GRK2(495-689)-Rit) were constructed as previously described (Irannejad and Wedegaertner, 2010). Adenoviruses were constructed using a vector expressing YFP from one promoter and PM-GRK2ct, Golgi-GRK2ct, or GFP-Tubby from a second cytomegalovirus promoter.

Antisera

Anti-GRK2 was from Santa Cruz Biotechnology (Dallas, TX; 1:100); rabbit polyclonal affinity-purified anti-ANF was from EMD Millipore. Golgi protein, 58K protein (1:250), monoclonal antibody (58K-9) was from Thermo Fisher Scientific (Waltham, MA). mAKAP antibody was kindly provided by Michael Kapiloff (University of Miami School of Medicine, Miami, FL).

nDKAR FRET

NRVMs were transduced with an nDKAR-expressing adenovirus. After 48 h, FRET was analyzed by confocal microscopy on an Olympus FV1000MP microscope. Transduced cells were identified by exciting CFP at 440 nm and emission monitored at 480 nm. FRET was determined by measuring the ratio of YFP emission at 535 nm to CFP emission at 480 nm after CFP excitation at 440 nm. Because the YFP and CFP are fused in this construct, it is not necessary to correct for expression or spectral bleedthrough of the various constructs. Data were normalized to the initial YFP/CFP ratio in each set at an arbitrary value of 1.

Isolation of cardiac myocytes

NRVMs were isolated from 2- to 3-d-old Sprague Dawley rats and cultured essentially as described (Zhang *et al.*, 2011). Adult myocytes were isolated from 4- to 6-mo-old wild-type mice as previously described (Oestreich *et al.*, 2009).

Transfection and transduction of NRVMs and AVMs

NRVMs cultured for 2 d were transfected with 1–2 μ g of the indicated plasmid constructs or transduced with 50 multiplicity of infection (MOI) of the indicated viruses. Adenoviral transduction




FIGURE 7: Blocking G $\beta\gamma$ signaling in the Golgi or PM prevents ET-1–stimulated ANF expression in AVMs. (A) AVMs were infected with adenoviruses expressing either Tubby-GFP (top) or GFP-FAPP-PH (bottom). (B) AVMs were infected with GFP-FAPP-PH and stimulated with 100 nM ET-1, and perinuclear GFP-FAPP-PH was monitored over time as indicated. Each trace represents pooled data from at least five cells from five separate myocyte preparations \pm SEM. (C) AVMs were infected with viruses (50 MOI) expressing YFP, Golgi-GRK2ct, or PM-GRK2ct (only cells expressing YFP or Golgi-GRK2ct are shown). Viruses expressing Golgi-GRK2ct and PM-GRK2ct also expressed YFP from a separate cassette. Cells were stimulated for 24 h with 100 nM ET-1 and fixed and stained for ANF expression. (D) ANF expression was quantitated as mean \pm SEM for all treatments as described in *Materials and Methods*. **** $p < 0.0001$, statistically different from YFP control. None of the other samples is statistically different from YFP control (one-way ANOVA). Scale bars, 10 μ m.

efficiency was >90%, and plasmid-based transfection efficiency was ~5%. Cells were examined 24 h later by microscopy or as indicated for measurement of nDKAR FRET or GFP-FAPP-PH fluorescence.

Confocal microscopy

Myocytes transfected with plasmids or transduced adenoviruses expressing enhanced GFP (EGFP)-, YFP-, or mCherry-tagged proteins were analyzed in an Olympus FV1000MP microscope in confocal mode with a LUMPLFL 40x/0.8 numerical aperture W (Olympus, Tokyo, Japan) lens. EGFP was excited at 488 nm and emission monitored at 510 nm, YFP was excited at 515 nm and emission monitored at 527 nm, mCherry and monomeric red fluorescent protein were excited at 559 nm and emission monitored at 618 nm, CFP was excited at 440 nm and emission monitored at 476 nm, and 4',6'-diamidino-2-phenylindole was excited at 405 nm and emission monitored at 461 nm. During imaging, cells were in culture medium containing serum.

Western blotting

Membranes were blocked in 5% milk in Tris-buffered saline/0.1% Tween 20 (TBST) for 2 h at room temperature. After three washes (5 min each) with TBST, the membranes were incubated with anti-GRK2 antibody (1:3000) in 3% bovine serum albumin (BSA) in TBST overnight at 4°C. After incubation, membranes were washed four times with TBST (10 min each) and incubated with anti-rabbit secondary antibody (1:10,000, IR-800; LI-COR Biosciences, Lincoln, NE) in 5% milk in TBST for 1 h at room temperature. Blocking was for 1 h at room temperature, and primary antibody was incubated for 2 h at room temperature and secondary antibody for 1 h at room temperature. The membranes were washed three times with TBST (10 min each) and scanned on a LI-COR Odyssey infrared imaging system.

Immunocytochemistry for ANF induction

NRVMs were transfected with 1 µg of empty vector, GRK2ct, PM-GRK2ct, or Golgi-GRK2ct and 1 µg of a YFP-expressing plasmid using Lipofectamine 2000 to allow identification of transfected cells. Sixteen hours after transfection 100 nM ET-1 was added for 24 h to induce hypertrophy. Cells were briefly washed with PBS and fixed for 25 min in a solution of 2% paraformaldehyde (Sigma-Aldrich) with 0.025% glutaraldehyde (J.T. Baker, Center Valley, PA) in PBS. After permeabilization in 0.2% Triton X-100 for 15 min at room temperature, cells were blocked in a solution of 10% BSA in PBS for 1 h at room temperature. Primary antibody (rabbit anti-ANF [EMD Millipore]) was incubated at a dilution of 1:2500 in PBS overnight at 4°C. After three washes in PBS, cells were incubated with the secondary antibody (goat anti-rabbit Alexa Fluor 546 [Life Technologies, Grand Island, NY]) at a 1:2500 dilution in PBS for 3 h at room temperature. Fluorescence images were captured by confocal microscopy and analyzed for ANF expression.

For AVMs, freshly isolated adult myocytes from mice were infected with adenoviruses expressing YFP, PM-GRK2ct, or Golgi-GRK2ct for 2 h, after which the virus was removed by washing and the contraction inhibitor blebbistatin was added to the culture medium. At 16 h later, cells were treated with either vehicle or 100 nM ET-1 for 24 h. Cells were fixed and stained for ANF as described.

To quantitate ANF induction, fluorescence images were captured by confocal microscopy, and transfected YFP fluorescent cells were identified and analyzed for ANF fluorescence intensity using Olympus FluoView software, version 2.0a. Perinuclear area was traced, and peak ANF intensity was measured for each cell minus background subtraction from area in the cytosol. Data were

averaged based on individual cells (>40 cells total) pooled from three separate experiments for NRVMs and AVMs.

ACKNOWLEDGMENTS

This research was supported by National Institutes of Health Grants R01GM53536 and R01GM81772 (A.V.S.) and R01 GM56444 (P.B.W.).

REFERENCES

- Balla T, Szentepey Z, Kim YJ (2009). Phosphoinositide signaling: new tools and insights. *Physiology* 24, 231–244.
- Balla A, Tuymetova G, Tsiomenko A, Várnai P, Balla T (2005). A plasma membrane pool of phosphatidylinositol 4-phosphate is generated by phosphatidylinositol 4-kinase type-iii α : studies with the PH domains of the oxysterol binding protein and FAPP1. *Mol Biol Cell* 16, 1282–1295.
- Balla T, Várnai P (2002). Visualizing cellular phosphoinositide pools with GFP-fused protein-modules. *Sci STKE* 2002, pl3.
- Bkaily G, Avedanian L, Al-Khoury J, Provost C, Nader M, D'Orléans-Juste P, Jacques D (2011). Nuclear membrane receptors for ET-1 in cardiovascular function. *Am J Physiol Regul Integr Comp Physiol* 300, R251–R263.
- Blumer JB, Smrcka AV, Lanier SM (2007). Mechanistic pathways and biological roles for receptor-independent activators of G-protein signaling. *Pharmacol Ther* 113, 488–506.
- Bonacci TM, Mathews JL, Yuan C, Lehmann DM, Malik S, Wu D, Font JL, Bidlack JM, Smrcka AV (2006). Differential targeting of G $\beta\gamma$ -subunit signaling with small molecules. *Science* 312, 443–446.
- Bos JL (2003). Epac: a new cAMP target and new avenues in cAMP research. *Nat Rev Mol Cell Biol* 4, 733–738.
- Campelo F, Malhotra V (2012). Membrane fission: the biogenesis of transport carriers. *Annu Rev Biochem* 81, 407–427.
- Casey LM, Pistner AR, Belmonte SL, Migdalovich D, Stolpnik O, Nwakanma FE, Vorobiof G, Dunaevsky O, Matavel A, Lopes CM, et al. (2010). Small molecule disruption of G $\beta\gamma$ signaling inhibits the progression of heart failure. *Circ Res* 107, 532–539.
- Chisari M, Saini DK, Kalyanaraman V, Gautam N (2007). Shuttling of G protein subunits between the plasma membrane and intracellular membranes. *J Biol Chem* 282, 24092–24098.
- D'Angelo DD, Sakata Y, Lorenz JN, Boivin GP, Walsh RA, Liggett SB, Dorn GW II (1997). Transgenic α_q overexpression induces cardiac contractile failure in mice. *Proc Natl Acad Sci USA* 94, 8121–8126.
- Diaz Anel AM (2007). Phospholipase C $\beta 3$ is a key component in the G $\beta\gamma$ /PKC η /PKD-mediated regulation of trans-Golgi network to plasma membrane transport. *Biochem J* 406, 157–165.
- Diaz Anel AM, Malhotra V (2005). PKC η is required for $\beta_1 \gamma_2 / \beta_3 \gamma_2$ - and PKD-mediated transport to the cell surface and the organization of the Golgi apparatus. *J Cell Biol* 169, 83–91.
- Fielitz J, Kim M-S, Shelton JM, Qi X, Hill JA, Richardson JA, Bassel-Duby R, Olson EN (2008). Requirement of protein kinase D1 for pathological cardiac remodeling. *Proc Natl Acad Sci USA* 105, 3059–3063.
- Filtz TM, Grubb DR, McLeod-Dryden TJ, Luo J, Woodcock EA (2009). Gq-initiated cardiomyocyte hypertrophy is mediated by phospholipase C $\beta 1b$. *FASEB J* 23, 3564–3570.
- Irannejad R, Wedegaertner PB (2010). Regulation of constitutive cargo transport from the trans-Golgi network to plasma membrane by Golgi-localized G protein $\beta\gamma$ subunits. *J Biol Chem* 285, 32393–32404.
- Jamora C, Yamanoue N, Van Lint J, Laudenslager J, Vandenheede JR, Faulkner DJ, Malhotra V (1999). G $\beta\gamma$ -mediated regulation of Golgi organization is through the direct activation of protein kinase D. *Cell* 98, 59–68.
- Knowlton KU, Michel MC, Itani M, Shubeita HE, Ishihara K, Brown JH, Chien KR (1993). The $\alpha 1A$ -adrenergic receptor subtype mediates biochemical, molecular, and morphologic features of cultured myocardial cell hypertrophy. *J Biol Chem* 268, 15374–15380.
- Koch WJ, Rockman HA, Samama P, Hamilton RA, Bond RA, Milano CA, Lefkowitz RJ (1995). Cardiac function in mice overexpressing the β -adrenergic receptor kinase or a β ARK inhibitor. *Science* 268, 1350–1353.
- Kunkel MT, Toker A, Tsien RY, Newton AC (2007). Calcium-dependent regulation of protein kinase D revealed by a genetically encoded kinase activity reporter. *J Biol Chem* 282, 6733–6742.
- Lorenz K, Schmitt JP, Schmitteckert EM, Lohse MJ (2009). A new type of ERK1/2 autophosphorylation causes cardiac hypertrophy. *Nat Med* 15, 75–83.
- Naga Prasad SV, Esposito G, Mao L, Koch WJ, Rockman HA (2000). G $\beta\gamma$ -dependent phosphoinositide 3-kinase activation in hearts with in vivo pressure overload hypertrophy. *J Biol Chem* 275, 4693–4698.

- Nienaber JJ, Tachibana H, Naga Prasad SV, Esposito G, Wu D, Mao L, Rockman HA (2003). Inhibition of receptor-localized PI3K preserves cardiac β -adrenergic receptor function and ameliorates pressure overload heart failure. *J Clin Invest* 112, 1067–1079.
- Oestreich EA, Malik S, Goonasekera SA, Blaxall BC, Kelley GG, Dirksen RT, Smrcka AV (2009). Epac and phospholipase C ϵ regulate Ca²⁺ release in the heart by activation of protein kinase C δ and calcium-calmodulin kinase II. *J Biol Chem* 284, 1514–1522.
- Oestreich EA, Wang H, Malik S, Kaproth-Joslin KA, Blaxall BC, Kelley GG, Dirksen RT, Smrcka AV (2007). Epac-mediated activation of phospholipase C ϵ plays a critical role in β -adrenergic receptor-dependent enhancement of Ca²⁺ mobilization in cardiac myocytes. *J Biol Chem* 282, 5488–5495.
- O'Neill PR, Karunaratne WKA, Kalyanaraman V, Silvius JR, Gautam N (2012). G-protein signaling leverages subunit-dependent membrane affinity to differentially control $\beta\gamma$ translocation to intracellular membranes. *Proc Natl Acad Sci USA* 109, E3568–E3577.
- Oner SS, Vural A, Lanier SM (2013). Translocation of activator of G-protein signaling 3 to the Golgi apparatus in response to receptor activation and its effect on the trans-Golgi network. *J Biol Chem* 288, 24091–24103.
- Perrino C, Rockman HA, Chiariello M (2006). Targeted inhibition of phosphoinositide 3-kinase activity as a novel strategy to normalize β -adrenergic receptor function in heart failure. *Vasc Pharmacol* 45, 77–85.
- Prasad SVN., Perrino C, Rockman HA (2003). Role of phosphoinositide 3-kinase in cardiac function and heart failure. *Trends Cardiovasc Med* 13, 206–212.
- Rengo G, Lympelopoulous A, Leosco D, Koch WJ (2011). GRK2 as a novel gene therapy target in heart failure. *J Mol Cell Cardiol* 50, 785–792.
- Rockman HA, Chien KR, Choi DJ, Iaccarino G, Hunter JJ, Ross J Jr, Lefkowitz RJ, Koch WJ (1998). Expression of a β -adrenergic receptor kinase 1 inhibitor prevents the development of myocardial failure in gene-targeted mice. *Proc Natl Acad Sci USA* 95, 7000–7005.
- Ruppert C, Deiss K, Herrmann S, Vidal M, Oezkur M, Gorski A, Weidemann F, Lohse MJ, Lorenz K (2013). Interference with ERKThr188 phosphorylation impairs pathological but not physiological cardiac hypertrophy. *Proc Natl Acad Sci USA* 110, 7440–7445.
- Saini DK, Kalyanaraman V, Chisari M, Gautam N (2007). A family of G protein $\beta\gamma$ subunits translocate reversibly from the plasma membrane to endomembranes on receptor activation. *J Biol Chem* 282, 24099–24108.
- Saini DK, Karunaratne WK, Angaswamy N, Saini D, Cho JH, Kalyanaraman V, Gautam N (2010). Regulation of Golgi structure and secretion by receptor-induced G protein $\beta\gamma$ complex translocation. *Proc Natl Acad Sci USA* 107, 11417–11422.
- Santagata S, Boggon TJ, Baird CL, Gomez CA, Zhao J, Shan WS, Myszkowski DG, Shapiro L (2001). G-protein signaling through tubby proteins. *Science* 292, 2041–2050.
- Smrcka AV, Brown JH, Holz GG (2012). Role of phospholipase C ϵ in physiological phosphoinositide signaling networks. *Cell Signal* 29, 1333–1343.
- Smrcka AV, Hepler JR, Brown KO, Sternweis PC (1991). Regulation of polyphosphoinositide-specific phospholipase C activity by purified Gq. *Science* 251, 804–807.
- Taylor SJ, Chae HZ, Rhee SG, Exton JH (1991). Activation of the β 1 isozyme of phospholipase C by α subunits of the G q class of G proteins. *Nature* 350, 516–518.
- Teruel MN, Blanpied TA, Shen K, Augustine GJ, Meyer T (1999). A versatile microportation technique for the transfection of cultured CNS neurons. *J Neurosci Methods* 93, 37–48.
- Thal DM, Homan KT, Chen J, Wu EK, Hinkle PM, Huang ZM, Chuprun JK, Song J, Gao E, Cheung JY, et al. (2012). Paroxetine is a direct inhibitor of G protein-coupled receptor kinase 2 and increases myocardial contractility. *ACS Chem Biol* 7, 1830–1839.
- Thal DM, Yeow RY, Schoenau C, Huber J, Tesmer JJ (2011). Molecular mechanism of selectivity among G protein-coupled receptor kinase 2 inhibitors. *Mol Pharmacol* 80, 294–303.
- Vega RB, Harrison BC, Meadows E, Roberts CR, Papst PJ, Olson EN, McKinsey TA (2004). Protein kinases C and D mediate agonist-dependent cardiac hypertrophy through nuclear export of histone deacetylase 5. *Mol Cell Biol* 24, 8374–8385.
- Wang H, Oestreich EA, Maekawa N, Bullard TA, Vikstrom KL, Dirksen RT, Kelley GG, Blaxall BC, Smrcka AV (2005). Phospholipase C ϵ modulates β -adrenergic receptor-dependent cardiac contraction and inhibits cardiac hypertrophy. *Circ Res* 97, 1305–1313.
- Wing MR, Houston D, Kelley GG, Der CJ, Siderovski DP, Harden TK (2001). Activation of phospholipase C- ϵ by heterotrimeric G protein $\beta\gamma$ -subunits. *J Biol Chem* 276, 48257–48261.
- Wu X, Zhang T, Bossuyt J, Li X, McKinsey TA, Dedman JR, Olson EN, Chen J, Brown JH, Bers DM (2006). Local InsP₃-dependent perinuclear Ca²⁺ signaling in cardiac myocyte excitation-transcription coupling. *J Clin Invest* 116, 675–682.
- Zhang L, Malik S, Kelley GG, Kapiloff MS, Smrcka AV (2011). Phospholipase C ϵ scaffolds to muscle-specific A kinase anchoring protein (mAKAP β) and integrates multiple hypertrophic stimuli in cardiac myocytes. *J Biol Chem* 286, 23012–23021.
- Zhang L, Malik S, Pang J, Wang H, Park K, Yule D, Blaxall B, Smrcka AV (2013). Phospholipase C ϵ hydrolyzes perinuclear phosphatidylinositol 4-phosphate to regulate cardiac hypertrophy. *Cell* 153, 216–227.