

3-1-2016

Predicting Revision Following In Situ Ulnar Nerve Decompression for Patients With Idiopathic Cubital Tunnel Syndrome.

Michael P. Gaspar

The Philadelphia Hand Center; Thomas Jefferson University

Patrick M. Kane

Thomas Jefferson University; Philadelphia Hand Center

Dechporn Putthiwara

Philadelphia Hand Center

Sidney M. Jacoby

Thomas Jefferson University; Philadelphia Hand Center

A. Lee Osterman

Thomas Jefferson University; Philadelphia Hand Center

Follow this and additional works at: <https://jdc.jefferson.edu/orthofp>



Part of the [Orthopedics Commons](#), and the [Surgery Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Gaspar, Michael P.; Kane, Patrick M.; Putthiwara, Dechporn; Jacoby, Sidney M.; and Osterman, A. Lee, "Predicting Revision Following In Situ Ulnar Nerve Decompression for Patients With Idiopathic Cubital Tunnel Syndrome." (2016). *Department of Orthopaedic Surgery Faculty Papers*. Paper 90.
<https://jdc.jefferson.edu/orthofp/90>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Orthopaedic Surgery Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

1 **ABSTRACT**

2 PURPOSE

3 To determine the incidence of revision and potential risk factors for needing
4 revision surgery following in situ ulnar nerve decompression for patients with
5 idiopathic cubital tunnel syndrome (CuTS).

6

7 METHODS

8 We conducted a retrospective chart review of all patients treated at one specialty
9 hand center with an open in situ ulnar nerve decompression for idiopathic CuTS
10 from January 2006 through December 2010. Revision incidence was determined by
11 identifying patients who underwent additional surgeries for recurrent or persistent
12 ulnar nerve symptoms. Bivariate analysis was performed to determine which
13 variables had a significant influence on the need for revision surgery.

14

15 RESULTS

16 Revision surgery was required in 3.2% (7 of 216) of all cases. Age of less than 50
17 years at the time of index decompression was the lone significant predictor of need
18 for revision surgery. Other patient factors, including sex, diabetes, smoking history,
19 and worker compensation status were not predictive of the need for revision
20 surgery. Disease-specific variables including nerve conduction velocities, McGowan
21 grading, and predominant symptom type were also not predictive of revision.

22

23 CONCLUSIONS

24 For patients with idiopathic cubital tunnel syndrome, the risk of revision surgery
25 following in situ ulnar nerve decompression is low. However, this risk was
26 increased in patients who were younger than 50 years of age at the time of the index
27 procedure. The findings of this study suggest that, in the absence of underlying
28 elbow arthritis or prior elbow trauma, in situ ulnar nerve decompression is an
29 effective, minimal-risk option for the initial surgical treatment of CuTS.

30

31 LEVEL OF EVIDENCE

32 Prognostic Level III

33

34 INTRODUCTION

35 Cubital tunnel syndrome (CuTS) is second only to carpal tunnel syndrome in
36 incidence among compression neuropathies of the upper extremity. [1-3] Despite its
37 commonality, there is no established consensus regarding the optimal surgical
38 treatment. This is evidenced by a wide range of surgical options including in situ
39 decompression, medial epicondylectomy and subcutaneous, intramuscular or
40 submuscular transposition of the ulnar nerve. Additionally, in recent years surgeons
41 have also advocated for endoscopic or minimal-incision release of the ulnar nerve,
42 with or without transposition, to further minimize soft tissue trauma and potential
43 vascular insult to the nerve, while allowing for faster recovery, thus further
44 expanding the number of treatment options. [4-6]

45

46 Technique selection can depend on a variety of factors including surgeon
47 preference, patient anatomy, patient desires, underlying pathology, and
48 complication rates. Transposition, for example, often requires extensive dissection
49 around the nerve, which may compromise its extrinsic vascular supply. Thus, it may
50 be contraindicated in patients with diabetes for instance who may have a tenuous
51 vascular system at the level of the cubital tunnel. [7, 8] In addition, with an
52 increasing focus on healthcare economics in the United States, the relative cost-
53 effectiveness of different treatment options for CuTS may progressively factor into
54 surgical decision-making, thus potentially clouding the treatment decision even
55 further. [9-11]

56

57 Generally, in situ decompression offers the least invasive surgical option but may
58 increase the risk of revision surgery. [12, 13] A recent study found that prior history
59 of trauma around the elbow was a notable predictor of need for revision after in situ
60 decompression of the ulnar nerve, while other postulated factors including patient
61 age had no effect. [14] However, risk factors for revision in patients with *idiopathic*
62 CuTS, that is, those without an underlying traumatic, arthritic, or other pre-
63 disposing etiology, remain unclear. As revision surgery yields inferior outcomes
64 versus primary surgery for CuTS, information on risk factors leading to revision in
65 these patients with idiopathic CuTS could provide a valuable addition to the overall
66 treatment algorithm. [15]

67 The purpose of this study was to determine the incidence of needed revision after in
68 situ ulnar nerve decompression for patients with idiopathic CuTS and to investigate
69 which patient risk factor(s) may contribute to an increased likelihood of needing
70 revision.

71

72 MATERIALS and METHODS

73 This study was approved by our institutional review board. Using our departmental
74 electronic billing database search for Current Procedural Terminology (American
75 Medical Association, Chicago, IL, USA) code 64718 (surgery on ulnar nerve at
76 elbow), we identified all patients who had undergone in situ ulnar nerve
77 decompression surgery from January 2006 through December 2010. Patients who

78 demonstrated intraoperative subluxation of the ulnar nerve following in situ
79 decompression were excluded, as these patients subsequently underwent either
80 anterior transposition of the ulnar nerve or medial epicondylectomy. Patients were
81 also excluded if they underwent in situ ulnar nerve decompression for reasons other
82 than treatment of CuTS symptoms (e.g., prophylactic release performed in
83 conjunction with elbow arthroplasty or fracture fixation) or had previously
84 undergone operative treatment for CuTS. In addition, patients with a prior history of
85 fracture or trauma at the elbow were excluded, as were those with a history of
86 degenerative, post-traumatic, or inflammatory arthritis at the elbow. However,
87 patients with a known history of inflammatory or systemic arthritis without
88 evidence of local arthritic changes at the surgical elbow were not excluded. Finally,
89 patients with less than 6 months of follow-up at our institution were excluded from
90 data analysis unless a revision surgery occurred in that time interval. Records for
91 those patients with less than 6 months of follow-up were reviewed in an effort to
92 predict their clinical course. In addition, attempts were made to contact those
93 patients via telephone with the goal of identifying any patients that may have had
94 additional surgery performed elsewhere.

95

96 **Diagnostic workup**

97 Patients seen at our institution are generally evaluated by the treating surgeon prior
98 to obtaining additional studies, including imaging or electrodiagnostic testing.
99 Exceptions to this practice typically only occur in patients who are seen at our
100 institution for a second opinion and have already undergone electrodiagnostic

101 testing prior to presentation. During initial evaluation, a comprehensive clinical
102 examination, including disease-specific tests and provocative maneuvers, is
103 performed. This includes 2 point-discrimination, vibratory discrimination testing,
104 comparative grip strength testing, cross-finger testing, Froment sign, Tinel sign,
105 elbow flexion-compression test, and testing for nerve mobility. When a patient is
106 suspected of having CuTS based on clinical history and physical examination,
107 standard elbow radiographs are routinely obtained to rule out contributory bony
108 abnormalities or deformities in addition to electrodiagnostic testing. Nerve
109 conduction tests are considered abnormal if conduction velocity across the affected
110 elbow is less than 50 meters per second or is decreased by more than 10 meters per
111 second across the elbow . The diagnosis of CuTS is based on clinical findings in
112 conjunction with nerve testing results.

113

114 Additionally, effort is made to elucidate any nerve symptoms not originating at the
115 elbow, such as proximally based cervical pathology or distal compression of the
116 ulnar and median nerves at the wrist. When the diagnostic workup suggests
117 pathology at those distal sites, it is not uncommon in our practice to perform
118 concomitant release of the ulnar and median nerves at the Guyon canal and the
119 carpal tunnel. However, for those patients with findings of ipsilateral cervical
120 radiculopathy, the cervical pathology is generally addressed prior to any operative
121 management of CuTS-related symptoms.

122

123 **Operative indications**

124 Indications for primary in situ decompression generally involve nerve symptoms
125 consistent with CuTS that have failed a trial of conservative management, have
126 positive electrodiagnostic findings, and have a stable ulnar nerve. At our institution,
127 ulnar nerve hypermobility, manifested as nerve subluxation or dislocation during
128 preoperative or intraoperative assessment, is considered a contraindication to
129 performing in situ decompression alone. Thus, when such hypermobility is noted,
130 alternative surgical options such as anterior ulnar nerve transposition or medial
131 epicondylectomy are considered.

132

133 The decision to operate on patients with CuTS in the revision setting is a joint-
134 agreement between the patient and surgeon. Although this is normally approached
135 on a case-by-case basis, the typical scenario involves persistent or incomplete-
136 resolution of symptoms compared to preoperatively. Workup for recurrent or
137 persistent CuTS is largely the same as in primary CuTS described above.

138

139 **Surgical technique and postoperative protocol**

140 All surgeries were performed by one of 8, fellowship-trained orthopedic hand
141 surgeons. A posteromedial incision measuring 5 to 10 centimeters centered about
142 the epicondylar groove is used for exposure. As the incision is carried
143 subcutaneously, care is taken to identify and protect branches of the medial
144 antebrachial cutaneous nerve. Upon identification of the ulnar nerve,
145 decompression is performed via surgical release of the Osborne ligament and fascia
146 overlying the flexor carpi ulnaris with blunt dissection carried roughly 8

147 centimeters proximally to the level of the arcade of Struther. In those patients found
148 to have an anconeus epitrochlearis, the anomalous muscle is generally split or
149 excised depending on its involvement in compression of the ulnar nerve. Care is
150 taken to avoid circumferential dissection around the nerve to preserve its vascular
151 supply. Following release, the elbow is taken through its full range-of-motion to
152 confirm stability of the ulnar nerve. Postoperatively, the limb is placed in a well-
153 padded posterior long-arm orthosis with the elbow positioned in approximately 70
154 degrees of flexion. Active range-of-motion is typically initiated subsequent to the
155 first postoperative visit one week following surgery. Nerve conduction testing is not
156 routinely performed postoperatively except in cases of persistent, recurrent, or
157 worsening symptoms.

158

159 **Data collection and statistical analysis**

160 For those patients satisfying inclusion in the study, demographic, medical, and
161 surgical data were obtained from departmental records. We defined our primary
162 outcome of interest to be revision cubital tunnel surgery performed after in situ
163 ulnar nerve decompression. Thus any patients, who at the time of data analysis had
164 not had revision surgery, were designated to the control cohort. Bivariate analysis
165 was performed for categorical variables of sex, diabetes history, smoking history,
166 presence of bilateral symptoms, predominant preoperative symptom, modified pre-
167 and postoperative modified McGowan grade, concomitant surgery, and worker
168 compensation status using Chi-square or Fisher exact testing. Continuous variables
169 recorded preoperatively including symptom duration, body mass index (BMI), and

170 nerve conduction velocity (NCV), were compared using Student t-test or Mann-
171 Whitney *U* test. Age was analyzed as both a categorical variable (less than 50 years
172 versus greater-than-or-equal-to 50 years) and as a continuous variable.

173

174 RESULTS

175 A total of 216 elbows in 201 patients satisfied inclusion in this study. (See Figure 1)
176 The mean age at the time of surgery for all 216 cases was 53 +/- 14 years, with
177 mean follow-up duration of 22 +/- 21 months. Continuous and categorical
178 demographic variables of the entire study cohort are represented in Tables 1 and 2,
179 respectively.

180

181 Revision surgery was required in 7 (3.2%) cases, with the first revision occurring at
182 a median interval of 10 months from the index surgery (range 3 to 59 months). Five
183 of those patients were revised with anterior subcutaneous transposition, one with
184 submuscular transposition, and one with intramuscular transposition. Two patients
185 required more than one revision for persistent or recurrence of symptoms.

186 Treatment course and demographic characteristics of those patients requiring
187 revision surgery are outlined in Table 3.

188

189 **Bivariate analysis**

190 Younger age had a statistically significant effect on need for revision surgery when
191 analyzed as a continuous variable, mean age non-revised = 53 +/- 14 years versus
192 revised = 43 +/- 7 years; $P = 0.009$, (see Table 1) and as a categorical variable (age \geq

193 50 years vs. age < 50 years; Fisher exact test, $P = 0.002$, see [Table 2](#)). The duration of
194 preoperative symptoms in the revised cohort was roughly double that of the
195 controls, although this association only approached statistical significance ($12 \pm$
196 11 months versus 26 ± 17 months; $P = 0.08$, [Table 1](#)). Patient sex, diabetes history,
197 smoking history, predominant symptom at this time of surgery, modified McGowan
198 grade, concomitant surgery, worker compensation status, body mass index, and
199 ulnar nerve conduction velocity values were not statistically different between
200 those patients requiring revision and those who did not. [Figure 2](#) illustrates the
201 change in modified McGowan grade for the entire study cohort. [Tables 1 and 2](#) detail
202 the respective relationships of continuous and categorical variables and the need for
203 revision surgery.

204

205 **Subjective and validated outcomes**

206 No patients reported worsening of their symptoms following ulnar nerve in situ
207 decompression compared to preoperatively. Of the 209 patients who did not
208 undergo revision surgery, 3 patients complained of persistent sensory symptoms
209 and were offered revision surgery, but they declined. A fourth patient reported
210 recurrence of her symptoms and expressed desire to undergo revision surgery, but
211 she was subsequently lost to follow-up. Multiple attempts to contact that patient via
212 telephone were unsuccessful. The remaining 205 patients reported subjective
213 improvement and general satisfaction following their operation. [Table 3](#) details the
214 treatment course of the revision cohort.

215

216

217 DISCUSSION

218 Selecting the optimal surgical treatment plan for patients with idiopathic cubital
219 tunnel syndrome remains a difficult task. Though numerous studies have explored
220 differences in outcomes among the various surgical options, results have often been
221 inconclusive, and at times, contradictory. [3, 16-18]

222

223 Need for revision surgery is a particularly important outcome to investigate, as it
224 not only represents a sub-optimal clinical result but has important economic
225 considerations as well. With a lack of high quality, adequately powered prospective
226 randomized-control trials comparing the multitude of surgical options for CuTS,
227 cost-effectiveness and decision analyses may afford clinicians a useful tool for
228 comparisons when real-world studies fall short or may simply be impractical. [9, 11,
229 19] A decision analysis study concluded that in situ decompression of the ulnar
230 nerve had the highest utility of 4 tested surgical procedures, while medial
231 epicondylectomy fared worst. [10] These results were later supported by Song et al,
232 who explored the same four surgical treatments for CuTS and found that in situ
233 decompression to be superior to the other options in cost-effectiveness. [11] Both
234 studies used literature available at the time to account for expected incidences of
235 complications and revision for each of the 4 surgical treatments examined. In a
236 randomized-control trial comparing ulnar nerve in situ decompression with
237 anterior subcutaneous transposition, Bartels et al found in situ decompression to be

238 superior from a cost perspective, while also demonstrating a lower incidence of
239 complications. [3, 9]

240

241 Despite the findings of these studies supporting in situ decompression as a first-
242 option for CuTS, the question remains as to which patients are best suited for this
243 versus other surgical options for CuTS, particularly in regards to circumventing the
244 need for revision surgery. Determining which patients are most likely to need
245 revision surgery after initial decompression could be equally as valuable as the
246 previously mentioned cost and decision-based analyses in avoiding the medical and
247 economic costs associated with a second surgery. Krogue and colleagues studied
248 factors leading to revision after in situ ulnar nerve decompression for CuTS and
249 found that a prior history of elbow trauma was the most notable variable predicting
250 the need for revision surgery after simple decompression. [14] In light of those
251 findings, we determined that further investigation into risk factors leading to
252 revision for patients with idiopathic would provide additional information to
253 surgeons contemplating surgical options for CuTS.

254

255 In this study, we report an overall revision incidence of 3.2%, which is lower than
256 previous studies of in situ decompression. At least one potential factor for this
257 difference is the exclusion of patients with traumatic or arthritic etiology. However,
258 this is not completely unlike a previous study by Goldfarb et al, who excluded
259 patients with elbow arthritis, medial epicondylitis, and ulnar nerve subluxation, and

260 reported a revision incidence of 7%. [12] When Kroguet al implemented even less
261 stringent inclusion criteria, they reported a revision incidence of 19%. [14] Taken
262 together, these 3 studies suggest that, in the absence of both traumatic and arthritic
263 conditions, simple in situ decompression of the ulnar nerve for CuTS has a low
264 incidence of revision. A comparative overview of these studies is included in
265 [Table 4](#).

266

267 Our study also provides statistically significant evidence that younger age is a risk
268 factor for needing revision surgery in these patients. Although the clinical meaning
269 of this finding is less clear, the relationship of younger age as a pre-disposing factor
270 to complications after in situ decompression is not novel. Murata et al demonstrated
271 younger age to be predictive of increased incidence of ulnar nerve dislocation, as
272 simulated intra-operatively by placing patients' elbows in full-flexion after ulnar
273 nerve decompression. [20] They suggested that anatomical differences in the size of
274 the medial epicondyle and the shape of the ulnar groove played a role in the higher
275 nerve dislocation incidence in younger patients. All elbows in our study were
276 confirmed to have a ulnar nerve that neither subluxed or dislocated when tested
277 intra-operatively after release had been performed during the index procedure.
278 However, of the 7 cases requiring revision, 4 were noted to have a subluxating ulnar
279 nerve at the time of revision surgery. None of these 4 patients was noted to have
280 nerve instability in their latest physical examination prior to undergoing revision. It
281 remains unclear as to the mechanism by which a confirmed stable ulnar nerve

282 would later become unstable without any further intervention. In addition, we were
283 unable to account for the fact that these nerves appeared stable during examination
284 and only after surgical re-exposure were they unstable. We speculate that perhaps
285 some of the soft tissue and scarring that was released to gain exposure at the time of
286 revision surgery may have also had a tethering effect on the nerve. Regardless of the
287 means through which younger age predicts a higher revision incidence following in
288 situ decompression for treatment of CuTS, these findings suggest a consideration for
289 surgeons to discuss with younger patients seeking operative treatment for CuTS.

290

291 This study has limitations. Its retrospective nature required that we rely strictly on
292 medical records, which were not always complete and could be subject to
293 interpretation. In addition, though we only included patients who had at least 6
294 months of follow-up at our institution, there is potential for bias if any patients
295 sought care involving revision surgery elsewhere after that initial period. We sought
296 to minimize this possibility by attempting to reach patients via telephone while also
297 reviewing records for those patients to predict which, if any, would be likely to seek
298 care elsewhere. We were unable to contact over one-third of those patients with less
299 than 6 months of follow-up (see [Figure 1](#)). Furthermore, relying solely on clinical
300 documentation to speculate on this type of information is imperfect. Lastly, while
301 our specific aim was to investigate risk factors specific to idiopathic CuTS, exclusion
302 of patients with post-traumatic or arthritic etiologies may have led to us to
303 underestimate a clinically relevant revision incidence.

304

305 Despite these limitations, our results may be useful in establishing a treatment
306 algorithm for uncomplicated idiopathic CuTS. In particular, for patients confirmed
307 to have CuTS without arthritis or history of trauma to the involved elbow, our
308 findings strongly support in situ decompression as a reliable, first-line surgical
309 treatment option. The risk of revision increased somewhat in patients younger than
310 50 years of age, though the underlying mechanism of this relationship remains
311 unclear.

312

313 **REFERENCES**

- 314 1. Shuman S, Osterman AL, Bora FW. Compression neuropathies. *Semin Neurol.*
315 1987 Mar;7(1):76-87. Review.
- 316 2. Karthik K, Nanda R, Storey S, Stothard J. Severe ulnar nerve entrapment at
317 the elbow: functional outcome after minimally invasive in situ
318 decompression. *J Hand Surg Eur Vol.* 2012;37(2):115-122.
- 319 3. Bartels RHM a, Verhagen WIM, Van Der Wilt GJ, Meulstee J, Van Rossum LGM,
320 Grotenhuis J a. Prospective randomized controlled study comparing simple
321 decompression versus anterior subcutaneous transposition for idiopathic
322 neuropathy of the ulnar nerve at the elbow: Part 1. *Neurosurgery.*
323 2005;56(3):522-529.
- 324 4. Morse LP, Hons B, Mcguire DT, Ortho FC, Bain GI, A F a O. Endoscopic Ulnar
325 Nerve Release and Transposition. *Tech Hand Up Extrem Surg.* 2014;18(1):10-
326 14.
- 327 5. Zajonc H, Momeni A. Endoscopic release of the cubital tunnel. *Hand Clin.*
328 2014;30(1):55-62.
- 329 6. Adkinson JM, Chung KC. Minimal-incision in situ ulnar nerve decompression
330 at the elbow. *Hand Clin.* 2014;30(1):63-70.
- 331 7. Nakamura K, Uchiyama S, Ido Y, et al. The effect of vascular pedicle
332 preservation on blood flow and clinical outcome following ulnar nerve
333 transposition. *J Hand Surg Am.* 2014;39(2):291-302.
- 334 8. Ogata K, Manske PR, Lesker PA. The effect of surgical dissection on regional
335 blood flow to the ulnar nerve in the cubital tunnel. *Clin Orthop Relat Res.*
336 1985; 193(Mar):195-198.


- 337 9. Bartels RHM a, Termeer EH, Van Der Wilt GJ, et al. Simple decompression or
338 anterior subcutaneous transposition for ulnar neuropathy at the elbow: A
339 cost-minimization analysis - Part 2. *Neurosurgery*. 2005;56(3):531-535.
- 340 10. Brauer CA, Graham B. The surgical treatment of cubital tunnel syndrome: a
341 decision analysis. *J Hand Surg Eur Vol*. 2007;32(6):654-662.
- 342 11. Song JW, Chung KC, Prosser LA. Treatment of ulnar neuropathy at the elbow:
343 cost-utility analysis. *J Hand Surg Am*. 2012 Aug;37(8):1617-1629.e3.
- 344 12. Goldfarb C, Sutter M, Martens E, Manske P. Incidence of Re-Operation and
345 Subjective Outcome Following in Situ Decompression of the Ulnar Nerve *J*
346 *Hand Surg Eur Vol*. 2012;2009(3):379-383.
- 347 13. LeRoux P, Ensign T, Burchiel K. Surgical Decompression without
348 Transposition for Ulnar Neuropathy: Factors Determining Outcome.
349 *Neurosurgery*. 1990;27(5):709-714.
- 350 14. Krogue JD, Aleem AW, Osei D a., Goldfarb C a., Calfee RP. Predictors of
351 surgical revision after in situ decompression of the ulnar nerve. *J Shoulder*
352 *Elbow Surg*. 2015;24(4):634-639.
- 353 15. Aleem AW, Krogue JD, Calfee RP. Outcomes of Revision Surgery for Cubital
354 Tunnel Syndrome. *J Hand Surg Am*. 2014;39(11):2141-2149.
- 355 16. Charles YP, Coulet B, Rouzaud JC, Daures JP, Chammas M. Comparative
356 Clinical Outcomes of Submuscular and Subcutaneous Transposition of the
357 Ulnar Nerve for Cubital Tunnel Syndrome. *J Hand Surg Am*. 2009;34(5):866-
358 874.
- 359 17. Macadam SA., Gandhi R, Bezuhly M, Lefaivre KA. Simple Decompression
360 Versus Anterior Subcutaneous and Submuscular Transposition of the Ulnar

- 361 Nerve for Cubital Tunnel Syndrome: A Meta-Analysis. *J Hand Surg Am.*
362 2008;33(8):1-12.
- 363 18. Mitsionis GI, Manoudis GN, Paschos NK, Korompilias AV, Beris AE.
364 Comparative study of surgical treatment of ulnar nerve compression at the
365 elbow. *J Shoulder Elbow Surg.* 2010 Jun;19(4):513-9.
- 366 19. Hill SR. Cost-effectiveness analysis for clinicians. *BMC Med.* 2012;10(1):10.
- 367 20. Murata K, Omokawa S, Shimizu T, Nakanishi Y, Kawamura K, Yajima H,
368 Tanaka Y. Risk factors for dislocation of the ulnar nerve after simple
369 decompression for cubital tunnel syndrome. *Hand Surg.* 2014;19(1):13-8.
- 370

371
372
373
374


FIGURES

Figure 1. Flow chart of inclusion and exclusion criteria applied to potential study subjects.



375

376 **Figure 2.** Graphical representation of change between pre- to postoperative
 377 Modified McGowan grade. Aside from 2 patients with preoperative grade of IIa who
 378 improved to normal postoperatively (thick dashed arrow), all other patients either
 379 improved by one grade (solid arrow) or remained the same (dotted arrow).



380