

4-1-2018

Differences in work environment for staff as an explanation for variation in central line bundle compliance in intensive care units.

Yuna S.H. Lee
Yale University

Patricia W. Stone
Columbia University

Monika Pogorzelska-Maziarz
Thomas Jefferson University

Ingrid M. Nembhard
Yale University

Follow this and additional works at: <https://jdc.jefferson.edu/nursfp>



Part of the [Nursing Commons](#), and the [Public Health Commons](#)

[Let us know how access to this document benefits you](#)

Recommended Citation

Lee, Yuna S.H.; Stone, Patricia W.; Pogorzelska-Maziarz, Monika; and Nembhard, Ingrid M., "Differences in work environment for staff as an explanation for variation in central line bundle compliance in intensive care units." (2018). *College of Nursing Faculty Papers & Presentations*. Paper 97.

<https://jdc.jefferson.edu/nursfp/97>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in College of Nursing Faculty Papers & Presentations by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.



Published in final edited form as:

Health Care Manage Rev. 2018 ; 43(2): 138–147. doi:10.1097/HMR.0000000000000134.

Differences in Work Environment for Staff as an Explanation for Variation in Central Line Bundle Compliance in ICUs

Yuna S.H. Lee, MPH,

Doctoral Candidate, Yale School of Public Health, 60 College Street, P.O. Box 208034, New Haven, CT 06520-8034; Phone: 646 708 1097; Fax: 203-785-6287

Patricia W. Stone, PhD, FAAN,

Centennial Professor in Health Policy, Columbia University School of Nursing, 617 W. 168th Street, New York, NY 10032; Phone 212-305-1738 Fax 212 305-3659

Monika Pogorzelska-Maziarz, PhD, MPH, and

Assistant Professor, Jefferson School of Nursing, Thomas Jefferson University 130 South 9th Street, Room 847, Philadelphia, PA 19107; Phone: 215-503-5613; Fax: 215-503-0376

Ingrid Nembhard, PhD, MS

Associate Professor, Yale School of Public Health, Yale School of Management, 60 College Street, P.O. Box 208034, New Haven, CT 06520-8034; Phone: 203-785-3778; Fax: 203-785-6287

Abstract

Background—Central line-associated bloodstream infections (CLABSIs) are a common and costly quality problem, and their prevention is a national priority. A decade ago, researchers identified an evidence-based bundle of practices that reduce CLABSIs. Compliance with this bundle remains low in many hospitals.

Purpose—To assess whether differences in core aspects of work environments—workload, quality of relationships, and prioritization of quality—are associated with variation in maximal CLABSI bundle compliance i.e., compliance 95–100% of the time in intensive-care units (ICUs).

Methodology/Approach—Cross-sectional study of hospital medical-surgical ICUs in the United States. Data on work environment and bundle compliance were obtained from the Prevention of Nosocomial Infections & Cost Effectiveness Refined (P-NICER) Survey completed in 2011 by infection prevention directors, and data on ICU and hospital characteristics from the National Healthcare Safety Network (NHSN). Factor and multi-level regression analyses were conducted.

Findings—Reasonable workload and prioritization of quality were positively associated with maximal CLABSI bundle compliance. High-quality relationships, although a significant predictor when evaluated apart from workload and prioritization of quality, had no significant effect after accounting for these two factors.

Practice Implications—Aspects of the staff work environment are associated with maximal CLABSI bundle compliance in ICUs. Our results suggest that hospitals can foster improvement in

ensuring maximal CLABSI bundle compliance—a crucial precursor to reducing CLABSI infection rates—by establishing reasonable workloads and prioritizing quality.

Keywords

Central line-associated bloodstream infection (CLABSI); work environment; maximal CLABSI bundle compliance; implementation; infection control

Introduction

Central line-associated bloodstream infections (CLABSIs) are a common and costly quality problem (Berwick, Calkins, McCannon, & Hackbarth, 2006; Centers for Disease Control and Prevention, 2014; Institute of Medicine, 2001). In United States (U.S.) hospitals, 41,000 patients experience a CLABSI annually, with 1 in 4 affected patients dying as a result (Centers for Disease Control and Prevention, 2014; Weaver, Weeks, Pham, & Provonost, 2014). Caring for infected patients costs the healthcare system \$2 to \$3 billion annually (Warren et al., 2006). That so many patients and the healthcare system endure these health and financial costs respectively, is of great concern because CLABSIs are preventable (Berwick et al., 2006).

A decade ago, researchers identified and began disseminating information about an evidence-based bundle of practices aimed at reducing the CLABSI rate (Berwick et al., 2006; Mermel, Farr, & Sheretz, 2001; Provonost, 2008). The bundle, termed “the CLABSI bundle”, consists of five practices: appropriate hand hygiene, use of chlorhexidine for skin preparation, full-barrier precautions when inserting central venous catheters, avoidance of femoral line placement, and removal of unnecessary lines (Provonost et al., 2006). In the most extensive study to date, National On the CUSP: Stop BSI Project, which involved more than 1,100 adult ICUs across 44 states, researchers calculated that use of the bundle was associated with a 43% decrease in CLABSI rates, 1,500 saved lives, and \$175 million in reduced healthcare costs (Agency for Healthcare Research and Quality, 2012; Weaver et al., 2014). While on average the CUSP program succeeded (the median reported infection rate per 1,000 catheter days decreased to zero and was sustained for eighteen months), participants exhibited variation in the decrease of CLABSI incidence (Dixon-Woods, Bosk, Aveling, Goeschel, & Pronovost, 2011; Provonost, 2008). Researchers found maximal bundle compliance – defined as all practices in the bundle were used 95% or more of the time - is key for reduction: only when hospitals use practices in the bundle at this highest level of compliance does CLABSI incidence decrease (Furuya et al., 2011; Zachariah et al., 2014).

Despite the effectiveness of the CLABSI bundle and widespread promotion by federal agencies such as the Centers for Disease Control and Prevention (CDC) (Berwick et al., 2006), bundle compliance in the U.S. remains low. Having a written policy in place does not necessarily translate into high compliance (Zachariah et al., 2014). Maximal bundle compliance estimates range from 28% to 38% of U.S. hospitals (Provonost, 2008). Low compliance is puzzling given that the bundle’s practices are evidence-based and do not require specialized expertise or equipment for implementation (Provonost, 2008). It is also

puzzling because some hospitals achieve and maintain high compliance, indicating that it is attainable; something other than feasibility impacts compliance (Rangachari et al., 2015).

The aim of this study is to assess whether differences in work environment for staff are associated with variation in maximal CLABSI bundle compliance. Work environment is the “inner setting” of the organization where staff interplay with the organization within which they work (Damschroder et al., 2009). The study of work environment facilitates understanding of how organizations can promote staff behavior change, and how staff behavior can result in organizational variation (Lukas et al., 2007). We focus on work environment because it creates opportunities or constraints on workers’ behavior (Johns, 2006) and has been shown to influence other aspects of patient safety outcomes (Stone et al., 2007), including CLABSI rates (e.g. Dixon-Woods et al. 2011; McAlearney et al. 2015). This past work that linked work environment to CLABSI rates found that aspects of the work environment were qualitatively linked to lower CLABSI rates, including for example, leader goal setting and support (McAlearney, Hefner, Robbins, & Garman, 2013) and inter-professional collaboration (McAlearney, Hefner, Robbins, Harrison, & Garman, 2015). An ex-post analysis of the CUSP project identified that cultural influences within the organization were the likely source of variation in CLABSI rates (Dixon-Woods et al., 2011). This prompted the CUSP team to add a cultural component to their intervention, with programs focused on quality at the unit, executive and community levels (Provonost, Marsteller, & Goeschel, 2011). Hospitals that have used this intervention have seen greater reduction in their CLABSI rates (Provonost et al., 2011).

Given that CLABSI bundle compliance is a precursor to lower CLABSI rates, we anticipate that work environment might also matter to CLABSI bundle compliance. This possibility has not yet been examined, nor tested empirically. Organizational scholars assert that three core factors construct work environments, also termed work contexts: the work to be performed (task context), relationships with co-workers (social context), and the setting where work occurs (culture and physical context) (Johns, 2006). We focus on a defining element of each: staff workload (amount of work to perform – task context), quality of staff relationships (social context), and hospitals’ prioritization of quality (culture and physical context). Past qualitative work by McAlearney et al. (2015, 2016) identified these elements as similar aspects of work environment to be important for reducing CLABSI incidence (e.g. resource discussions, staff communication with each other, leader alignment). In this manuscript, we provide a quantitative complement and extension to the past work by focusing on these aspects of work environment in relation to the preceding step of CLABSI bundle compliance.

Our study responds to the call for research on bundle compliance and the role of organizational factors (Hsu, Weeks, Yang, Sawyer, & Marsteller, 2014) as well as practitioner calls for insight that begets strategies to improve compliance (Rangachari et al., 2015; Zachariah et al., 2014). Improvement has become a greater priority for hospitals because CLABSIs are no longer reimbursable in many value-based payment systems (Bodily, McMullen, Walker, & Warren, 2011; Rangachari et al., 2015).

Conceptual Framework: Why Work Environment Might Affect Compliance

Drawing on the organizations literature, there is logic for why each work environment factor—staff workload, quality of staff relationships, and hospitals' prioritization of quality (infection prevention, in particular)—may influence CLABSI bundle compliance.

Staff workload—Research shows that when workers have a reasonable workload (i.e., the amount of work they are expected to do can be accomplished in the time allotted (Nembhard & Edmondson, 2006), they are more likely to take time to perform all tasks as specified. When workload is excessive, they tend to forget steps and take shortcuts to accomplish their total work faster (Chong, Van Eerde, Chai, & Rutte, 2011). Excessive workload has been correlated with lower nurses' compliance with patient care plans (Daud-Gallotti et al., 2012; Pakyz et al., 2014) and hand hygiene mandates (Institute of Medicine, 2001). Similar to these tasks, CLABSI bundle compliance may be neglected when health professionals are overburdened. In contrast, we expect CLABSI bundle compliance to be higher when professionals have reasonable workloads, as it frees time and attention to devote to compliance.

Quality of staff relationships—Lower quality relationships are characterized by interactions that are limited to transactional exchanges (this-for-that), while higher quality relationships are characterized by trust, cooperation, and interpersonal concern (Carmeli & Gittell, 2009; Nembhard, Northrup, Shaller, & Cleary, 2012). According to social exchange theory, quality of relationships with others in the workplace is positively associated with workers' willingness to behave in ways that support colleagues and their task completion (Blau, 1986; Emerson, 1976). The higher the quality of the relationship, the more likely workers are to communicate and coordinate with each other, including about sensitive issues (Carmeli & Gittell, 2009). Comfort in communication is likely to aid CLABSI bundle compliance because it enables workers to feel safe reminding others about practices and asking for help to perform them. In high-quality relationships, neither party in the communication feels threatened by such conversation (Carmeli & Gittell, 2009). Past research has shown that staff relationships characterized by trust and collegiality were shown to facilitate the face-to-face communication needed for compliance with another type of bundle, the *Clostridium difficile* bundle (Pakyz et al., 2014). High-quality relationships between staff responsible for infection-related issues (e.g. providers, nurses or infection prevention specialists) should facilitate CLABSI bundle compliance as well, as some of the bundle's practices (e.g., removal of unnecessary lines) can involve communication and coordination between care team members.

Prioritization of quality—Organizational priorities are reflected in what the organization and its leaders supports, encourages and rewards, which in turn influences staff perception of priorities, and their behaviors (Johns, 2006). Staff perception of priorities is theorized to exert subtle control over staff by creating an environment (“organizational culture”) that sets acceptable norms and behaviors. When an organization prioritizes quality, an emphasis on quality permeates the organization's mission and action (Nembhard et al., 2012) and facilitates staff actions in support of quality (Rangachari et al. 2015). Thus, with respect to CLABSI, organizations' and their leaders' prioritization of quality, and infection prevention

specifically, is likely to affect bundle compliance because when quality is prioritized, organizations tend to take actions that support bundle compliance (e.g. placing handwashing stations in key areas), which cultivates and reinforces a culture of quality (Provonost, 2008), to which staff respond. A systematic review of healthcare associated infection (HAI) reduction strategies (e.g., ventilator-associated pneumonia interventions) – that did not include use of the CLABSI bundle - identified prioritization of safety efforts and organizational support as predictors of the use of these strategies (Mauger et al., 2014). A recent qualitative study of management practices to prevent HAIs identified top-level commitment to quality as the key management practice related to CLABSI outcomes (McAlearney et al., 2015). Thus, we hypothesize that reasonable workload, high-quality staff relationships, and prioritization of quality are each positively associated with CLABSI bundle compliance.

Methods

Sample and Data Collection

We examined the effect of staff workload, high-quality staff relationships and prioritization of quality on maximal CLABSI bundle compliance in ICUs, using data from 507 adult medical-surgical intensive care units (ICUs). These ICUs were located in hospitals that participated in the Prevention of Nosocomial Infections and Cost-Effectiveness Refined (P-NICER) study. The P-NICER survey was web-based and assessed the existence of evidence-based infection prevention policies for reducing HAIs including CLABSI, compliance with each evidence-based practice, and the work environment in U.S. hospitals. Work environment questions were drawn from the Leading a Culture of Quality-Infection Prevention Survey (LCQ-IP) (Pogorzelska-Maziarz, Nembhard, Schnall, Nelson, & Stone, 2015), an infection prevention-focused version of the LCQ survey, which has been used to assess the work environment for staff in other healthcare settings (e.g., primary care) (Nembhard et al., 2012).

As described in detail elsewhere (Furuya et al., 2011; Stone, Pogorzelska-Maziarz, Herzig, Furuya, & Dick, 2014; Zachariah et al., 2014), all nonveteran hospitals enrolled in the CDC National Healthcare Safety Network (NHSN) in 2011 (N= 3,374) were invited to participate in P-NICER. NHSN is the nation's largest HAI surveillance system (Zachariah et al., 2014). Hospitals were invited to participate in the survey via an e-mailed invitation, followed by weekly reminders and a last chance letter (a modified Dillman technique), all of which occurred between October and December 2011. The invitation explained that participation entailed survey completion, and providing the research team with access to data from the NHSN annual survey and up to six years (2006–2011) of CLABSI rate data. Each participating hospital was entered into a lottery for 10 \$100 prizes per week for 8 weeks.

Invitations were directed to the director or manager of each hospital's infection prevention (IP) and control department, who was asked to complete the survey. IP directors were selected as informants because prior research has shown infection preventionists (IPs) to be valid informants on hospital work environments and infection-related issues in ICUs (Hazamy et al., 2013). In hospitals, IPs are often the designated leader of infection control programs and have been required to report HAI rates and other data to the NHSN, a task

designated to IPs by the Centers for Disease Control and Prevention over 30 years ago, with the belief and after analysis showing that IPs are well-positioned to reliably report HAI-related performance (Bryant et al., 2016). In the course of their work, IPs work closely with other hospital staff on infection-related issues, as members of the clinical departments. They track infections in the ICU, provide consultation to ICU staff, and implement and monitor interventions and policies in the ICU – all of which makes them knowledgeable about the state of infection prevention practices such as bundle compliance. Because IPs' work requires them to spend considerable time in ICUs as “pseudo” team members, they are also able to provide an informed, evaluation of staff work environment. In using IPs as single informants in HAI research, we follow several national studies (Furuya et al., 2011; Krein et al., 2007; Saint et al., 2008) and the CUSP program, which has regarded their perspective as “providing a common unifying thread that ties activities at the front line of care to an organization's management” (pg. 571, (Goeschel, Wachter, & Provonost, 2010).

Of the 3,374 invited hospitals, 975 responded to the survey (29% overall response rate). Prior work found these hospitals did not differ significantly from those that did not respond to the survey, including on the hospital characteristics included in this study (Stone et al., 2014; Zachariah et al., 2014). In our study, we focused on the 2,768 hospitals with medical-surgical ICUs, of which 644 hospitals responded to the survey (23% response rate). Medical-surgical ICUs treat medical and surgical patients and are a high-risk setting for CLABSI due to the frequent use of catheters during critical care procedures (Provonost, 2008). They were the largest ICU type in the P-NICER sample (Stone et al., 2014), and are the most common type of ICU in U.S. hospitals (Stone et al., 2007). The 644 hospitals provided survey data on 703 ICUs. We excluded 196 of these ICUs (multiple reasons may apply): 135 because of incomplete data on the ICU and hospital characteristics used as covariates in our analyses, 75 because of incomplete bundle compliance responses, 2 because “don't know” was provided as the bundle compliance response, 17 because of incomplete work environment responses. Table 1 shows the characteristics of the 507 medical-surgical ICUs in our final sample (72% of the 703 ICUs), originating from 455 hospitals.

Measures

Maximal CLABSI Bundle Compliance—We assessed the degree of compliance with the CLABSI bundle in each ICU by aggregating survey respondents' answer to the question: *“During the last period monitored, what proportion of the time was this policy correctly implemented?”* There was a question for each policy in the bundle: “Cleaning patient's skin with chlorhexidine at an insertion site”, “Monitoring hand hygiene practices at insertion”, “Checking the line daily for necessity”, “Using maximal barrier precautions upon insertions for patients with a central venous catheter” and “Selecting an optimal catheter site” (Stone et al., 2014). Respondents indicated: no monitoring, we monitor but don't know proportion, rarely or never (<25%), sometimes (25%–74%), usually (75%–94%) and all of the time (95%–100%). “Don't know” responses (n=2) were excluded. We created a dichotomous variable to indicate whether an ICU complied all of the time with all practices, i.e. respondents selected all of the time for each practice, indicating maximal compliance that re-coded the categories (1= maximal compliance and 0=otherwise) to assess the effect of work environment on maximal (the highest level of) bundle compliance, which past work

showed has the greatest association with CLABSI reduction (Furuya et al., 2011; Zachariah et al., 2014).

Work Environment—We assessed our focal work environment factors – staff workload, high-quality staff relationships and prioritization of quality in each ICU on a 5-point Likert response scale (1=strongly disagree; 5 = strongly agree). Table 2 shows the items used to assess each work environment factor. Exploratory factor analyses were conducted to determine the discriminant validity of each measure. The appropriateness of factor analysis was supported by the Kaiser-Meyer-Olkin Measure of Sampling Adequacy test value (0.93, $p < 0.001$) that exceeded the 0.90 threshold for Bartlett’s Test of Sphericity (Hatcher & O’Rourke, 1994). We conducted an exploratory factor analysis on correlation matrices using the principal axis factor method with squared multiple correlation estimates as initial communality estimates, a Promax (oblique) rotation to account for significant correlations between factors ($r = 0.32$, $p < .05$), and item assignment to factors on which the item had a pattern loading greater than 0.40. As recommended by experts, we excluded two items because of cross-loading (Guttman, 1954). The presence of three factors/survey scales was supported by the discontinuity of the scree plot occurring at $n=3$, and eigenvalues for each factor (4.67, 1.41 and 1.02) exceeding Guttman’s threshold criteria of 1 (Guttman, 1954). Cronbach’s alpha for each measure (reported in Table 2) was above 0.70, indicating satisfactory reliability of the measures.

ICU-Level Covariates—We controlled for ICU characteristics that could be related to maximal CLABSI bundle compliance: central line utilization ratio, number of central line days, and number of ICU beds. We included ICU’s central line utilization ratio, which is a ratio of the days in which central lines were used to patient days as an indicator of patients’ illness severity (Stone et al., 2007) because sicker patients may be more susceptible to CLABSI. We included the number of ICU beds and number of central line days (i.e., how frequently central line insertion occurs) because they could impact the degree to which the ICU is familiar with central line prevention practices.

Hospital-Level Covariates—We also controlled for hospital-level characteristics that could be related to maximal bundle compliance: teaching hospital, geographic region, urbanicity of location, number of patient days, number of admitted patients and participation in HAI quality improvement initiatives. Teaching hospitals, which we considered all hospitals that were members of the Council of Teaching Hospitals, may have higher compliance because their educational efforts cause them to emphasize compliance. Geographic region (east, central, south or west) and urbanicity (urban, suburb, rural) may impact external resources available for infection prevention efforts (Furuya et al., 2011). The number of patient days and number of admitted patients indicates patient volume, which has been linked to experience implementing evidence-based processes of care (Williams, Koss, Morton, Schmaltz, & Loeb, 2008) and might impact the likelihood of maximal bundle compliance. Participation in one or more of the many quality improvement and demonstration programs aimed at reducing HAIs may also be related to bundle compliance. Hence, we included an indicator variable addressing whether the hospital participates in IHI, CUSP or other initiatives.

Statistical Analysis

We began our analysis by calculating descriptive statistics for maximal bundle compliance to gain a sense of the range of compliance in our sample. Next, we conducted bivariate Spearman correlation analyses between work environment variables and covariates. Then we used a series of multivariate multi-level regression models to estimate the associations between the work environment variables, covariates, and maximal CLABSI bundle compliance. We accounted for the 507 ICUs nested in 455 hospitals with a multi-level model using SAS PROC GENMOD, with a logit link function to account for the binary outcome variable and a cluster indicator (REPEATED command) to model the effect of ICU membership. In the first model, we included covariates only, allowing us to assess how much these structural characteristics of the ICUs and hospitals alone account for the variation in compliance. In the next three models, we assessed the association between each work environment factor and maximal CLABSI bundle compliance separately, while controlling for ICU and hospital covariates. In the final model, all work environment variables were included to assess the robustness of results and ascertain the relative importance of these factors after controlling for hospital and ICU characteristics. We examined Odds Ratios and 95% CI to evaluate the significance of each factor. All analyses were conducted in SAS version 9.3.

Findings

Maximal bundle compliance (95–100% compliance with all practices in the bundle) occurred in 17.64% of the ICUs in our sample; the remaining 82.36% of ICUs were below this benchmark.

As reported in Table 2, the average staff workload in the study ICUs was moderate (mean=3.06 on 5-point scale, S.D.=0.83), while quality of staff relationships and prioritization of quality were evaluated as high (mean=4.32, S.D.=0.68 and mean=4.00, S.D.=0.61, respectively). Correlations between these work environment factors were significant (p -values < .05), however, variance inflation factors were all less than 6 (under the standard threshold of 10), indicating that problematic multicollinearity was not present (Neter, Wassermann, & Kutner, 1989).

Table 3 presents our regression analyses results. Model 1, the covariates-only model, shows that none of the structural characteristics of the hospitals and their ICUs considered had a significant association with maximal bundle compliance (all p -values>0.2). Models 2, 3, and 4 show that the work environment for staff was associated with CLABSI bundle compliance. In sequence, the models show that reasonable workload (OR=1.67; 95% CI=1.27– 2.21; p <0.01), high-quality staff relationships (OR = 1.36; 95% CI=1.01–1.86; p =0.04), and prioritization of quality (OR = 2.27; 95% CI=1.58–3.25; p <0.001), were each positively associated with maximal CLABSI bundle compliance. When these three factors were included in the same model (Model 5), only reasonable workload and prioritization of quality were positively associated with maximal bundle compliance. In the study ICUs, for a one unit increase in reasonable workload (e.g., from agree to strongly agree), the odds of achieving maximal CLABSI bundle compliance increased by 1.34 (95% CI=1.30–1.84; p =0.03). For a one unit increase in prioritization of quality (e.g., from agree to strongly agree),

the odds of achieving maximal CLABSI bundle compliance increased by 2.04 (95% CI =1.34– 3.06; $p<0.001$). High-quality staff relationships were not significantly associated with compliance (OR=0.99, 95% CI=0.70– 1.38, $p=0.92$).

Discussion

To our knowledge, this study is the first to examine whether there is an association between the work environment for staff and maximal CLABSI bundle compliance in U.S. ICUs. Our results provide evidence of a link. Staff workload and hospitals' prioritization of quality were positively associated with maximal CLABSI bundle compliance. The significance of these across models suggests that the overall work to be performed by staff (task context), which affects time and attention to compliance, and the setting where work occurs (culture context), which influences staff appreciation for the importance of compliance, persistently determines the extent to which staff comply with the CLABSI bundle. This observation is consistent with prior work that finds that workload and prioritization play a role in compliance with other bundles (Fakih et al., 2014; Pakyz et al., 2014). Our study adds to this evidence by identifying work environment as a key contributor to use of the CLABSI bundle specifically, which has been a challenge for many hospitals. It provides evidence that complements past qualitative work that has linked the work environment to CLABSI rates (e.g. McAlearney et al., 2013; McAlearney et al., 2015).

This study provides evidence of the importance of two elements in the “inner work setting” in the implementation of evidence-based practice. Past implementation frameworks have theorized about their importance (Damschroder et al., 2009; Kitson et al., 2008). This study provides evidence in a national sample. Our study finds that the odds of maximal bundle compliance, an evidence-based practice for reducing CLABSI, can double if hospitals prioritize quality and increase by a third if they institute reasonable workload.

Our study also shows that not all work environment factors are equally influential on staff compliance with the CLABSI bundle, as the presence of high-quality staff relationships had limited relation to maximal CLABSI bundle compliance once we accounted for the other two factors. We had theorized that high-quality staff relationships influenced bundle compliance by facilitating communication and coordination. These actions are clearly important to patient safety and quality (Institute of Medicine, 2001) but may be less central for the CLABSI bundle. Some of the practices in the CLABSI bundle (e.g. hand hygiene, chlorhexidine use) are enacted independently by clinicians; communication and coordination are not required for successful completion. This contrasts with key practices in other infection prevention bundles. For example, compliance with the *Clostridium difficile* bundle requires communication between clinicians with different expertise (physicians, pharmacists, microbiologists, etc.) for practices such as preauthorization of antimicrobials, and coordination of assessments between clinicians is necessary to fulfill many of the bundles' tasks. The difference in degree of communication and coordination required may explain why high-quality relationships are strongly associated with compliance with this bundle but not the CLABSI bundle. Despite the lower impact of high-quality relationships for the CLABSI bundle, we caution against dismissing their value. Certain practices, such as the removal of unnecessary lines, may still be facilitated by effective working relationships

between clinicians, even if not deterministic of compliance. Given the contrast in findings between bundles, future work should examine how the nature of bundles affects which work factors influence compliance.

Notably, whereas the work environment factors studied had a significant association with maximal CLABSI bundle compliance, the ICU and hospital-level structural factors did not. Our analyses included ICU and hospital-level covariates found to have associations with CLABSI-infection rates (e.g. patient volume, geographic location) (Zachariah et al., 2014) as well as some not previously considered though seen as potentially important (e.g. patient severity) (Furuya et al., 2011). Our finding that none of these structural factors were significantly associated with maximal CLABSI bundle compliance suggests that broader structural factors have a weaker relationship to bundle compliance than to CLABSI rates, and that compliance may be influenced more proximately by work environment. More research is needed to understand how structural and work environment factors can be leveraged to achieve better compliance and infection rates.

As with all studies, this study has limitations. First, although based upon a national study, this analysis utilized cross-sectional data. Second, although the sample size is large, the response rate for this study is low (29%). Prior studies using this data demonstrated that respondents in our sample of medical-surgical ICUs resemble non-respondents on relevant infection-control related variables including the control variables used in our analyses (Furuya et al., 2011); nevertheless, our results may not generalize. Our study should be replicated with a larger sample and other types of ICUs. Third, compliance and work environment factors were both self-reported by a single informant, the infection preventionist (IP), which may raise the issues of validity and common-method bias, even though we did not rely on a single survey item for any of our measures, which increases the reliability of our measures (Neter et al., 1989). As noted earlier, past work has found IPs to be valid informants (Hazamy et al., 2013), especially regarding work environment and organizational issues (McAlearney & Hefner, 2014). A qualitative study of 8 organizations and 76 staff showed that the observations and themes discussed by IPs about their hospitals converge with those of frontline staff, although IPs discuss CLABSI prevention using organizational constructs such as leadership, processes and resources while frontline staff illustrate issues with patient examples (McAlearney & Hefner, 2014). Moreover, the low report of compliance suggests that our data was not subject to social desirability bias, a risk of self-report. Nevertheless, the possibility of differences in IP-staff reports and inaccurate reporting of compliance remains. We hope that future research will have access to objective data on compliance and multiple informants regarding work environment. Last, this study was limited to a few core characteristics of the work environment. There are promising avenues of research in the exploration of other work environment characteristics (e.g., unit leader role-modeling) and their impact on compliance (Hsu et al., 2014).

Practice Implications

Our results suggest that hospital leadership can foster improvement in ensuring maximal bundle compliance by establishing reasonable workloads and prioritizing quality. Excessive workload is associated with patient safety and worker stress problems in many hospitals and

ICUs (Daud-Gallotti et al., 2012). Our results show another area of quality that is affected by this factor, and provides additional impetus for pursuing reasonable workload. A number of effective strategies exist including increasing staff-to-patient ratios, reducing the length of workshifts, and reforming other scheduling practices – all of which can increase the time and attention available to staff to complete important tasks (Ernst, Jiang, Krishnamoorthy, & Sier, 2004). For those organizations with scarce financial resources that limits their ability to enact the aforementioned strategies for workload reduction, past research suggests that managers acknowledging staff's contributions in the face of excessive workload can provide some aid. Manager acknowledgement can lessen the experience of workload psychologically, making it more bearable for staff and minimizing frustration that distracts from task completion (Parker et al., 2008).

With respect to prioritizing quality, past work suggests that leader behavior is key as staff attend to their actions (Morrison & Phelps, 1999; Yukl, 1981). A key behavior is to be cognizant of the artifacts, signals, and messages propagated about the importance of quality and specifically infection control in their hospitals (Rangachari et al., 2015). Proven leadership strategies for conveying prioritization include creating strategic goals (e.g., centered on quality) with milestones, enacting related initiatives throughout the organization (e.g., zero-CLABI campaign), and communicating periodically (e.g., about infection prevention practices) (Mauger et al., 2014; Rangachari et al., 2015).

Conclusion

Ensuring maximal bundle compliance is a crucial precursor to reducing CLABSI infection rates, making the study of bundle compliance an important endeavor. We found that differences in staff workload and hospitals' prioritization of quality were central work environment factors associated with variation in maximal CLABSI bundle compliance. This work provides a basis for further research to evaluate more aspects of work environment that may contribute to variation in compliance with CLABSI and other bundles, and for interventions focused on the work environment for staff.

Acknowledgments

This study was funded by the National Institute of Nursing Research (R01NR010107). Dr. Nembhard's work on this paper was supported by a grant (#K01HS018987) from the Agency for Healthcare Research and Quality (AHRQ). The content is solely the responsibility of the authors and does not necessarily represent the official views of funding agencies. Study procedures were approved by the Human Investigations Committee at Yale University and the Columbia University Medical Center, RAND Corporation and New York University Medical Center Institutional Review Boards.

References

- Agency for Healthcare Research and Quality. AHRQ Patient Safety Project Reduces Bloodstream Infections by 40 Percent. Rockville, MD: 2012.
- Berwick DM, Calkins DR, McCannon CJ, Hackbarth AD. The 100,000 lives campaign: setting a goal and a deadline for improving health care quality. *JAMA*. 2006; 295(3):324–327. DOI: 10.1001/jama.295.3.324 [PubMed: 16418469]
- Blau, P. Exchange and power in social life. John Wiley & Sons; 1986.
- Bodily MM, McMullen K, Walker R, Warren D. The use of administrative data in detecting central line infections: Implications for hospital reimbursement. *Am J Infect Control*. 2011; 39(5):E. 137.

- Bryant K, Harris A, Gould C, Humphreys E, Lundstrom T, Murphy D, ... Zerr D. Necessary Infrastructure of Infection Prevention and Healthcare Epidemiology Programs: A Review. *Infect Control Hosp Epidemiol*. 2016; 37(4):371–380. [PubMed: 26832072]
- Carmeli A, Gittel J. High-quality relationships, psychological safety, and learning from failures in organizations. *Journal of Organizational Behavior*. 2009; 30:708–729.
- Centers for Disease Control and Prevention. Central Line-Associated Bloodstream Infection (CLABSI) Event. 2014. Retrieved from
- Chong D, Van Eerde W, Chai K, Rutte C. A double-edged sword: The effect of challenge and hindrance time pressure on new product development teams. *IEEE Transactions on Engineering Management*. 2011; 58(1):71–86.
- Damschroder LJ, Aron DC, Keith RE, Kirsh SR, Alexander JA, Lowery JC. Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science. *Implement Sci*. 2009; 4:50.doi: 10.1186/1748-5908-4-50 [PubMed: 19664226]
- Daud-Gallotti R, Costa S, Guimares T, Tadihla K, Inoue E, Vasconcelo T, ... Levin A. Nursing workload as a risk factor for healthcare associated infections: A prospective study. *PLoS One*. 2012 Dec 27.
- Dixon-Woods M, Bosk CL, Aveling EL, Goeschel CA, Pronovost PJ. Explaining Michigan: developing an ex post theory of a quality improvement program. *Milbank Q*. 2011; 89(2):167–205. DOI: 10.1111/j.1468-0009.2011.00625.x [PubMed: 21676020]
- Emerson R. Social Exchange Theory. *Annual Review of Sociology*. 1976; 2:335–362.
- Ernst AT, Jiang H, Krishnamoorthy J, Sier D. Staff scheduling and rostering: A review of applications, methods and models. *European Journal of Operational Research*. 2004; 153(1):3–27.
- Fakih MSK, Edson B, Watson S, Barries J, Saint S. Engaging health care workers to prevent catheter-associated urinary tract infection and avert patient harm. *Am J Infect Control*. 2014; 42:S223–S229. [PubMed: 25239714]
- Furuya EY, Dick A, Perencevich EN, Pogorzelska M, Goldmann D, Stone PW. Central line bundle implementation in US intensive care units and impact on bloodstream infections. *PLoS One*. 2011; 6(1):e15452.doi: 10.1371/journal.pone.0015452 [PubMed: 21267440]
- Goeschel C, Wachter RM, Pronovost P. Responsibility for quality improvement and patient safety: hospital board and medical staff leadership challenges. *Chest*. 2010; 138(1):171–178. [PubMed: 20605815]
- Guttman L. Some necessary conditions for common-factor analyses. *Psychometrika*. 1954; 19:149–161.
- Hatcher, L., O'Rourke, N. A step-by-step approach to using SAS for factor analysis and structural equation modeling. 2. 1994.
- Hazamy PA, Van Antwerpen C, Tserenpuntsag B, Haley VB, Tsivitis M, Doughty D, ... Stricof RL. Trends in validity of central line-associated bloodstream infection surveillance data, New York State, 2007–2010. *Am J Infect Control*. 2013; 41(12):1200–1204. DOI: 10.1016/j.ajic.2013.06.006 [PubMed: 24035214]
- Hsu Y, Weeks W, Yang T, Sawyer M, Marsteller J. Impact of self-reported guideline compliance: Bloodstream infection prevention in a national collaborative. *Am J Infect Control*. 2014; 42:S191–S196. [PubMed: 25239709]
- Institute of Medicine. *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington DC: New Academy Press; 2001.
- Johns G. The essential impact of context on organizational behavior. *Academy of Management Review*. 2006; 31(2):386–408.
- Kitson AL, Rycroft-Malone J, Harvey G, McCormack B, Seers K, Titchen A. Evaluating the successful implementation of evidence into practice using the PARIHS framework: theoretical and practical challenges. *Implementation Science*. 2008; 3(1):1–12. [PubMed: 18179688]
- Krein SL, Hofer TP, Kowalski CP, Olmsted R, Kauffman C, Forman J, ... Saint S. Use of central venous catheter-related bloodstream infection prevention practices by US hospitals. *Mayo Clinic Proceedings*. 2007; 82(6):672–678. [PubMed: 17550746]

- Lukas CV, Holmes SK, Cohen AB, Restuccia J, Cramer IE, Shwartz M, Charns MP. Transformational change in health care systems: an organizational model. *Health Care Manage Rev.* 2007; 32(4): 309–320. DOI: 10.1097/01.HMR.0000296785.29718.5d [PubMed: 18075440]
- Mauger B, Marbella A, Pines E, Chopra R, Black E, Aronson N. Implementing quality improvement strategies to reduce healthcare-associated infections: A systematic review. *Am J Infect Control.* 2014; 42:S274–283. [PubMed: 25239722]
- McAlearney AS, Hefner J. Facilitating central line-associated bloodstream infection prevention: A qualitative study comparing perspectives of infection control professionals and frontline staff. *Am J Infect Control.* 2014; 42(10):S216–S222. [PubMed: 25239713]
- McAlearney AS, Hefner J, Robbins J, Garman AN. The role of leadership in eliminating health care-associated infections: a qualitative study of eight hospitals. *Adv Health Care Manag.* 2013; 14:69–94. [PubMed: 24772883]
- McAlearney AS, Hefner J, Robbins J, Harrison MI, Garman AN. Preventing central line-associated bloodstream infections: a qualitative study of management practices. *Infect Control Hosp Epidemiol.* 2015; 36(5):557–563. [PubMed: 25703102]
- Mermel L, Farr B, Sheretz R. Guidelines for the management of intravascular catheter-related infections. *Infect Control Hosp Epidemiol.* 2001; 22(4):222–242. [PubMed: 11379714]
- Morrison EW, Phelps CC. Taking charge at work: Extrarole effort to initiate workplace change. *Academy of Management Journal.* 1999; 42(4):403–419.
- Nembhard IM, Edmondson AC. Making it safe: The effects of leader inclusiveness and professional status on psychological safety and improvement efforts in health care teams. *Journal of Organizational Behavior.* 2006; 27:941–966.
- Nembhard IM, Northrup V, Shaller D, Cleary PD. Improving organizational climate for quality and quality of care: Does membership in a collaborative help? *Medical care.* 2012; 50:S74–S82.
- Neter, J., Wassermann, W., Kutner, M. *Applied Linear Regression Models.* New York: Wiley; 1989.
- Pakyz A, Moczygamba L, VanderWielen L, Edmond M, Stevens M, Kuzel A. Facilitators and barriers to implementing antimicrobial stewardship strategies: Results from a qualitative study. *Am J Infect Control.* 2014:S257–263. [PubMed: 25239719]
- Parker L, Kirchner J, Bonner L, Fickel J, Ritchie M, Simons C, Yano E. Creating a quality-improvement dialogue: Utilizing knowledge from frontline staff, managers, and experts to foster health care quality improvement. *Quality Health Research, OnlineFirst.* 2008
- Pogorzelska-Maziarz M, Nembhard IM, Schnall R, Nelson S, Stone PW. Psychometric Evaluation of an Instrument for Measuring Organizational Climate for Quality: Evidence From a National Sample of Infection Preventionists. *American Journal of Medical Quality.* 2015 May 21.
- Provonost P. Interventions to decrease catheter-related bloodstream infections in the ICU: the Keystone Intensive Care Unit Project. *Am J Infect Control.* 2008; 36:171–175.
- Provonost P, Marsteller J, Goeschel C. Preventing bloodstream infections: A measurable national success story in quality improvement. *Health Affairs.* 2011; 30(4):628–634. [PubMed: 21471482]
- Provonost P, Needham D, Berenholtz S, Sinopoli D, Chu H, Cosgrove S, ... Goeschel C. An intervention to decrease catheter-related bloodstream infections in the ICU. *New England Journal of Medicine.* 2006; 355(26):2725–2732. [PubMed: 17192537]
- Rangachari P, Madaio M, Rethemeyer RK, Wagner P, Hall L, Roy S, Rissing P. Cumulative impact of periodic top-down communications on infection prevention practices and outcomes in two units. *Health Care Management Review.* 2015; 40(4):324–336. [PubMed: 25120195]
- Saint S, Kowalski CP, Kaufman SR, Hofer TP, Kauffman C, Olmsted RN, ... Krein SL. Preventing hospital-acquired urinary tract infection in the United States: a national study. *Clinical Infectious Diseases.* 2008; 46(2):243–250. [PubMed: 18171256]
- Stone PW, Mooney-Kane C, Larson E, Horan T, Glance L, Zwanziger J, AD. Nurse working conditions and patient safety outcomes. *Medical care.* 2007; 45:571–578. [PubMed: 17515785]
- Stone PW, Pogorzelska-Maziarz M, Herzig CT, Furuya EY, Dick A. State of infection prevention in US hospitals enrolled in the NHSN. *Am J Infect Control.* 2014; 42:94–99. [PubMed: 24485365]
- Warren D, Quadir WW, Hollenbeak CS, Elward AM, Cox MJ, Fraser VJ. Attributable cost of catheter-associated bloodstream infections among intensive care patients in a nonteaching hospital. *Crit Care Med.* 2006; 34(8):2084–2089. [PubMed: 16763511]

- Weaver S, Weeks K, Pham J, Provonost P. On the CUSP, stop BSI: Evaluating the relationship between central line-associated bloodstream infection rate and patient safety climate profile. *Am J Infect Control*. 2014; 42:S203–S208. [PubMed: 25239711]
- Williams S, Koss R, Morton D, Schmaltz S, Loeb J. Case volume and hospital compliance with evidence-based processes of care. *International journal for quality in health care : journal of the International Society for Quality in Health Care / ISQua*. 2008; 20(2):79–87.
- Yukl, G. *Leadership in Organizations*. Englewood Cliffs Prentice Hall; 1981.
- Zachariah P, Furuya EY, Edwards J, Dick A, Liu H, Herzig CT, ... Saiman L. Compliance with prevention practices and their association with central line-associated bloodstream infections in neonatal intensive care units. *Am J Infect Control*. 2014; 42(8):847–851. DOI: 10.1016/j.ajic.2014.04.020 [PubMed: 25087136]

Table 1

Characteristics of ICUs in Sample (N = 507)

Characteristic	Mean (S.D)/ N (%)
<i>ICU-level characteristics</i>	
Central line utilization rate	0.40 (0.21)
Number central line days/yr	1514.25 (1453.18)
Number of ICU Beds	32.13 (38.81)
<i>Hospital-level characteristics</i>	
Teaching hospital	
No	327 (65%)
Yes	180 (35%)
Geographic region	
Northeast (9 states)	102 (20%)
Midwest (12 states)	175 (35%)
South (17 states)	138 (27%)
West (11 states)	92 (18%)
Urbanicity	
Urban	121 (24%)
Rural	386 (76%)
Number of hospital beds	237.25 (209.64)
Number of admitted patients	12,124.64 (10,615.51)
Infection preventionists per bed	1.13 (1.13)

Summary statistics are reported as means with standard deviations in parentheses for continuous variables and sample sizes with percentages in parentheses for categorical variable.

Table 2
Measures for Work Environment with Item Loadings from Exploratory Factor Analysis (N=507)

Measures			
Survey Items	Staff Workload	High-quality Staff Relationships	Prioritization of Quality (Infection Prevention)
The quality of work suffers because of the amount of work staff are expected to do (Reverse-scored)	0.70	0.26	0.32
Most people in this organization are so busy that they have little time to devote to infection prevention efforts (Reverse-scored)	0.71	0.24	0.39
I observe a high level of cooperation among all members of my work unit or department	0.32	0.95	0.39
There is a climate of trust in my department or work unit	0.29	0.84	0.40
Results of our infection prevention efforts are measured and communicated regularly to staff	0.34	0.35	0.65
HAI prevention goals and strategic plan of our organization are clear and well communicated	0.34	0.35	0.65
Employees are encouraged to become involved in infection prevention	0.39	0.20	0.60
There is good information flow among departments to provide high quality patient safety and care	0.49	0.29	0.69
People here feel a sense of urgency about preventing HAI	0.39	0.29	0.72
Cronbach's Alpha	0.75	0.88	0.83
Mean (S.D.)	3.06 (0.83)	4.32 (0.68)	4.00 (0.61)

Table 3
Associations of Maximal CLABSI Bundle Compliance and Work Environment in United States ICUs in 2011 (N=507)

Independent variables	Model 1: Covariates Only	Model 2: Reasonable Workload	Model 3: High-Quality Staff Relationships	Model 4: Prioritization of Quality	Model 5: All Work Environment Factors
	Odds Ratio (95% CI)	Odds Ratio (95% CI)	Odds Ratio (95% CI)	Odds Ratio (95% CI)	Odds Ratio (95% CI)
Reasonable workload		1.67 (1.27, 2.21)**			1.34 (1.30, 1.84)*
High-quality staff relationships			1.36 (1.01, 1.86)*		0.99 (0.70, 1.38)
Prioritization of quality				2.27 (1.58, 3.25)**	2.04 (1.34, 3.06)**
ICU and hospital covariates					
Central line utilization rate	0.73 (0.15, 2.46)	0.74 (0.14, 2.01)	0.79 (0.15, 2.04)	0.80 (0.14, 2.20)	0.82 (0.15, 2.36)
Number of central line days per year	1.00 (0.99, 1.03)	1.00 (0.98, 1.02)	1.00 (0.99, 1.03)	1.01 (0.99, 1.03)	1.00 (0.99, 1.03)
Number of ICU beds	0.99 (0.98, 1.01)	1.00 (0.99, 1.02)	0.99 (0.98, 1.01)	1.00 (0.99, 1.02)	0.98 (0.97, 1.00)
Teaching hospital	0.98 (0.97, 1.00)	0.99 (0.98, 1.01)	0.99 (0.98, 1.01)	0.99 (0.98, 1.01)	0.99 (0.98, 1.01)
Region: Northeast	0.40 (0.30, 1.51)	0.39 (0.29, 1.49)	0.40 (0.30, 1.50)	0.43 (0.31, 1.52)	0.44 (0.35, 1.58)
Region: Midwest	0.22 (0.02, 1.60)	0.18 (0.01, 1.18)	0.20 (0.02, 1.57)	0.20 (0.02, 1.57)	0.23 (0.02, 1.79)
Region: South	0.18 (0.15, 1.90)	0.17 (0.14, 1.80)	0.18 (0.15, 1.91)	0.20 (0.16, 2.02)	0.21 (0.18, 2.06)
Region: West	0.30 (0.02, 1.40)	0.27 (0.02, 1.37)	0.35 (0.25, 1.42)	0.40 (0.31, 1.43)	0.41 (0.32, 1.46)
Rural setting	1.17 (0.86, 1.53)	1.15 (0.84, 1.50)	1.10 (0.80, 1.45)	1.11 (0.81, 1.47)	1.15 (0.84, 1.49)
Urban setting	1.03 (0.57, 1.88)	1.09 (0.62, 1.93)	1.05 (0.59, 1.90)	1.05 (0.59, 1.91)	1.02 (0.56, 1.85)
Number of beds	1.00 (0.99, 1.01)	1.00 (0.99, 1.01)	1.00 (0.99, 1.01)	1.00 (0.99, 1.01)	1.00 (0.99, 1.01)
Number of admitted patients	1.00 (0.99, 1.01)	1.00 (0.99, 1.01)	1.00 (0.99, 1.01)	1.00 (0.99, 1.01)	1.00 (0.99, 1.01)
Participation in IHI or CUSP	1.20 (0.96, 1.41)	1.17 (0.88, 1.33)	1.16 (0.90, 1.30)	1.16 (0.92, 1.32)	1.13 (0.87, 1.27)

* $P < 0.05$;

** $P < 0.01$