

10-2018

Synthetic DNA immunotherapy in biochemically relapsed prostate cancer

N. Shore
Carolina Urologic Research Center

E. I. Heath
Karmanos Cancer Center

L. T. Nordquist
Urology Cancer Center and GU Research Network

H. Cheng
University of Washington School of Medicine
Follow this and additional works at: <https://jdc.jefferson.edu/medoncfp>

K. Bhatt
Part of the Oncology Commons
in vivo Pharmaceuticals Inc.

[Let us know how access to this document benefits you](#)

See next page for additional authors
Recommended Citation

Shore, N.; Heath, E. I.; Nordquist, L. T.; Cheng, H.; Bhatt, K.; Morrow, M.; McMullan, T.; Kraynyak, K.; Lee, J.; Sacchetta, B.; Liu, L.; Rosencranz, S.; Tagawa, S. T.; Appleman, L. J.; Tutrone, R.; Garcia, J.; Whang, Y.; Kelly, W.; Csiki, I.; and Bagarazzi, M. L., "Synthetic DNA immunotherapy in biochemically relapsed prostate cancer" (2018). *Department of Medical Oncology Faculty Papers*. Paper 114.

<https://jdc.jefferson.edu/medoncfp/114>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Medical Oncology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

N. Shore, E. I. Heath, L. T. Nordquist, H. Cheng, K. Bhatt, M. Morrow, T. McMullan, K. Kraynyak, J. Lee, B. Sacchetta, L. Liu, S. Rosencranz, S. T. Tagawa, L. J. Appleman, R. Tutrone, J. Garcia, Y. Whang, W. Kelly, I. Csiki, and M. L. Bagarazzi

830P Synthetic DNA immunotherapy in biochemically relapsed prostate cancer

N. Shore¹, E.I. Heath², L.T. Nordquist³, H. Cheng⁴, K. Bhatt⁵, M. Morrow⁶, T. McMullan⁷, K. Krainyak⁶, J. Lee⁸, B. Sacchetta⁸, L. Liu⁶, S. Rosencranz⁹, S.T. Tagawa⁹, L.J. Appleman¹⁰, R. Tutrone¹¹, J. Garcia¹², Y. Whang¹³, W. Kelly¹⁴, I. Csiki⁸, M.L. Bagarazzi⁸

¹Urology, Carolina Urologic Research Center, Myrtle Beach, SC, USA, ²Oncology, Karmanos Cancer Center, Detroit, MI, USA, ³Medical Oncology, Urology Cancer Center and GU Research Network, Omaha, NE, USA, ⁴Medical Oncology, University of Washington School of Medicine, Seattle, WA, USA, ⁵Clinical, Inovio Pharmaceuticals Inc, Plymouth Meeting, PA, USA, ⁶Immunology, Inovio Pharmaceuticals, Inc., Plymouth Meeting, PA, USA, ⁷Biostatistics, Inovio Pharmaceuticals, Inc., Plymouth Meeting, PA, USA, ⁸Clinical, Inovio Pharmaceuticals, Inc., Plymouth Meeting, PA, USA, ⁹Medical Oncology, Weill Cornell Medical College, New York, NY, USA, ¹⁰Medical Oncology, University of Pittsburgh Cancer Institute, Pittsburgh, PA, USA, ¹¹Medical Oncology, Chesapeake Urology Associates, Towson, MD, USA, ¹²Medical Oncology, Cleveland Clinic, Cleveland, OH, USA, ¹³Medical Oncology, Lineberger Comprehensive Cancer Center, Chapel Hill, NC, USA, ¹⁴Medicine, Thomas Jefferson University, Philadelphia, PA, USA

Background: INO-5150 (PSA and PSMA) +/- INO-9012 (IL-12), a synthetic DNA immunotherapy, was assessed for safety, immunogenicity and efficacy in biochemically recurrent prostate cancer patients (pts).

Methods: Phase I, open-label, multi-center study in the US included pts with rising PSA after surgery and/or RT, PSA doubling time (PSADT) >3 months (mos), testosterone >150 ng/dL and no concurrent ADT. Safety, immunogenicity and efficacy (PSA kinetics, PFS) were evaluated in 4 treatment arms of 15 pts each. Arms A: 2mg INO-5150, B: 8.5 mg INO-5150, C: 2mg INO-5150 + 1mg INO-9012 and D: 8.5mg INO-5150 + 1mg INO-9012. Pts received 4 IM doses of vaccine followed by electroporation on day 0, wks 3, 12 and 24 and were followed for 72 wks.

Results: 50/61 (82%) pts completed all visits and treatments were well tolerated with no safety concerns. Median PFS for overall population [N = 61, baseline (D0) PSADT range (mos) 1.5-217.1, median 9.8] and for a subset of pts with D0 PSADT ≤12mos (N = 36) has not yet been reached (FU 3-19 mos). 86% of pts with D0 PSADT ≤12 mos were progression free through 19mos FU. 27 out of 36 (75%) pts with D0 PSADT ≤ 12 mos had disease stabilization at wks 27 evidenced by significant improvement in log₂PSA change over time (slope) and PSADT from D0 (Slope=0.19 declined to 0.1, PSADT=5.3 improved to 10.1 mos, p = <0.0001). This effect was maintained at wk 72 (Slope=0.09, PSADT=10.6, p = <0.0001). Immunogenicity was observed in 77% (47/61) of pts by multiple immunologic assessments. Patient immunogenicity to INO-5150 as determined by CD38 and Perforin + CD8 T cell immune reactivity correlated with attenuated % PSA rise compared to pts without reactivity (p = 0.05, n = 50).

Conclusions: INO-5150 +/- INO-9012 was safe, well tolerated and immunogenic. Clinical efficacy was observed in the patients with D0 PSADT ≤ 12 mos as evidenced by a significant dampening of log₂PSA change over time and increased PSADT up to 72 weeks FU. Additional genomic analyses are ongoing to further elucidate the correlation of immunologic efficacy and clinical benefit. (NCT02514213).

Clinical trial identification: FDA IND number 15870.

Legal entity responsible for the study: Inovio Pharmaceuticals, Inc.

Funding: Inovio Pharmaceuticals, Inc.

Disclosure: N. Shore: Consulting or advisory role for multiple pharmaceutical companies. E.I. Heath: Consulting or advisory role and honoraria or funding received for multiple pharmaceutical companies. H. Cheng, Y. Whang: Funding received from multiple pharmaceutical companies. K. Bhatt, M. Morrow, T. McMullan, K. Krainyak, J. Lee, B. Sacchetta, L. Liu, S. Rosencranz, I. Csiki, M.L. Bagarazzi: Employee of the pharmaceutical company sponsoring this research trial. S.T. Tagawa, R. Tutrone, J. Garcia, W. Kelly: Consulting or advisory role and funding received from multiple pharmaceutical companies. All other authors have declared no conflicts of interest.